

Dissolved Oxygen Patterns Across NCOS Sites

Spring 2026

Phoebe Dupa, Rebecca Martinez, and Shucan Zhao

Table of contents

Introduction	1
Methods	2
Results	4
Discussion	4
Conclusion	4
References	4

Introduction

North Campus Open Space (NCOS) is a restored wetland connected to the Devereux Slough system in Goleta, California. The site includes several channels, bridges, and wetland areas, so water quality may differ across the system. The North Campus Open Space Restoration Project Monitoring Report provides background on restoration progress and monitoring at the site (Rickard 2024). At a restored wetland like NCOS, water quality monitoring helps show how conditions vary across different parts of the system.

Dissolved oxygen is a useful water quality measurement because it shows how much oxygen is available in the water for aquatic organisms. In shallow wetlands and estuaries, dissolved oxygen can vary by location, season, time of day, and depth in the water column. These patterns can be influenced by water movement, temperature, salinity, vegetation, photosynthesis, respiration, and decomposition (Gale et al. 2006, Nezlin et al. 2009). These oxygen conditions are also tied to habitat quality. Low dissolved oxygen can reduce the suitability of wetland and marsh habitats for fish and other nekton (Smith and Able 2003, Clark et al. 2025). Low oxygen

near the bottom of shallow wetlands can also affect decomposition and where invertebrates are active in the water column (Lee et al. 2017). For these reasons, dissolved oxygen can be used as an indicator of water quality and aquatic habitat conditions.

Research from other wetland and estuary systems helps explain why both site and depth are important to examine. In the Carmel River Lagoon, a Central California bar-built estuary, dissolved oxygen varied across different branches of the lagoon as mouth state and hydrologic conditions changed (Dawson et al. 2023). That pattern is relevant to NCOS because East Channel, Phelps Bridge, and Venoco Bridge are different locations within the same coastal wetland system. Other studies show that shallow lagoons and estuaries can develop vertical differences in dissolved oxygen when mixing is limited. Under these conditions, bottom water can become lower in oxygen while surface water remains more oxygenated (Gale et al. 2006, Nezlin et al. 2009). Together, these studies provide a basis for looking at both site-level differences and vertical differences in dissolved oxygen at NCOS.

The analysis focuses on three NCOS sites: East Channel, Phelps Bridge, and Venoco Bridge. These sites represent different parts of the wetland and may differ in water movement, depth, vegetation, salinity, and connection to other parts of the slough. The analysis also uses elevation code to examine possible dissolved oxygen stratification within each site. Elevation code 0 represents bottom measurements, elevation code 1 represents middle measurements, and elevation code 2 represents upper measurements.

This study asks two main questions:

1. How does dissolved oxygen differ across East Channel, Phelps Bridge, and Venoco Bridge?
2. How does dissolved oxygen stratification differ across elevation-code measurement positions within each site?

Methods

This analysis used NCOS water quality monitoring data from the 2024 and 2025 water years. The analysis focused on three main sites: East Channel, Phelps Bridge, and Venoco Bridge. These sites were selected because they represent different parts of the wetland system and had enough dissolved oxygen measurements to compare. The North Campus Open Space Restoration Project Monitoring Report provides background on the restoration project and monitoring at the site (Rickard 2024).

The analysis used NCOS water quality monitoring metadata and YSI water quality measurements. The metadata included survey information such as site name, monitoring date, weather notes, water year, and location information. The YSI dataset included water quality measurements such as dissolved oxygen, temperature, salinity, conductivity, and elevation code. NOAA weather data were also available and provided temporal context for the monitoring dates.

The metadata and YSI water quality data were cleaned and joined so that each dissolved oxygen measurement was connected to the correct site, date, water year, and elevation-code position. The final dissolved oxygen dataset was filtered to include East Channel, Phelps Bridge, and Venoco Bridge during the 2024 and 2025 water years. The dataset was also filtered to elevation codes 0, 1, and 2. Elevation code 0 represented bottom measurements, elevation code 1 represented middle measurements, and elevation code 2 represented upper measurements.

Dissolved oxygen, measured in milligrams per liter, was the main response variable. Site was used to compare dissolved oxygen across the three NCOS locations, and elevation code was used to compare dissolved oxygen across bottom, middle, and upper measurement positions within each site. Descriptive statistics were calculated before statistical testing. Dissolved oxygen was summarized by site and by site/elevation-code combinations using mean, median, first quartile, third quartile, standard deviation, variance, and sample size. Stratification difference was also calculated for each site as upper median dissolved oxygen minus bottom median dissolved oxygen. Positive values indicated that median dissolved oxygen was higher at the upper measurement position than at the bottom measurement position.

Figures were created to show dissolved oxygen across sites, dissolved oxygen across elevation codes within each site, median dissolved oxygen elevation profiles, and median dissolved oxygen through time by elevation code and site. Kruskal-Wallis tests were used to compare dissolved oxygen across groups. For the first question, a Kruskal-Wallis test compared dissolved oxygen across East Channel, Phelps Bridge, and Venoco Bridge. When that result was significant, Dunn post-hoc tests were used to compare site pairs. For the stratification question, separate Kruskal-Wallis tests were run within each site to compare dissolved oxygen across elevation codes. When a within-site result was significant, Dunn post-hoc tests were used to compare elevation-code pairs. Effect sizes were also calculated to describe the strength of the patterns.

The dataset included repeated monitoring observations across sampling dates rather than continuous dissolved oxygen sensor measurements. As a result, the analysis shows patterns across sites, elevation codes, and sampling dates, but it does not capture all short-term changes that may happen within a day.

Results

Discussion

Conclusion

References

- Clark, A., W. T. Ellis, A. R. Rodriguez, and R. Baker. 2025. [Dissolved Oxygen May Limit the Suitability of Salt Marsh as Nekton Habitat](#). *Estuaries and Coasts* 48.
- Dawson, J., M. M. Orescanin, R. Clark, and K. O'Connor. 2023. [Spatiotemporal variability of dissolved oxygen in response to morphological state in a central California coast bar-built estuary](#). *Estuarine, Coastal and Shelf Science* 282:108241.
- Gale, E., C. Pattiaratchi, and R. Ranasinghe. 2006. [Vertical mixing processes in Intermittently Closed and Open Lakes and Lagoons, and the dissolved oxygen response](#). *Estuarine, Coastal and Shelf Science* 69:205–216.
- Lee, G. H. van der, M. H. S. Kraak, R. C. M. Verdonschot, J. A. Vonk, and P. F. M. Verdonschot. 2017. [Oxygen drives benthic-pelagic decomposition pathways in shallow wetlands](#). *Scientific Reports* 7.
- Nezlin, N. P., K. Kamer, J. Hyde, and E. D. Stein. 2009. [Dissolved oxygen dynamics in a eutrophic estuary, Upper Newport Bay, California](#). *Estuarine, Coastal and Shelf Science* 82:139–151.
- Rickard, A. 2024. [North campus open space restoration project monitoring report: Year 7, december 2024](#). UC Santa Barbara: Cheadle Center for Biodiversity and Ecological Restoration.
- Smith, K. J., and K. W. Able. 2003. [Dissolved Oxygen Dynamics in Salt Marsh Pools and Its Potential Impacts on Fish Assemblages](#). *Marine Ecology Progress Series* 258:223–232.