

Visualization Draft

Phoebe Dupa, Rebecca Martinez, and Shucan Zhao

2026-05-22

1. Set up

Packages

Data

2. Clean data

Cleaning metadata

Goal

Clean the metadata dataset so it includes survey date, site information, water year, and readable site names.

Cleaning water parameter data

Goal

Clean the water quality parameter dataset so it includes elevation code, elevation labels, and dissolved oxygen measurements.

3. Join data frames

Water quality = metadata + water parameters

4. Filter data for dissolved oxygen analysis

Goal

Create a filtered dissolved oxygen dataset that focuses on the three main NCOS sites, the 2024 and 2025 water years, and the main elevation codes used for the stratification question.

5. Summarize dissolved oxygen

These summaries answer the two main project questions:

1. How does dissolved oxygen differ across sites?
2. How does dissolved oxygen stratification differ across sites?

Dissolved oxygen by site

Table 1: Dissolved oxygen summary by site

Site	Mean DO	Median DO	SD	Variance	n
East Channel	7.22	7.28	3.43	11.76	109
Phelps Bridge	4.30	3.70	2.51	6.33	166
Venoco Bridge	5.68	5.98	4.01	16.09	257

Dissolved oxygen by site and elevation code

Table 2: Dissolved oxygen summary by site and elevation code

Site	Elevation code	Mean DO	Median DO	SD	n
East Channel	0	7.02	6.66	3.62	54
East Channel	1	7.53	7.96	3.20	46
East Channel	2	6.86	7.17	3.64	9
Phelps Bridge	0	4.53	4.14	2.34	76
Phelps Bridge	1	4.19	3.63	2.61	81


Site	Elevation code	Mean DO	Median DO	SD	n
Phelps Bridge	2	3.31	2.01	3.08	9
Venoco Bridge	0	4.01	4.31	3.40	89
Venoco Bridge	1	6.00	6.38	4.16	99
Venoco Bridge	2	7.38	7.62	3.72	69

6. Visualizing

These figures focus on dissolved oxygen. Figure 1 follows the original proposed site comparison. Figure 2 follows the original proposed elevation-code stratification figure. Figure 3 replaces the scrapped temperature figure with a highlighted DO stratification figure. Figure 4 adds an exploratory time component by showing dissolved oxygen through time by site and elevation code.

Figure 1: Dissolved oxygen across sites


Caption: Figure 1 shows dissolved oxygen measurements across the three main NCOS sites. Points represent individual observations, and red diamonds represent site medians.



Interpretation: Dissolved oxygen varied by site. East Channel had the highest median DO, Phelps Bridge had the lowest, and Venoco Bridge had the widest range of values.

Figure 2: Dissolved oxygen stratification by site


Caption: Figure 2 shows dissolved oxygen measurements across elevation-code positions within each site. Boxplots show the distribution of DO values, points represent individual observations, and red diamonds represent median DO values.



Interpretation: DO patterns across elevation codes were different by site. East Channel was fairly similar across depths, Phelps Bridge stayed lower overall, and Venoco Bridge showed the clearest increase from bottom to upper measurements.

Figure 3: Dissolved oxygen stratification by focal site


Caption: Figure 3 highlights dissolved oxygen measurements across elevation-code positions for each focal site. In each panel, white points show the focal site and gray points show the other sites for comparison. Elevation code is shown on the y-axis, with bottom measurements at the bottom and upper measurements at the top.



Interpretation: Stratification is most visible at Venoco Bridge, where DO is generally lower at the bottom and higher in the middle and upper elevations.

Figure 4: Dissolved oxygen through time by elevation code

Caption: Figure 4 shows dissolved oxygen measurements through time, separated by elevation code. Points represent individual dissolved oxygen observations, and colors represent sites.



Interpretation: DO changed through time, but the site pattern was still visible. East Channel was generally higher, Phelps Bridge was generally lower, and Venoco Bridge showed more separation between bottom, middle, and upper measurements.

7. Statistical analysis

The statistical analyses follow the class workflow for comparing more than two groups: summarize the data, visualize the pattern, run a Kruskal-Wallis test, use Dunn post-hoc tests when the Kruskal-Wallis result is significant, calculate effect size, and interpret the result. The main response variable is dissolved oxygen. The grouping variables are site and elevation code.

Question 1: How does dissolved oxygen differ across sites?

This analysis tests whether dissolved oxygen differs across East Channel, Phelps Bridge, and Venoco Bridge.

Kruskal-Wallis test

```
Kruskal-Wallis rank sum test
```

```
data: do_mg_l by site_full
```

```
Kruskal-Wallis chi-squared = 45.667, df = 2, p-value = 1.212e-10
```

The Kruskal-Wallis test showed that dissolved oxygen differed significantly across sites, $\chi^2 = 45.67$, $df = 2$, $p < 0.001$. This means that DO was not distributed the same way across East Channel, Phelps Bridge, and Venoco Bridge.

Dunn post-hoc test

```
# A tibble: 3 x 9
```

.y.	group1	group2	n1	n2	statistic	p	p.adj	p.adj.signif
* <chr>	<chr>	<chr>	<int>	<int>	<dbl>	<dbl>	<dbl>	<chr>
1 do_mg_l	East Chan~	Phelp~	109	166	-6.74	1.57e-11	4.72e-11	****
2 do_mg_l	East Chan~	Venoc~	109	257	-4.03	5.62e- 5	1.12e- 4	***
3 do_mg_l	Phelps Br~	Venoc~	166	257	3.72	1.98e- 4	1.98e- 4	***

Because the Kruskal-Wallis test was significant, Dunn post-hoc tests were used to compare site pairs. All three site pairs were significantly different after adjustment: East Channel and Phelps Bridge, East Channel and Venoco Bridge, and Phelps Bridge and Venoco Bridge.

Effect size

```
# A tibble: 1 x 5
```

.y.	n	effsize	method	magnitude
* <chr>	<int>	<dbl>	<chr>	<ord>
1 do_mg_l	532	0.0825	eta2[H]	moderate

The effect size for the site comparison was $\eta^2[H] = 0.083$, which was interpreted as moderate in the test output. This suggests that site explained a moderate amount of variation in DO ranks in this filtered dataset.

Interpretation

Dissolved oxygen differed across the three main NCOS sites. East Channel had the highest median DO, Phelps Bridge had the lowest median DO, and Venoco Bridge had the widest spread of values. The Dunn post-hoc test showed that all site pairs were significantly different, so DO patterns were not the same across sites.

Question 2: How does dissolved oxygen stratification differ across sites?

This analysis tests whether dissolved oxygen differs across elevation-code measurement positions within each site. This addresses stratification because elevation code represents the measurement position in the water column.

Kruskal-Wallis tests within each site

```
Kruskal-Wallis rank sum test
```

```
data: do_mg_l by elevation_code
```

```
Kruskal-Wallis chi-squared = 1.1119, df = 2, p-value = 0.5735
```

```
Kruskal-Wallis rank sum test
```

```
data: do_mg_l by elevation_code
```

```
Kruskal-Wallis chi-squared = 3.0201, df = 2, p-value = 0.2209
```

```
Kruskal-Wallis rank sum test
```

```
data: do_mg_l by elevation_code
```

```
Kruskal-Wallis chi-squared = 28.641, df = 2, p-value = 6.034e-07
```

The within-site Kruskal-Wallis tests showed no significant DO differences across elevation codes at East Channel or Phelps Bridge. Venoco Bridge did show a significant difference across elevation codes, $\chi^2 = 28.64$, $df = 2$, $p < 0.001$. This means that Venoco Bridge had the clearest statistical evidence of DO stratification.

Dunn post-hoc test for Venoco Bridge

```
# A tibble: 3 x 9
  .y.    group1 group2    n1    n2 statistic      p      p.adj p.adj.signif
* <chr>  <chr>  <chr>  <int> <int>    <dbl>    <dbl>    <dbl> <chr>
1 do_mg_l 0      1      89    99      3.39 0.000688    1.38e-3 **
2 do_mg_l 0      2      89    69      5.26 0.000000145 4.36e-7 ****
3 do_mg_l 1      2      99    69      2.22 0.0266     2.66e-2 *
```

Because Venoco Bridge had a significant Kruskal-Wallis result, Dunn post-hoc tests were used to compare elevation-code pairs at that site. All three pairs were significantly different after adjustment: bottom and middle, bottom and upper, and middle and upper. This supports the pattern of higher median DO from bottom to upper measurements at Venoco Bridge.

Effect size for Venoco Bridge

```
# A tibble: 1 x 5
  .y.      n effsize method magnitude
* <chr>  <int>  <dbl> <chr>    <ord>
1 do_mg_l 257  0.105 eta2[H] moderate
```

The effect size for the Venoco Bridge elevation-code comparison was $\eta^2[H] = 0.105$, which was interpreted as moderate in the test output. This suggests that elevation code explained a moderate amount of variation in DO ranks at Venoco Bridge.

Interpretation

Statistical evidence for DO stratification was clearest at Venoco Bridge. East Channel and Phelps Bridge did not show significant DO differences across elevation codes, while Venoco Bridge did. At Venoco Bridge, median DO was lower at the bottom and higher at the middle and upper positions.

8. Overall interpretation

Overall, dissolved oxygen differed across the three main NCOS sites, and statistical evidence for stratification was clearest at Venoco Bridge. East Channel had the highest median DO, Phelps Bridge had the lowest median DO, and Venoco Bridge showed the clearest increase from bottom to upper measurements. These results suggest that DO conditions and stratification patterns were not the same across the three sites.