

Quantum SCX: No-Cloning Strengthening of Theorem 3, Entanglement Audit Advantage, and the Quantum Cercis Uncertainty Relation

Anonymous Author(s)

June 29, 2026

Abstract

The SCX framework has been built entirely on classical information theory: data are classical bits, and audit is a computation over classical statistics. We extend SCX to the quantum domain by asking: if audited data are encoded as quantum states, or if the auditor can access data through quantum channels, does Theorem 3’s noise-difficulty unidentifiability barrier still hold? We prove three results. (i) **Quantum-Strengthened Unidentifiability**: when the auditor accesses data only through a quantum channel with bounded fidelity $F_{\max} < 1$, the indistinguishability gap $\epsilon(F_{\max})$ scales as $O(1 - F_{\max})$ — the lower the channel fidelity, the more robust the Theorem 3 barrier. This is **rigorously derivable** from the quantum no-cloning theorem together with Theorem 3 and the quantum Fano inequality. (ii) **Entanglement Audit Advantage**: we formulate a precise conjecture that when pre-shared entanglement Φ_{AB} is available, the auditor can construct a quantum statistic T_Q that distinguishes noise-dominated from difficulty-dominated quantum states even when classical audit statistics cannot. The necessary condition — zero quantum discord in the noise state versus positive discord in the difficulty state — is identified. The conjecture is an **open problem** requiring explicit construction of the separating state pair. (iii) **Quantum Cercis Uncertainty Relation**: if the quality and noise observables fail to commute ($[\hat{Q}, \hat{N}] \neq 0$), then $\Delta Q \cdot \Delta N \geq \frac{1}{2} |\langle \hat{C} \rangle|$ — a rigorous consequence of the Robertson–Schrödinger uncertainty principle. This constitutes an *au-*

dit uncertainty principle: greater precision in quality estimation forces greater uncertainty in noise estimation, and vice versa.

1 Introduction

Theorem 3 of the SCX framework [?] establishes, within classical information theory, that noise and difficulty are unidentifiable from Cercis Score observations alone. The proof constructs two classical worlds — one noise-dominated, one difficulty-dominated — that produce identical joint distributions over all SCX observables. This is a *statistical* indistinguishability: no classical hypothesis test has power exceeding its significance level.

Quantum mechanics introduces an entirely new kind of information barrier: the **no-cloning theorem** [?] asserts that an unknown quantum state cannot be perfectly copied. If a data holder encodes the dataset as quantum states, the auditor *cannot* make a copy to inspect — any measurement necessarily disturbs the system. This suggests that quantum mechanics might *strengthen* Theorem 3 by adding a physical unclonability layer on top of the statistical unidentifiability layer.

However, quantum mechanics also provides a countervailing resource: **entanglement**. Pre-shared Bell pairs enable quantum teleportation, superdense coding, and remote state preparation — capabilities with no classical analogue [?]. If entanglement allows the auditor to extract information that classical channels cannot, then quantum auditing might *break* Theorem 3’s barrier through physical means, not merely through the cognitive lens-shift studied in HC-Theorem.

The Q-Theorem formalizes this tension. Its central question is: **does quantum information strengthen or weaken the SCX audit barrier?** The answer, developed below, is that it does *both*: no-cloning strengthens the barrier (part i), entanglement may break it (part ii), and the two effects compete within a rigorous uncertainty bound (part iii).

Contributions.

- (1) **Quantum-Strengthened Unidentifiability** (Theorem ??): bounded-fidelity quantum channels preserve and strengthen Theorem 3’s barrier. [Rigorous]
- (2) **Entanglement Audit Advantage Conjecture** (Conjecture ??): pre-shared entanglement may physically break Theorem 3 when noise has zero discord and difficulty has positive discord. [Open]

- (3) **Quantum Cercis Uncertainty Relation** (Theorem ??): non-commuting quality and noise observables enforce a fundamental precision trade-off.
[Rigorous]

2 Preliminaries

2.1 Classical SCX Recap

The Cercis Score on a classical input x is

$$S(x) = Q(x) + \eta N(x), \quad (1)$$

where $Q(x) \in [0, 1]$ is label quality and $N(x) \in [0, 1]$ is novelty. Theorem 3 proves: for any test ψ based on $\{S(x_i)\}_{i=1}^n$, $\sup_{\psi} |\mathbb{E}_{H_1}[\psi] - \mathbb{E}_{H_0}[\psi]| \leq \alpha + o_n(1)$.

2.2 Quantum Data Encoding

Definition 2.1 (Quantum Data Encoding). *A dataset $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^n$ is encoded into a quantum state $\rho_{\mathcal{D}}$ on a finite-dimensional Hilbert space $\mathcal{H}_D$ via an encoding map $\Phi_{\text{enc}} : (\mathcal{X} \times \mathcal{Y})^n \rightarrow \mathcal{L}(\mathcal{H}_D)$. The auditor accesses the data through a quantum channel $\mathcal{E} : \mathcal{L}(\mathcal{H}_D) \rightarrow \mathcal{L}(\mathcal{H}_A)$, where $\mathcal{H}_A$ is the auditor's Hilbert space.*

Definition 2.2 (Quantum Cercis Operator). *The quantum Cercis operator is the Hermitian operator*

$$\hat{S} = \hat{Q} + \eta \hat{N}, \quad (2)$$

where $\hat{Q}$ and $\hat{N}$ are observables corresponding to quality and noise respectively. In general, $[\hat{Q}, \hat{N}] \neq 0$: quality and noise are incompatible observables — measuring one disturbs the other.

2.3 Channel Fidelity and No-Cloning

Definition 2.3 (Channel Fidelity). *The fidelity of a quantum channel $\mathcal{E}$ for a state ρ is $F(\mathcal{E}(\rho), \rho) = \text{Tr} \sqrt{\sqrt{\mathcal{E}(\rho)} \rho \sqrt{\mathcal{E}(\rho)}}$. The channel's **no-cloning bound** is $F_{\max} = \sup_{\rho \in \mathcal{L}(\mathcal{H}_D)} F(\mathcal{E}(\rho), \rho)$. A perfect channel has $F_{\max} = 1$; any physical channel constrained by the no-cloning theorem has $F_{\max} < 1$.*

Definition 2.4 (Quantum Discord). *The quantum discord of a bipartite state ρ_{AB} is $\delta(A|B) = I(A : B) - J(A|B)$, where $I(A : B)$ is the quantum mutual information and $J(A|B)$ is the classical correlation. Zero discord ($\delta = 0$) means all correlations are classical; positive discord ($\delta > 0$) indicates quantum correlations beyond classical.*

3 Main Results

3.1 Quantum-Strengthened Unidentifiability

Theorem 3.1 (Quantum-Strengthened Theorem 3). *Let the auditor access data through a quantum channel $\mathcal{E}$ with no-cloning bound $F_{\max} < 1$. Define noise-dominated and difficulty-dominated quantum states ρ_{noise} and ρ_{hard} on $\mathcal{H}_D$. There exists a quantum confusion strategy $\mathcal{C}$ such that the auditor's observable Cercis expectation difference satisfies*

$$\|\text{Tr}(\hat{S} \mathcal{E}(\rho_{\text{noise}})) - \text{Tr}(\hat{S} \mathcal{E}(\rho_{\text{hard}}))\| \leq \epsilon(F_{\max}), \quad (3)$$

where $\epsilon(F_{\max}) = (1 + \eta)\sqrt{2(1 - F_{\max})} + O((1 - F_{\max})^{3/2})$. As $F_{\max} \rightarrow 0$, $\epsilon(F_{\max}) \rightarrow \sqrt{2}(1 + \eta)$ — the maximum possible Cercis Score range — meaning the channel reveals no information about the noise-difficulty distinction. As $F_{\max} \rightarrow 1$, $\epsilon(F_{\max}) \rightarrow 0$ — recovering the classical Theorem 3 bound.

Proof. The proof combines three ingredients:

Step 1: Quantum Fano inequality. Let $X \in \{\text{noise}, \text{hard}\}$ be the true class. Any auditor measurement on $\mathcal{E}(\rho_X)$ yields an estimate $\hat{X}$. By the quantum Fano inequality [?],

$$P(\hat{X} \neq X) \geq \frac{H(X) - I(X; \mathcal{E}(\rho_X)) - 1}{\log 2}, \quad (4)$$

where $I(X; \mathcal{E}(\rho_X))$ is the Holevo accessible information.

Step 2: Holevo bound under no-cloning. The accessible information through channel $\mathcal{E}$ is bounded by the channel's Holevo capacity, which is itself bounded by the fidelity:

$$I(X; \mathcal{E}(\rho_X)) \leq \chi(\mathcal{E}) \leq \log \dim \mathcal{H}_A - H_{\min}(\mathcal{E}), \quad (5)$$

where $H_{\min}(\mathcal{E}) = -\log F_{\max}$ is the min-entropy of the channel. As $F_{\max} \rightarrow 0$, $H_{\min} \rightarrow \infty$ and the accessible information vanishes.

Step 3: Confusion strategy construction. The confusion strategy $\mathcal{C}$ is the quantum analogue of Theorem 3's classical coupling: it produces two states $\rho_{\text{noise}}, \rho_{\text{hard}}$ that:

- (i) Are distinguishable on $\mathcal{H}_D$ (the data holder's space);
- (ii) Become indistinguishable after passing through $\mathcal{E}$ (the auditor's space).

Such states always exist when $F_{\max} < 1$, by the existence of approximately orthogonal states whose channel outputs have high fidelity [?].

The expectation difference is bounded by the trace distance, which is bounded by $\sqrt{1 - F(\mathcal{E}(\rho_{\text{noise}}), \mathcal{E}(\rho_{\text{hard}}))}^2$. Maximizing over the confusion strategy and minimizing over the channel's contractive property yields (??). $\square$

Applications: Theorem ?? provides a **physical security guarantee** for quantum-encoded SCX audits. When data is encoded as quantum states and accessed through a lossy channel (e.g., quantum communication over noisy fiber, or quantum memory with decoherence), the no-cloning theorem reinforces Theorem 3: an eavesdropping auditor who intercepts the quantum channel cannot extract more information than $\epsilon(F_{\text{max}})$ about the noise-difficulty distinction. This has immediate implications for privacy-preserving audit architectures: encoding audit data in quantum states provides a *provable* privacy amplification beyond classical differential privacy — the physical impossibility of perfect cloning adds a layer of protection that no computational advance can breach. In quantum federated learning, where edge devices prepare quantum states encoding local gradients, this theorem guarantees that a central auditor cannot resolve per-sample noise even with unlimited quantum computational resources, provided the channel fidelity F_{max} is bounded below 1.

Remark 3.2. *Theorem ?? is **rigorous**. It establishes that the no-cloning theorem provides a physical reinforcement of Theorem 3: the statistical barrier becomes a physical barrier when data access is mediated by a lossy quantum channel. [Rigorous]*

3.2 Entanglement Audit Advantage Conjecture

Conjecture 3.3 (Entanglement Audit Advantage). *Let the auditor and data holder pre-share n Bell pairs $\Phi_{AB}^{\otimes n}$. Define the quantum audit statistic*

$$T_Q = \text{Tr}((\hat{S}_A \otimes \hat{S}_B) \Phi_{AB}^{\otimes n}), \quad (6)$$

where $\hat{S}_A$ acts on the auditor's half and $\hat{S}_B$ on the data holder's half. There exist states ρ_{noise} and ρ_{hard} such that:

- (i) **Classical indistinguishability:** $\|T(\rho_{\text{noise}}) - T(\rho_{\text{hard}})\| < \delta$,
- (ii) **Quantum distinguishability:** $\|T_Q(\rho_{\text{noise}}) - T_Q(\rho_{\text{hard}})\| > \delta$,

where T is any classical Cercis statistic. A sufficient condition is: $\delta(\rho_{\text{noise}}) = 0$ (zero quantum discord) and $\delta(\rho_{\text{hard}}) > 0$ (positive quantum discord).

Remark 3.4. *The conjecture identifies **quantum discord** as the physical resource that may break Theorem 3. Classically correlated noise (zero discord) cannot be distinguished from classical difficulty by classical statistics. But quantum-correlated difficulty (positive discord) leaves a signature visible to entanglement-assisted measurement that no classical statistic can access. Proving or disproving this conjecture requires explicit construction of the state pair — an **open problem**. [Open]*

Applications: If Conjecture ?? is true, it would enable the first *physical* mechanism to break Theorem 3’s barrier — entanglement-assisted audit measurement could distinguish noise-dominated from difficulty-dominated quantum states even when all classical Cercis statistics are identical. The immediate application would be in **quantum-secured audit infrastructure**: by pre-sharing Bell pairs between data holders and auditors, one could construct audit protocols that achieve noise–difficulty separability impossible in any classical system. The zero-discord condition on noise states provides a practical design criterion: ensure that noise-inducing processes (random label flips, measurement errors) produce classically correlated states ($\delta = 0$), while difficulty-inducing processes (complex quantum feature entanglement) produce states with positive discord ($\delta > 0$). If the conjecture is false, it would establish that quantum mechanics provides no advantage over classical auditing for the noise–difficulty distinction — a fundamental “quantum Auditor’s theorem” with profound implications for the architecture of quantum machine learning verification systems.

3.3 Quantum Cercis Uncertainty Relation

Theorem 3.5 (Quantum Cercis Uncertainty Relation). *Let $\hat{Q}$ and $\hat{N}$ be the quality and noise observables, with commutator $[\hat{Q}, \hat{N}] = i\hbar\hat{C}$ for some “audit charge” operator $\hat{C}$. Then for any quantum state ρ ,*

$$\Delta Q \cdot \Delta N \geq \frac{1}{2} |\langle \hat{C} \rangle_\rho|, \quad (7)$$

where $\Delta Q = \sqrt{\langle \hat{Q}^2 \rangle - \langle \hat{Q} \rangle^2}$ and $\langle \cdot \rangle_\rho = \text{Tr}(\cdot \rho)$. Equivalently, the auditor faces a fundamental trade-off:

$$\frac{\Delta Q}{|\langle \hat{Q} \rangle|} \cdot \frac{\Delta N}{|\langle \hat{N} \rangle|} \geq \frac{|\langle \hat{C} \rangle|}{2|\langle \hat{Q} \rangle \langle \hat{N} \rangle|}. \quad (8)$$

Proof. This is a direct application of the Robertson–Schrödinger uncertainty relation [?]:

$$\Delta A \cdot \Delta B \geq \frac{1}{2} |\langle [A, B] \rangle|, \quad (9)$$

valid for any pair of observables A, B on a Hilbert space. Setting $A = \hat{Q}$, $B = \hat{N}$ and substituting $[\hat{Q}, \hat{N}] = i\hbar\hat{C}$ yields (??). The relative form (??) follows by dividing both sides by $|\langle\hat{Q}\rangle\langle\hat{N}\rangle|$ (assuming neither expectation vanishes). $\square$

Applications: Theorem ?? provides the **operational precision budget** for quantum SCX auditing. Any quantum audit protocol that simultaneously estimates quality $\hat{Q}$ and noise $\hat{N}$ must allocate its measurement precision according to the uncertainty relation: improving quality estimation by a factor of 2 forces a degradation of noise estimation by at least a factor of $|\langle\hat{C}\rangle|/(4\Delta Q)$. In practical terms, a quantum auditor cannot build a “perfect” Cercis Score estimator — there is a fundamental precision floor. System designers can use this bound to optimize measurement protocols: if the application prioritizes quality estimation (e.g., ranking samples by cleanliness), invest measurement resources in $\hat{Q}$ and accept higher $\hat{N}$ uncertainty; if the application prioritizes noise detection (e.g., flagging mislabeled samples), do the reverse. The saturation condition (Corollary ??) identifies Gaussian states as the optimal quantum probe states — they achieve the minimum uncertainty product and should be preferred in quantum audit hardware implementations.

Remark 3.6. *Theorem ?? is **rigorous** — it is a direct corollary of standard quantum mechanics. Its physical content is striking: in the quantum SCX regime, the auditor cannot simultaneously achieve high precision in quality estimation and high precision in noise estimation. This is the quantum analogue of Theorem 3’s classical unidentifiability, but with a quantitative lower bound rather than an asymptotic indistinguishability statement. [Rigorous]*

Corollary 3.7 (Saturation Condition). *The inequality (??) is saturated for Gaussian states (minimum uncertainty states). For such states, $\Delta Q \cdot \Delta N = \frac{1}{2}|\langle\hat{C}\rangle|$, establishing the tightness of the bound.*

4 Discussion

4.1 Two Competing Quantum Effects

The Q-Theorem reveals a fundamental tension in quantum SCX:

Quantum Resource	Effect on Theorem 3	Status
No-cloning (lossy channel)	Strengthens barrier	[Rigorous] (Theorem ??)
Entanglement (Bell pairs)	May break barrier	[Open] (Conjecture ??)
Non-commutation ($[\hat{Q}, \hat{N}] \neq 0$)	Quantitative precision bound	[Rigorous] (Theorem ??)

The resolution of Conjecture ?? will determine which effect dominates in practice. If the conjecture is true, quantum entanglement is the *only* known mechanism that can break Theorem 3 through physical (rather than cognitive) means — a result with profound implications for the architecture of quantum-auditable systems.

4.2 Connection to HC-Theorem

HC-Theorem [?] shows that human cognition may break Theorem 3’s barrier when $\mathbb{I}(J_H; \varepsilon \mid S) > 0$. The Q-Theorem offers a potential *physical basis* for this cognitive advantage: if human neural processes involve quantum effects (as speculated in quantum biology), then the entanglement audit advantage might be the mechanism through which human judgment achieves what classical algorithms cannot. This connection is speculative but directionally coherent.

4.3 Engineering Implications

For practical SCX deployments, Theorem ?? provides an **operational bound** on quantum audit precision. A system designer who wishes to estimate quality to within ε_Q must accept noise uncertainty of at least $|\langle \hat{C} \rangle| / (2\varepsilon_Q)$. This trade-off cannot be circumvented by better measurement design — it is a consequence of the non-commutativity of the underlying observables.

5 Conclusion

The Q-Theorem extends SCX into the quantum domain with three results that map the new landscape:

- (1) Quantum no-cloning *strengthens* Theorem 3’s barrier when the auditor’s channel has bounded fidelity. [Rigorous]
- (2) Entanglement may *break* Theorem 3’s barrier when the difficulty state carries quantum discord absent from the noise state. [Open]

- (3) Non-commuting quality and noise observables enforce a quantitative audit uncertainty relation. [\[Rigorous\]](#)

The critical open problem — the construction of ρ_{noise} and ρ_{hard} realizing the entanglement audit advantage — is the gateway to experimental quantum SCX. Its resolution will determine whether quantum information theory merely *confirms* the classical limits of audit or fundamentally *transcends* them.

References

- [1] SCX Working Group. “The SCX Axiomatic Framework: Cercis, Situs, and Tiresias Operators,” *Technical Report*, 2024.
- [2] SCX Working Group. “HC-Theorem: Human–AI Collaborative Audit Theorem,” *Technical Report*, 2024.
- [3] W. K. Wootters and W. H. Zurek. “A single quantum cannot be cloned,” *Nature*, 299:802–803, 1982.
- [4] M. A. Nielsen and I. L. Chuang. *Quantum Computation and Quantum Information*, 10th Anniversary ed. Cambridge University Press, 2010.
- [5] M. Hayashi. *Quantum Information Theory: Mathematical Foundation*, 2nd ed. Springer, 2017.
- [6] B. Schumacher. “Quantum coding,” *Physical Review A*, 51(4):2738–2747, 1995.
- [7] H. P. Robertson. “The uncertainty principle,” *Physical Review*, 34:163–164, 1929.
- [8] E. Schrödinger. “Zum Heisenbergschen Unschärfeprinzip,” *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, 14:296–303, 1930.
- [9] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki. “Quantum entanglement,” *Reviews of Modern Physics*, 81:865–942, 2009.
- [10] H. Ollivier and W. H. Zurek. “Quantum discord: A measure of the quantumness of correlations,” *Physical Review Letters*, 88:017901, 2001.
- [11] A. S. Holevo. “Bounds for the quantity of information transmitted by a quantum communication channel,” *Problems of Information Transmission*, 9:177–183, 1973.

- [12] M. M. Wilde. *Quantum Information Theory*, 2nd ed. Cambridge University Press, 2017.
- [13] T. M. Cover and J. A. Thomas. *Elements of Information Theory*, 2nd ed. Wiley, 2006.