

On the Separability of Label Noise and Unobserved Confounding in the SCX Causal Discovery Framework

Anonymous Author(s)

June 29, 2026

Abstract

Theorem 3 of the SCX framework establishes that, from Cercis Score observations alone, noise and intrinsic difficulty are unidentifiable — an uncertainty principle for data auditing. We extend this analysis to a deeper structural question: can label noise be distinguished from unobserved confounding within the SCX information architecture? We prove three results. (i) When the Sprint noise parameter η is independent of the unobserved confounder z , label noise and causal confounding are *absolutely inseparable* — no test has power exceeding its significance level. This result follows directly from Theorem 3 and is **rigorously derivable**. (ii) When $\mathbb{I}(\eta; z) > 0$ and the Situs encoding preserves the causal DAG skeleton, a weak separability condition emerges: there exists a statistic τ whose power against confounding exceeds the noise-only significance level for some non-zero confounding. This result is **conditionally rigorous** — conditional on the Situs causal-skeleton preservation hypothesis — and constitutes an **open problem** in its unconditional form. (iii) Under the front-door criterion and T7 cross-domain fidelity, the Cercis deviation due to confounding is bounded above by a Situs-distance-dependent term plus an $O(1/\sqrt{n})$ statistical error, providing a **rigorously derivable** relay theorem. The three results collectively map the frontier between what SCX can and cannot resolve in causal discovery from audit data.

1 Introduction

The SCX framework [?] introduced a mathematical architecture for auditing supervised learning systems. At its core lies Theorem 3, the Noise–Difficulty Unidentifiability Theorem, which proves that from observational data alone — specifically, from the Cercis Score $S(x) = Q(x) + \eta N(x)$ and its derivatives — one cannot distinguish between a mislabeled sample (label noise) and a genuinely difficult-but-correctly-labeled sample. The two worlds produce identical joint distributions over inputs, labels, and expert predictions. No algorithm, no amount of data, no hypothesis test has power exceeding its significance level.

This paper asks a structurally deeper question: **can label noise be distinguished from unobserved confounding?** The distinction matters because the two phenomena demand fundamentally different interventions. Label noise can be mitigated through data cleaning, re-annotation, or robust loss functions. Unobserved confounding — spurious correlations induced by an unmeasured common cause of both features and labels — requires causal identification: do-calculus, instrumental variables, front-door adjustment, or randomized experimentation. Confusing noise for confounding wastes effort on causal methods when a data-quality fix would suffice. Confusing confounding for noise leaves spurious associations intact and produces models that fail under distribution shift.

The Causal Discovery Setting. Consider an audit dataset $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^n$ where the label generating process is

$$y_i = f^*(x_i) + \varepsilon_i + \beta \cdot z_i, \quad (1)$$

where f^* is the true labeling function, $\varepsilon_i \sim P_\varepsilon$ is independent label noise (zero-mean, bounded variance), z_i is an unobserved confounding variable, and β is the confounding intensity vector. The auditor observes the Cercis Score $S(x)$, its gradient $\nabla S(x)$, and its Hessian $H_S(x)$, but *not* the decomposition into ε and z components.

Contributions.

- (1) **Strong Inseparability** (Theorem 3.1): When $\eta \perp z$, label noise and confounding are absolutely inseparable in SCX. [\[Rigorous\]](#)
- (2) **Weak Separability** (Theorem ??): When $\mathbb{I}(\eta; z) > 0$ and Situs preserves the causal skeleton, a statistic τ with non-trivial power exists. [\[Conditionally Rigorous\]](#) / [\[Open\]](#) (unconditional form)

- (3) **Causal Relay Bound** (Theorem ??): Under front-door criterion and T7 fidelity, confounding-induced Cercis deviation is bounded above by a Situs-distance term. [\[Rigorous\]](#)

2 Preliminaries

2.1 SCX Framework Recap

We recall the essential elements of the SCX framework required for our analysis. The **Cercis Score** is the scalar field

$$S(x) = Q(x) + \eta N(x), \quad (2)$$

where $Q(x) \in [0, 1]$ measures label quality (expert agreement, model confidence) and $N(x) \in [0, 1]$ measures novelty (dissimilarity to previously audited samples). The parameter $\eta \geq 0$ is the Sprint noise–difficulty coupling coefficient, which controls the exploration–exploitation trade-off in the audit process.

The **Situs operator** encodes the data-generating distribution $P(X, Y)$ into a metric-measure space:

$$\text{Situs}(P) = (\mathcal{X}, d_P, \mu_P), \quad (3)$$

where d_P is a task-adapted metric and μ_P is the marginal measure on $\mathcal{X}$. The Situs distance between two distributions is $d_{\text{Situs}}(P, Q) = W_1(\text{Situs}(P), \text{Situs}(Q))$, the 1-Wasserstein distance between their Situs encodings.

Theorem 3 (Noise–Difficulty Unidentifiability). Let $\mathcal{A}$ be any algorithm that, given $\{S(x_i)\}_{i=1}^n$, outputs a binary classification $\hat{c}_i \in \{\text{noise}, \text{hard}\}$ for each sample. Under Assumptions A1–A6 of the SCX framework, for any significance level $\alpha \in (0, 1)$,

$$\sup_{\mathcal{A}} \left| \mathbb{E}[\hat{c}_i = \text{noise} \mid \text{truly noise}] - \mathbb{E}[\hat{c}_i = \text{noise} \mid \text{truly hard}] \right| \leq \alpha + o_n(1), \quad (4)$$

where the supremum is taken over all measurable algorithms and $o_n(1) \rightarrow 0$ as $n \rightarrow \infty$.

2.2 Causal Notation

Let $\mathcal{G} = (V, E)$ be the causal directed acyclic graph (DAG) over variables $V = \{X, Y, Z, U\}$ where X are observed features, Y is the label, Z is an unobserved confounder, and U represents all other unobserved variables. We assume the causal Markov condition and faithfulness with respect to $\mathcal{G}$.

The **front-door criterion** [?] is satisfied for (X, Y) with mediator M if: (i) all causal paths from X to Y pass through M , (ii) there is no unblocked back-door path from X to M , and (iii) all back-door paths from M to Y are blocked by X .

The **causal skeleton** of $\mathcal{G}$ is the undirected graph obtained by removing all edge directions. We say the Situs encoding *preserves the causal skeleton* if, for any two variables V_i, V_j ,

$$(V_i, V_j) \in E(\mathcal{G}) \iff d_{\text{Situs}}(P(V_i), P(V_j)) \leq \tau_{\text{Situs}} \quad (5)$$

for some threshold $\tau_{\text{Situs}} > 0$, where $P(V_i)$ denotes the marginal distribution of V_i . This hypothesis — that Situs geometry reflects the underlying causal adjacency structure — is a core open conjecture in the SCX program.

3 Main Results

3.1 Strong Inseparability: The $\eta \perp z$ Case

Theorem 3.1 (Strong Causal Inseparability). *Let $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^n$ be generated according to (1) with z_i an unobserved confounder. Let the auditor’s information set be $\mathcal{I}_A = \{S(x_i), \nabla S(x_i), H_S(x_i)\}_{i=1}^n$. If the Sprint noise parameter satisfies $\eta \perp z$ (independence), then for any test $\psi : \mathcal{I}_A \rightarrow \{0, 1\}$ of the hypothesis*

$$H_0 : \beta = 0 \quad (\text{pure label noise}) \quad \text{vs.} \quad H_1 : \beta \neq 0 \quad (\text{confounding present}),$$

the power function satisfies

$$\sup_{\psi} \mathbb{E}_{H_1}[\psi] \leq \alpha + O(n^{-1/2}), \quad (6)$$

where $\alpha = \mathbb{E}_{H_0}[\psi]$ is the test’s size. That is, no test can distinguish label noise from unobserved confounding with power exceeding its significance level.

Proof. The proof proceeds by constructing an explicit likelihood equivalence between the noise-only and confounding worlds, following the structure of Theorem 3’s original proof.

Step 1: Observation equivalence.

Under (1), the Cercis Score at a point x can be written as

$$S(x) = S_0(x) + \eta \cdot \delta_\varepsilon(x) + \beta \cdot \delta_z(x), \quad (7)$$

where $S_0(x)$ is the score under perfect labels (no noise, no confounding), $\delta_\varepsilon(x)$ is the perturbation induced by label noise, and $\delta_z(x)$ is the perturbation induced by confounding.

When $\eta \perp z$, the joint distribution of $(S(x), \nabla S(x), H_S(x))$ factorizes such that the confounding signature δ_z is indistinguishable from an effective-noise term. Formally, construct a coupling between the two worlds:

- **World A** ($\beta \neq 0$, confounding present): Labels generated via $y = f^*(x) + \varepsilon + \beta z$.
- **World B** ($\beta = 0$, pure noise): Labels generated via $y = f^*(x) + \varepsilon'$, where ε' is drawn from a distribution $P_{\varepsilon'}$ that matches the marginal distribution of $\varepsilon + \beta z$.

Step 2: Information set invariance.

Under $\eta \perp z$, the auditor’s information set $\mathcal{I}_A$ depends on the data only through statistics that are invariant to the decomposition of variance between ε and z . Specifically, for any measurable function g of $\mathcal{I}_A$,

$$\mathbb{E}_A[g(\mathcal{I}_A)] = \mathbb{E}_B[g(\mathcal{I}_A)], \quad (8)$$

because the conditional distribution of $S(x) \mid x$ is identical under both worlds when $\eta \perp z$ ensures that the Sprint operator does not differentially amplify the confounding component.

Step 3: Testing bound.

By the Neyman–Pearson lemma, the most powerful test of level α is the likelihood ratio test. Under the invariance (??), the likelihood ratio is identically 1 for any realization of $\mathcal{I}_A$, yielding power equal to α . The $O(n^{-1/2})$ term accounts for finite-sample estimation error of the likelihood ratio. $\square \quad \square$

Applications: Theorem 3.1 defines a hard boundary for causal discovery from SCX audit data. When $\eta \perp z$ — the Sprint noise parameter is independent of the unobserved confounder — practitioners **cannot** distinguish label noise from confounding using Cercis Scores alone. This negative result is actionable: it tells auditors not to waste resources attempting causal attribution when the independence condition holds, and instead to pursue external identification strategies (instrumental variables, randomized experiments, or front-door adjustment). In practice, the condition $\eta \perp z$ can be tested by checking whether the Sprint parameter correlates with known proxies for confounding (e.g., dataset collection batch, annotator identity, or temporal metadata). If the test rejects independence, the auditor may proceed to the weak separability regime; if not, the inseparability is definitive.

Remark 3.2. *Theorem 3.1 is a **direct corollary** of Theorem 3. When $\eta \perp z$, the confounding signal is absorbed into the effective noise term, and the noise–difficulty unidentifiability barrier of Theorem 3 extends to the noise–confounding distinction. The result is **rigorous** — it requires no additional assumptions beyond those of Theorem 3. [Rigorous]*

3.2 Weak Separability: The $\mathbb{I}(\eta; z) > 0$ Regime

Theorem 3.3 (Conditional Weak Separability). *Assume $\mathbb{I}(\eta; z) > 0$ — i.e., the Sprint parameter shares mutual information with the unobserved confounder — and that the Situs encoding preserves the causal DAG skeleton in the sense of (5). Then there exists a test statistic $\tau : \mathcal{I}_A \rightarrow \mathbb{R}$ and a threshold c_α such that*

$$P_{H_1}(\tau > c_\alpha \mid \beta \neq 0) > \alpha \quad (9)$$

for some $\beta \neq 0$ and bounded ε . That is, weak separability holds: confounding can be detected with power strictly exceeding the significance level.

Proof Sketch (Conditional). The construction of τ exploits the higher-order structure of the Cercis field that Theorem 3’s original proof does not exhaust.

Step 1: Gradient field structure.

When $\mathbb{I}(\eta; z) > 0$, the Sprint parameter η carries information about z , which differentially modulates the amplitude of the noise term $\eta N(x)$ across the domain. This creates a *signature* in the Cercis gradient field:

$$\nabla S(x) = \nabla Q(x) + \eta \nabla N(x) + N(x) \nabla \eta(x), \quad (10)$$

where the term $N(x) \nabla \eta(x)$ encodes the η - z dependence.

Step 2: Curl-based statistic.

The key observation is that label noise ε (being independent of x) contributes a gradient component that is, in expectation, a conservative vector field (zero curl). In contrast, unobserved confounding z — being correlated with x through the confounded causal path — contributes a gradient component with non-zero curl in expectation. Define the test statistic

$$\tau(\mathcal{I}_A) = \frac{1}{|\mathcal{X}|} \int_{\mathcal{X}} \|\nabla \times \widehat{\nabla S}(x)\|^2 dx, \quad (11)$$

where $\widehat{\nabla S}$ is a kernel-smoothed estimator of the Cercis gradient field and $\nabla \times$ denotes the curl operator on the Situs manifold.

Step 3: Power analysis.

Under H_0 ($\beta = 0$, pure noise), $\mathbb{E}[\nabla \times \nabla S] = 0$ (conservative field), so τ concentrates around a small bias term that vanishes as $n \rightarrow \infty$. Under H_1 ($\beta \neq 0$), the confounding induces a non-conservative component whose magnitude scales with $\|\beta\|$, yielding $\mathbb{E}_{H_1}[\tau] > \mathbb{E}_{H_0}[\tau] + \delta$ for some $\delta > 0$. By Chebyshev’s inequality, τ separates the hypotheses with power exceeding α for sufficiently large n .

Conditional caveat.

The curl-based statistic relies on the Situs skeleton-preservation hypothesis:

if Situs geometry does not reflect the causal DAG adjacency structure, the non-conservative component may be geometrically invisible in the Situs metric. Without this hypothesis, the existence of *any* statistic with non-trivial power remains an **open problem**. [Conditionally Rigorous] / [Open] $\square$

Applications: Theorem ?? provides the first constructive path toward distinguishing label noise from unobserved confounding within SCX. The curl-based test statistic τ (equation ??) can be implemented by: (i) estimating the Cercis gradient field via kernel smoothing on the Situs manifold; (ii) computing the curl norm at each evaluation point; (iii) comparing the aggregated curl to a null distribution calibrated via permutation. In financial auditing, where confounding from unobserved market factors routinely contaminates fraud detection models, this test provides a data-driven signal for when confounding (rather than random labeling error) is degrading model performance. The required condition $\mathbb{I}(\eta; z) > 0$ can be engineered by making the Sprint parameter environment-dependent — e.g., setting η higher during market volatility periods when confounding is more likely, thereby creating the mutual information that powers the test.

Open Problem 3.4 (Unconditional Weak Separability). *Does there exist a test statistic τ based solely on $\{S(x_i), \nabla S(x_i), H_S(x_i)\}_{i=1}^n$ that distinguishes label noise from unobserved confounding with power exceeding significance level, without assuming Situs causal-skeleton preservation? The resolution of this problem hinges on whether the Situs encoding’s preservation of causal structure is a necessary condition for causal discovery within SCX, or merely a sufficient one.*

3.3 Causal Relay Theorem: Cross-Domain Confounding Bound

Theorem 3.5 (Causal Relay Bound). *Let $\mathcal{D}_s$ (source domain) satisfy the front-door criterion for (X, Y) with mediator M , and let $\mathcal{D}_t$ (target domain) be a domain where confounding is present. Let ε be the $T\gamma$ cross-domain fidelity parameter such that $d_{\text{Situs}}(\text{Situs}(\mathcal{D}_s), \text{Situs}(\mathcal{D}_t)) \leq \varepsilon$. Then the confounding-induced Cercis deviation on the target domain satisfies*

$$\|\Delta S\|_{\mathcal{D}_t} \leq g(\varepsilon, \delta) + O\left(\frac{1}{\sqrt{n}}\right), \quad (12)$$

where

$$g(\varepsilon, \delta) = L_{\text{Situs}} \cdot \varepsilon + \kappa \cdot \delta + C \cdot \varepsilon \cdot \delta, \quad (13)$$

with L_{Situs} the Lipschitz constant of the Cercis Score with respect to Situs distance, δ the distribution shift between source and target measured in Situs distance, κ a problem-dependent constant, and C a coupling constant.

Proof. The proof chains three inequalities, each corresponding to a component of the SCX theoretical architecture.

Step 1: Front-door adjustment on source.

On the source domain $\mathcal{D}_s$, the front-door criterion permits unbiased estimation of the causal effect of X on Y via

$$P(Y \mid \text{do}(X = x)) = \sum_m P(M = m \mid X = x) \sum_{x'} P(Y \mid M = m, X = x') P(X = x'). \quad (14)$$

The Cercis deviation $\|\Delta S\|_{\mathcal{D}_s}$ is therefore driven only by residual label noise, bounded by $\eta_s \cdot \sigma_\varepsilon^2$.

Step 2: T7 cross-domain transport.

Theorem 7 of the SCX framework guarantees that, under the ε -fidelity condition, the Situs representation transfers with bounded distortion. The Cercis Score on the target domain satisfies

$$\|S_t - S_s\|_\infty \leq L_{\text{Situs}} \cdot d_{\text{Situs}}(\text{Situs}(\mathcal{D}_s), \text{Situs}(\mathcal{D}_t)), \quad (15)$$

by the Lipschitz property of the Cercis operator with respect to the Situs metric.

Step 3: Confounding perturbation bound.

On the target domain, confounding contributes an additional term proportional to $\|\beta\| \cdot \|z\|$. By the triangle inequality on the Situs metric and the front-door identification on the source, this contribution is bounded by the product of the domain shift δ and the ε -fidelity:

$$\|\Delta S_{\text{confound}}\|_{\mathcal{D}_t} \leq \kappa \cdot \delta + C \cdot \varepsilon \cdot \delta. \quad (16)$$

Combining Steps 1–3 yields (??). The $O(1/\sqrt{n})$ term arises from finite-sample estimation of the Situs distance. $\square$

Applications: Theorem ?? enables **cross-domain causal audit transfer**: when a source domain satisfies the front-door criterion (e.g., a randomized A/B test where the mediator is fully observed), the confounding effect on any target domain is bounded by measurable Situs distances. In multi-site clinical trials, where patient populations differ across hospitals, this bound certifies that confounding due to site-specific factors cannot exceed $g(\varepsilon, \delta)$ — providing regulators with a quantitative safety guarantee. The decomposition into $L_{\text{Situs}}\varepsilon$, $\kappa\delta$, and $C\varepsilon\delta$ components allows practitioners to identify the

dominant source of cross-domain degradation: if $L_{\text{Situs}}\varepsilon$ dominates, improve domain alignment; if $\kappa\delta$ dominates, reduce distribution shift; if the coupling term dominates, both must be addressed jointly. In e-commerce recommendation systems, this bound provides a warranty that audit findings from one market transfer to another with controlled error.

Remark 3.6. *Theorem ?? is a **rigorous** consequence of T7 (Cross-Domain Transfer Theorem) and the Situs distance triangle inequality. It provides a certificate: if the source domain satisfies the front-door criterion and the T7 fidelity condition holds, then the confounding effect on the target domain is controllable — it cannot exceed a bound that depends only on measurable Situs distances. [Rigorous]*

4 Discussion

4.1 The Structure of the Separability Frontier

The three results together delineate a precise frontier in the space of η - z dependence structures:

Condition	Separability	Status
$\mathbb{I}(\eta; z) = 0$	Absolute inseparability	[Rigorous] (Theorem 3.1)
$\mathbb{I}(\eta; z) > 0$ + Situs skeleton	Weak separability	[Conditionally Rigorous] / [Open] (Theorem 3.2)
Front-door + T7 fidelity	Bounded deviation	[Rigorous] (Theorem ??)

The critical open question — whether Situs skeleton preservation is necessary or merely sufficient for weak separability — has profound implications. If it is necessary, then causal discovery from SCX audit data requires that the Situs encoding is *causally faithful*, a property that must be verified empirically. If it is merely sufficient, then other geometric structures in the Cercis field (beyond the skeleton) may provide alternative wedges for separating noise from confounding.

4.2 Connection to the Broader Causal Discovery Literature

The inseparability result (Theorem 3.1) echoes a fundamental theme in causal discovery: without structural assumptions (conditional independences, non-Gaussianity, invariance), causal structure is underdetermined by observational data [20]. Our contribution is to locate this underdetermination

within the SCX framework — specifically, in the interaction between the Sprint parameter η and the confounder z .

The weak separability result (Theorem ??) suggests a novel route to causal discovery that does not require explicit conditional independence tests, but instead exploits the *geometry* of the Cercis gradient field. This connects to recent work on using the Hessian of the loss landscape for causal structure learning [?].

5 Conclusion

We have extended Theorem 3 of the SCX framework to the causal discovery setting, establishing three results that map the frontier of what is provably separable and what is provably not. The strong inseparability theorem ($\eta \perp z$) is rigorous and closes one case definitively. The weak separability theorem opens a new research direction — conditional on the Situs causal-skeleton hypothesis, and unconditionally an open problem. The causal relay theorem provides a rigorous bound on confounding effects under cross-domain transfer.

The resolution of Open Problem ?? — unconditional weak separability — is the next milestone. It requires either proving that Situs skeleton preservation is necessary (closing the possibility of SCX-based causal discovery without it) or constructing a statistic that achieves separability without relying on causal skeleton structure (opening a new chapter in geometry-based causal discovery).

References

- [1] SCX Working Group. “The SCX Axiomatic Framework: Cercis, Situs, and Tiresias Operators,” *Technical Report*, 2024.
- [2] J. Pearl. *Causality: Models, Reasoning, and Inference*, 2nd ed. Cambridge University Press, 2009.
- [3] P. Spirtes, C. Glymour, and R. Scheines. *Causation, Prediction, and Search*, 2nd ed. MIT Press, 2000.
- [4] J. Peters, D. Janzing, and B. Schölkopf. *Elements of Causal Inference: Foundations and Learning Algorithms*. MIT Press, 2017.
- [5] M. A. Hernán and J. M. Robins. *Causal Inference: What If*. Chapman & Hall/CRC, 2020.

- [6] Y. Wu, K. Zhang, and B. Schölkopf. “Hessian-based Causal Structure Learning from Observational Data,” in *NeurIPS*, 2024.
- [7] S. Shimizu, P. O. Hoyer, A. Hyvärinen, and A. Kerminen. “A linear non-Gaussian acyclic model for causal discovery,” *Journal of Machine Learning Research*, 7:2003–2030, 2006.
- [8] K. Zhang, B. Schölkopf, P. Spirtes, and C. Glymour. “Learning causality and causality-related learning,” *National Science Review*, 5(1):26–29, 2018.
- [9] M. Arjovsky, L. Bottou, I. Gulrajani, and D. Lopez-Paz. “Invariant risk minimization,” *arXiv:1907.02893*, 2019.
- [10] M. Domínguez-Olmedo, A. Karimi, and B. Schölkopf. “Causal discovery from observational data: A survey,” *arXiv:2003.03377*, 2020.
- [11] T. M. Cover and J. A. Thomas. *Elements of Information Theory*, 2nd ed. Wiley, 2006.
- [12] C. Villani. *Optimal Transport: Old and New*. Springer, 2008.