



All-in-One Microbial Test

Patient Name	Pebless	Health Status	AD Sub tympanic mass	Sample ID	MI19104012
Owner's Name	NA	Ordered By	NA	Sample Type	Ear
Gender	M	Email	NA	Received Date	8/17/2021
Breed	Red Ear Slider	Hospital	NA	Report Date	9/3/2021
Age	10 years	Location	NA		
Species	Turtle	Account Number	NA		

Potential Clinically Relevant Microbes Detected:

Listed are those bacteria and fungi detected in the specimen that are of potential clinical relevance. Results from this report should be considered together with additional clinical data gathered by the veterinarian (physical examination, medical history, cytology, etc.) as the microbes detected may or may not be the cause of the clinical condition. For a comprehensive list of all microorganisms detected in this specimen see page 3 of this report.

1. Bacteria



Species Detected	Relative Abundance (%)	Cells per sample
Mycoplasmataceae sp	83.05	240000
Pseudomonas aeruginosa	16.91	49000

2. Fungi

None Detected

Microbial Overview:

Charts below depict *Bacteria vs Fungi*: an overview of the microbiome, *Bacteria*: the relative abundance of all clinically relevant species detected relative to the rest of the microbiome for bacteria, and *Fungi*: the relative abundance of all clinically relevant species detected relative to the rest of the microbiome for fungi. Each color represents a different species. The larger the size of the colored segment, the more abundant that specific species is in the specimen. The purpose of these graphs is to highlight if any clinically relevant organism is overgrown in the sample.





Patient Name
Owner's Name

Pebless
NA

Ordered By
Account Number

NA
NA

Page 2 of 4

Antibiotic Resistance Panel for Detected Clinically Relevant Microbes

The sample was screened for the presence of antibiotic resistance genes and intrinsic resistances of clinically relevant microorganisms. For this analysis more than 90 antibiotic resistance genes were screened. The **cautious** use of any antibiotic drug is highly recommended. Please follow the guidelines for antimicrobial stewardship in veterinary practice.

Antibiotics	Drug Tiers Follow Guidelines*	Pseudomonas aeruginosa (16.91%)	Suggested Dose for All Pathogens**	Drug Delivery
Cefazolin	1st Line Antibiotics for Common Infections.	NR	15 mg/kg, q 12 hrs	IV, SC
Cephalothin		F	4-20 mg/kg, q 8 hrs	PO
Cephalexin		NR	22 mg/kg, q 12 hrs	PO
Cefadroxil		NR	22 mg/kg, q 12 hrs	PO
Cefoxitin		NR	15 mg/kg, q 12 hrs	IV, SC
Penicillin		G	8-10 mg/kg, q 8 hrs	PO
Penicillin G		G	--	--
Oxacillin		NR	22 mg/kg, q 8 hrs	IV
Ampicillin		NR	22 mg/kg, q 8 hrs	IV, SC
Amoxicillin		NR	22 mg/kg, q 8 hrs	PO
Clavamox		NR	13.75 mg/kg, q 12 hrs	PO
Gentamicin		NR	6 mg/kg, q 24 hrs	IV, SC
Tobramycin		F	--	IV/Topical Use
Neomycin		NR	--	Topical Use
Clindamycin		NR	5.5 mg/kg, q 12 hrs	PO
Lincomycin		NR	15-25 mg/kg, q 24hrs	PO
Doxycycline		NR	5 mg/kg, q 12 hrs	PO
Minocycline		NR	10 mg/kg, q 12 hrs	PO
Tetracycline		NR	20 mg/kg, q 12 hrs	PO
Sulfonamide		G	30 mg/kg, q 12 hrs	PO
Trimethoprim		F	15-30 mg/kg, q 24 hrs	PO
Metronidazole		F	10 mg/kg, q 8 hrs	IV
Cefovecin		NR	8 mg/kg, once	SC
Cefpodoxime	2nd Line Use Caution to Avoid Resistance.	NR	5 mg/kg, q 24 hrs	PO
Ceftiofur		NR	2.2 mg/kg, q 24 hrs	SC
Timentin		NR	--	Topical Use
Azithromycin		NR	5 mg/kg q 12 hrs	PO
Orbifloxacin		G	2.5-7.5 mg/kg, q 24 hrs	PO
Chloramphenicol		F	35 mg/kg q 8 hrs	PO
Florfenicol	3rd Line Last Resort Options.	F	20 mg/kg, q 12 hrs	PO
Amikacin		NR	15 mg/kg, q 24 hrs	IV, SC
Rifampin		NR	5-10 mg/kg, q 12 hrs	PO
Imipenem		NR	10 or 20 mg/kg, q 8 hrs	--
Levofloxacin**		F	10-30 mg/kg, q 24 hrs	IV/PO
Marbofloxacin		F	2.75-5.5 mg/kg, q 24 hrs	PO
Pradofloxacin****		G	3.0 mg/kg, q 24 hrs	PO
Enrofloxacin		G	5 mg/kg, q 24 hrs	PO
Ciprofloxacin***		F	--	Topical Use
Nitrofurantoin		G	4.4-5mg/kg, q 24 hrs	PO
Colistin**		F	8-9g/kg, q 24 hrs	PO
Ceftazidime		G	3-30 mg/kg, q 6-8 hrs	IV
Mupirocin		F	--	Topical Use
Piperacillin		G	80-100 mg/kg, 30min q 8 hrs	IV
Ticarcillin		F	3.1 g, q4-6 hrs	IV

This table lists antibiotic sensitivities/resistances for the indicated bacteria based on detection of specific antibiotic resistance genes and naturally occurring, or intrinsic, resistance to specific antibiotics previously identified for that organism. To receive a list of the antibiotic resistance genes detected as well as intrinsic resistances for additional organisms not listed here, please contact MiDOG® customer support.

Abbreviation Key:

NR	Not Recommended (Due to either Intrinsic Resistance, or Resistance Gene Detection, or < 10% Effectiveness in Antibigram Studies)
P	Poor Performance (< 50% Effectiveness in Antibigram Studies)
F	Fair Performance (< 75% Effectiveness in Antibigram Studies)
G	Good Performance (> 75% Effectiveness in Antibigram Studies)
	No Literature Information Available

PO = Oral, By Mouth.

IV = Intravenous Injection. Injections involving direct injection into the vein.

SC = Subcutaneous Injection. The medication delivered to the tissues between the skin and the muscle.

TU = Topical Use

- * Reference: Antimicrobial Resistance and Stewardship Initiative University of Minnesota, Antibiotic Drug Tiers and Selection List for Companion Animals.
- ** Dosis may vary based on patient species and/or type of infection. Reference at: www.midogtest.com/antibiotics
- *** Variable bioavailability in canine patients
- **** Contraindicated in canine patients



Patient Name
Owner's Name

Pebless
NA


Ordered By
Account Number

NA
NA

Supplemental Data on Microbial Composition

Bacterial Analysis

Charts below depict the relative abundance of all detected bacterial species. Each color represents a different bacterial species. The larger the size of the colored segment, the more abundant that specific species is in the specimen.



The table below lists all bacterial species detected in the specimen within the limit of detection. The absolute and relative abundances of each species is shown.

Species Detected	Relative Abundance (%)	Cells per sample
Mycoplasmataceae sp	83.05	240000
Pseudomonas aeruginosa	16.91	49000

Fungal Analysis

No fungal species detected. Please Note: It is common to see no fungal load in healthy urine samples. For more information see the Methods Section under: 'When no Bacterial or Fungal Species are Detected'



Patient Name
Owner's Name

Pebless
NA

Ordered By
Account Number

NA
NA

Page 4 of 4

References

1. Rodrigues Hoffmann A, Patterson AP, Diesel A et al. The skin microbiome in healthy and allergic dogs. (2014) PLoS One, 9: e8197

Method

The MiDOG® All-in-One Microbial Test is a targeted, Next-generation DNA sequencing testing service able to identify molecular signatures unique to the identity and character of a specific microorganism. This test relies on safeguarded preservation and transport of collected samples, thorough extraction of DNA from all microbes present in the specimen, select amplification of microbial DNA followed by Next-generation DNA sequencing using the latest technologies from Illumina (Illumina, Inc., San Diego, CA). Data handling is done via curated microbial databases to accurately align DNA sequences to ensure precise and accurate (species-level) identification of all bacteria and fungi present in the specimen.

When no Bacterial or Fungal Species are Detected

When no bacterial or fungal species are detected in this test, this result may be due to a very low microbial load and/or low concentration of microbial DNA in the sample provided. In this case, we recommend re-sampling the area of interest and re-submitting specimen for analysis.

Disclaimer

The information contained in this MiDOG® report is intended only to be factor for use in a diagnosis and treatment regime for the canine patient. As with any diagnosis or treatment regime, you should use clinical discretion with each canine patient based on a complete evaluation of the canine patient, including history, physical presentation and complete laboratory data, including confirmatory tests. All test results should be evaluated in the context of the patients individual clinical presentation. The information in the MiDOG® report has not been evaluated by the FDA.

Customer Support

Tel: (833)456-4364
info@midogtest.com
www.midogtest.com