



All-in-One Microbial Test

Page 1 of 4

Patient Name	Bob	Health Status	NA	Sample ID	MI19104008
Owner's Name	Stephanie Klein	Ordered By	Dr. Malka	Sample Type	Skin, Abscess on bottom of foot
Gender	NA	Email	office@birdexoticsvet.com	Received Date	08/20/21
Breed	NA	Hospital	NA	Report Date	8/25/2021
Age	15 yrs	Location	NA		
Species	NA	Account Number	100A233		

Potential Clinically Relevant Microbes Detected:

Listed are those bacteria and fungi detected in the specimen that are of potential clinical relevance. Results from this report should be considered together with additional clinical data gathered by the veterinarian (physical examination, medical history, cytology, etc.) as the microbes detected may or may not be the cause of the clinical condition. For a comprehensive list of all microorganisms detected in this specimen see page 3 of this report.

1. Bacteria



Species Detected	Relative Abundance (%)	Cells per sample
Pseudomonas aeruginosa	98.0	690000

2. Fungi

None Detected

Microbial Overview:

Charts below depict *Bacteria vs Fungi*: an overview of the microbiome, *Bacteria*: the relative abundance of all clinically relevant species detected relative to the rest of the microbiome for bacteria, and *Fungi*: the relative abundance of all clinically relevant species detected relative to the rest of the microbiome for fungi. Each color represents a different species. The larger the size of the colored segment, the more abundant that specific species is in the specimen. The purpose of these graphs is to highlight if any clinically relevant organism is overgrown in the sample.





Patient Name Bob
Owner's Name Stephanie Klein

Ordered By
Account Number

Dr. Malka
100A233

Page 2 of 4

Antibiotic Resistance Panel for Detected Clinically Relevant Microbes

The sample was screened for the presence of antibiotic resistance genes and intrinsic resistances of clinically relevant microorganisms. We recommend the cautious use of any antibiotic drug and to follow the guidelines for antimicrobial stewardship in veterinary practice.

Antibiotics	<u><i>Pseudomonas aeruginosa</i></u> (98.0%)	Suggested Dose	Drug Delivery
Ampicillin	IR		
Amoxicillin	IR		
Oxacillin	ND	--	--
Benzylpenicillin	ND	--	--
Cefadroxil	IR		
Cefazolin	IR		
Cefpodoxime	IR		
Ceftiofur	IR		
Cephalexin	ND	22 mg/kg, q 12 hours	PO
Cefovecin	IR		
Cefoxitin	IR		
Enrofloxacin	ND	5 mg/kg, q 24hrs	PO
Marbofloxacin	ND	2.75 mg/kg, q 24hrs	PO
Ciprofloxacin	ND	--	--
Pradofloxacin	ND	--	--
Clindamycin	IR		
Lincomycin	ND	--	--
Erythromycin	IR		
Azithromycin	IR		
Sulfonamide	ND	30 mg/kg, q 12hrs	PO
Doxycycline	R		
Tetracycline	R		
Minocycline	R		
Neomycin	R		
Amikacin	R		
Gentamicin	R		
Clavamox	IR		
Timentin	ND	Topical use	Topical use
Ceftazadime	ND	--	--
Florfenicol	ND	--	--
Chloramphenicol	IR		
Imipenem	ND	--	--
Mupirocin	ND	Topical use	Topical use
Rifampicin	IR		

This table lists antibiotic sensitivities/resistances for the indicated bacteria based on detection of specific antibiotic resistance genes and naturally occurring, or intrinsic, resistance to specific antibiotics previously identified for that organism. To receive a list of the antibiotic resistance genes detected as well as intrinsic resistances for additional organisms not listed here, please contact MiDOG® customer support.

Abbreviation Key:

ND = Not Detected. According to our test the organism is most likely sensitive to this antibiotic. No antibiotic resistance gene or intrinsic resistance was detected.

R = Resistance. An Antibiotic resistance gene for the indicated antibiotic was detected and the organism is likely resistant to the suggested dose.

IR = Intrinsic Resistance. No resistance genes are detected or available for testing, would not recommend this antibiotic due to likely resistant to the indicated antibiotic based on intrinsic factors inferred from microbiological and clinical literature.

PO = Oral. By Mouth.

IV = Intravenous Injection. Injections involving direct injection into the vein.

SC = Subcutaneous Injection. The medication delivered to the tissues between the skin and the muscle.




Patient Name Bob
Owner's Name Stephanie Klein

Ordered By Dr. Malka
Account Number 100A233

Supplemental Data on Microbial Composition

Bacterial Analysis

Charts below depict the relative abundance of all detected bacterial species. Each color represents a different bacterial species. The larger the size of the colored segment, the more abundant that specific species is in the specimen.



The table below lists all bacterial species detected in the specimen within the limit of detection. The absolute and relative abundances of each species is shown.

Species Detected	Relative Abundance (%)	Cells per sample
Pseudomonas aeruginosa	98.00	690000
Dietzia sp	1.94	13000

Fungal Analysis

No fungal species detected. Please Note: It is common to see no fungal load in healthy urine samples. For more information see the Methods Section under: 'When no Bacterial or Fungal Species are Detected'



Patient Name Bob
Owner's Name Stephanie Klein

Ordered By Dr. Malka
Account Number 100A233

Page 4 of 4

References

1. Rodrigues Hoffmann A, Patterson AP, Diesel A et al. The skin microbiome in healthy and allergic dogs. (2014) PLoS One, 9: e8197

Method

The MiDOG® All-in-One Microbial Test is a targeted, Next-generation DNA sequencing testing service able to identify molecular signatures unique to the identity and character of a specific microorganism. This test relies on safeguarded preservation and transport of collected samples, thorough extraction of DNA from all microbes present in the specimen, select amplification of microbial DNA followed by Next-generation DNA sequencing using the latest technologies from Illumina (Illumina, Inc., San Diego, CA). Data handling is done via curated microbial databases to accurately align DNA sequences to ensure precise and accurate (species-level) identification of all bacteria and fungi present in the specimen.

When no Bacterial or Fungal Species are Detected

When no bacterial or fungal species are detected in this test, this result may be due to a very low microbial load and/or low concentration of microbial DNA in the sample provided. In this case, we recommend re-sampling the area of interest and re-submitting specimen for analysis.

Disclaimer

The information contained in this MiDOG® report is intended only to be factor for use in a diagnosis and treatment regime for the canine patient. As with any diagnosis or treatment regime, you should use clinical discretion with each canine patient based on a complete evaluation of the canine patient, including history, physical presentation and complete laboratory data, including confirmatory tests. All test results should be evaluated in the context of the patients individual clinical presentation. The information in the MiDOG® report has not been evaluated by the FDA.

Customer Support

Tel: (833)456-4364
info@midogtest.com
www.midogtest.com