



Weekly Quizzes

Review Test Submission: Week 06 Quiz

Review Test Submission: Week 06 Quiz

User Dong Gao

Subject Algorithms and Complexity

Test Week 06 Quiz

Started 15/04/16 4:31 PM

Submitted 15/04/16 4:48 PM

Due Date 20/04/16 11:59 PM

Status Completed

Attempt 4 out of 4 points

Score

Time 17 minutes

Elapsed

Instructions You should attempt the quiz after the lecture and your tutorial.

- The quiz is available for a period of 10 days.
- You may attempt the quiz multiple times (if you happen to get a question wrong, you can do it again)
- Your score on the quiz will be recorded in the grade book. The score is not used when determining your final mark in this subject
- The quiz might not display equations correctly in some browsers. If you experience problems, we recommend that you use Firefox.

Note: you must complete at least eight of the weekly quizzes to meet one of the hurdle requirements in this subject

Results All Answers, Submitted Answers, Feedback, Incorrectly Answered Questions

Displayed

Question 1

1 out of 1 points



Quicksort uses Hoare partitioning. Assume an array contains ten keys: 6 3 1 7 9 5 8 2 4 0. After a first round of simple Hoare partitioning (not median-of-three), the array looks like so:

Selected Answer: b. 2 3 1 0 4 5 6 8 9 7

Answers: a. 5 3 1 0 4 2 6 8 9 7

b. 2 3 1 0 4 5 6 8 9 7

c. 3 1 5 2 4 0 6 7 9 8

d. 2 3 0 1 5 4 6 7 8 9

e. 5 3 1 4 0 2 6 7 9 8

Response Feedback: Well done!

Question 2

1 out of 1 points



A complete binary tree has this inorder traversal sequence: 7, 4, 1, 0, 8, 5, 6, 3, 9, 2. What is the key at its root?

Selected Answer: 6

Response Feedback: Yes, that's right.

Question 3

1 out of 1 points



Consider this recurrence relation:

$$T(1) = 1$$

$$T(n) = 2 T(n/3) + 2n + 1 \quad \text{for } n > 1$$

The Master Theorem says that

Selected Answer: $T(n) \in \Theta(n)$

e.

Answers: $T(n) \in \Theta(n^3)$

a.

$$T(n) \in \Theta(n^2)$$

b.

$$T(n) \in \Theta(n \log n)$$

c.

$$T(n) \in \Theta(n \log \log n)$$

d.

$$T(n) \in \Theta(n)$$

e.

Response
Feedback:

That's right. In this case we have $a=2$, $b=3$, and $d=1$. And indeed $2 < 3$.

Question 4

1 out of 1 points



Consider this recurrence relation:

$$T(1) = 1$$

$$T(2) = 1$$

$$T(n) = 4 T(n-2) + 2n^2 \quad \text{for } n > 2$$

The Master Theorem tells us

Selected Answer: e. nothing

- Answers:
- a. $T(n) \in \Theta(n^3)$
 - b. $T(n) \in \Theta(n^2 \log n)$
 - c. $T(n) \in \Theta(n^2)$
 - d. $T(n) \in \Theta(n \log n)$
 - e. nothing

Response Feedback: That's right, the Master Theorem does not help here, as the recurrence is not of the required form.

Saturday, 4 June 2016 11:15:40 PM EST

← OK