Toward a Counterexample to Agrawal's Conjecture via Twin Smooth Integers

(working draft — v0.1)

Dan Shumow Claude (Anthropic Fable 5)

August 6, 2026

Audit notice (22 October 2026). The proof of Theorem 2.1 and the modulus normalization in Section 3 have been corrected on the joint-correctness branch. The remaining feasibility estimates are historical, not validated by this correction. The same-name PDF beside this source is the original draft; the rebuilt audit version is in `doc/audit-pdfs/`. Current audit status is recorded in `doc/integration-review-status.md`.

Abstract

Agrawal's conjecture asserts that for coprime n and prime r, the congruence $(X-1)^n \equiv X^n - 1 \pmod{n,\, X^r-1}$ forces n prime or $n^2 \equiv 1 \pmod r$; if true it yields a deterministic $\tilde{O}(\log^3 n)$ primality test. Lenstra and Pomerance gave a heuristic construction suggesting counterexamples exist but are astronomically large; none is known, and exhaustive search is hopeless. We pursue a *constructive* search for a counterexample of unrestricted size. Our contributions so far: (i) a reduction of the Lenstra–Pomerance construction to a two-sided subset-product problem over a partitioned smooth factor base; (ii) the observation that the candidate pool consists exactly of *twin smooth integers with prime sum* in a fixed congruence class, connecting the search to the isogeny-cryptography parameter literature and its datasets; (iii) feasibility estimates from that literature's empirical supply curves, isolating a single unmeasured quantity — the "coloring tax" — as the go/no-go criterion; and (iv) an obstruction: the boosting polynomials $2x^n - 1$ of the twin-smooth literature can never satisfy the local conditions of the Lenstra–Pomerance theorem as stated, motivating a re-derivation of its admissible local conditions.

1 Introduction

For coprime integers n, r write $T(-1, n, r)$ for the congruence

$$(X-1)^n \equiv X^n - 1 \pmod{n,\, X^r-1}. \tag{1}$$

Conjecture 1.1 (Agrawal; Bhattacharjee–Pandey [2])**.** *If r is a prime not dividing n and $T(-1,n,r)$ holds, then n is prime or $n^2 \equiv 1 \pmod r$.*

If Conjecture 1.1 were true, a single evaluation of (1) at one $r = O(\log n)$ would certify primality, giving a deterministic $\tilde{O}(\log^3 n)$ test, versus $\tilde{O}(\log^6 n)$ for the best proven algorithm [7]. Its status is strange: Lenstra and Pomerance [3] gave a heuristic argument for $\gg x^{1-\varepsilon}$ counterexamples below x, yet the conjecture has survived exhaustive search to 10^{17} (Primaboinca, 2010–2020) and folklore places the least counterexample at hundreds of digits. Popovych [4] conjectured that adding the hypothesis $(X+2)^n \equiv X^n + 2 \pmod{n,\, X^r-1}$ repairs the statement; his variant has no evidence against it.

Program. We abandon minimality. Any counterexample refutes Conjecture 1.1; the L–P density heuristic *improves* with size; and the construction of §2 is deterministic — when its arithmetic side conditions hold, (1) follows with no further luck required. The size of a counterexample produced this way is bounded only by the combinatorics of a subset-product problem (§3), whose raw material turns out to be the *twin smooth integers* of isogeny-based cryptography (§4). Verifying a candidate n of even 10^5 digits directly against (1) costs a few core-hours, and any find will also be tested against Popovych's second congruence.

2 The Lenstra–Pomerance construction

Theorem 2.1 (Lenstra–Pomerance [3]; statement following Váňa [5], who proves the $k \equiv 3 \pmod 4$ case)**.** *Let $n = p_1 \cdots p_k$ with $p_1, \ldots, p_k$ distinct primes such that*

(a) k is odd;

(b) $p_i \equiv 3 \pmod{80}$ for all i;

(c) $p_i - 1 \mid n - 1$ for all i *(n is Carmichael);*

(d) $p_i + 1 \mid n + 1$ for all i *(n is Lucas–Carmichael).*

Then $T(-1, n, 5)$ holds and $n^2 \not\equiv 1 \pmod 5$; in particular such an n with $k \geq 3$ is a counterexample to Conjecture 1.1.

Proof. Since $p \equiv 3 \pmod 5$ has order 4 in $(\mathbb{Z}/5)^\times$, over $\mathbb{F}_p$ the polynomial $X^5 - 1$ factors as $(X-1)\Phi_5(X)$ with Φ_5 irreducible, so $\mathbb{Z}[X]/(p, X^5 - 1) \cong \mathbb{F}_p \times \mathbb{F}_{p^4}$. The $\mathbb{F}_p$ component of (1) is trivial. In $\mathbb{F}_{p^4}$, put $\alpha = \zeta - 1$ for a primitive fifth root ζ. Since $p^2 \equiv -1 \pmod 5$, Frobenius gives $\alpha^{p^2} = -\zeta^{-1}\alpha$, so $\alpha^{10(p^2-1)} = 1$.

Let $j \in \{1, 3\}$ satisfy $j \equiv k \pmod 4$, and put $m = (p-1)/2$, $a = (p+1)/4$. The numbers $80, m, a$ are pairwise coprime: m, a are odd, neither is divisible by 5, and $\gcd(m, a) = 1$. Also $10(p^2 - 1) = 80ma$. Conditions (c),(d) imply $n \equiv p^j \pmod m$ and $n \equiv p^j \pmod a$, since $p \equiv 1 \pmod m$ and $p \equiv -1 \pmod a$. Finally $n \equiv 3^k \equiv 3^j \equiv p^j \pmod{80}$, using $3^4 \equiv 1 \pmod{80}$. CRT therefore yields $n \equiv p^j \pmod{10(p^2-1)}$ in both odd-cardinality cases. Consequently $\alpha^n = \alpha^{p^j} = \zeta^{p^j} - 1 = \zeta^n - 1$. This proves the congruence modulo each p_i; squarefreeness and CRT prove it modulo n. Lemma 2.2 gives $n^2 \not\equiv 1 \pmod 5$. □

Lemma 2.2. *If $p_i \equiv 3 \pmod 5$ for all i and k is odd, then $n \equiv 3^k \equiv \pm 2 \pmod 5$; hence $n^2 \equiv 4 \not\equiv 1 \pmod 5$.*

Proof. $3^k \bmod 5$ cycles $3, 4, 2, 1$ with period 4; odd k gives 3 or 2, i.e. $\pm 2 \pmod 5$. □

The difficulty of satisfying (c) and (d) simultaneously is concentrated in the following trivial but decisive observation.

Lemma 2.3 (Side-coprimality)**.** *If n satisfies (c) and (d), then $\gcd(p_i - 1, p_j + 1) \mid 2$ for all i, j (including $i = j$). Consequently the odd primes dividing $\prod_i (p_i - 1)$ and those dividing $\prod_i (p_i + 1)$ form disjoint sets.*

Proof. An odd prime q dividing $p_i - 1$ and $p_j + 1$ divides both $n - 1$ and $n + 1$, hence their difference 2, a contradiction. □

Remark 2.4. No integer that is simultaneously Carmichael and Lucas–Carmichael is known; none exists below 10^{18} [8]. Lemma 2.3 is the structural reason, and the reason the *least* counterexample is expected to be enormous. It does not, however, obstruct *large* solutions, which is what we seek.

Lemma 2.5 (2-adic bookkeeping)**.** *If* $p \equiv 3 \pmod{16}$ *then* $v_2(p-1) = 1$ *and* $v_2(p+1) = 2$*; if* $p \equiv 3 \pmod 5$ *then* $5 \nmid (p-1)(p+1)$*. If moreover every* $p_i \equiv 3 \pmod 4$ *and* k *is odd, then* $n \equiv 3 \pmod 4$*, so the 2-power parts of conditions (c),(d) hold automatically.*

Proof. $p \equiv 3 \pmod{16}$ gives $p-1 \equiv 2$ and $p+1 \equiv 4 \pmod{16}$, fixing the stated 2-adic valuations; $p \equiv 3 \pmod 5$ gives $p \mp 1 \equiv 2, 4 \pmod 5$. For the last claim, $n \equiv 3^k \equiv 3 \pmod 4$ for odd k, so $2 \mid n-1$ with $v_2(n-1) \geq 1 = v_2(p_i - 1)$ and $4 \mid n+1$ with $v_2(n+1) \geq 2 = v_2(p_i+1)$. □

Thus under condition (b) the entire problem lives in odd moduli prime to 5.

3 Reduction to a two-sided subset-product problem

Fix disjoint finite sets Q_-, Q_+ of odd primes with $5 \notin Q_- \cup Q_+$, and define the *pool*

$$\mathcal{P} \;=\; \Big\{ p \text{ prime} : p \equiv 3 \pmod{80},\ \tfrac{p-1}{2} \text{ is } Q_-\text{-smooth},\ \tfrac{p+1}{4} \text{ is } Q_+\text{-smooth} \Big\}.$$

For a nonempty pool let

$$\Lambda_- = \operatorname{lcm}\{p-1 : p \in \mathcal{P}\} = 2\,\operatorname{lcm}\{(p-1)/2 : p \in \mathcal{P}\},$$
$$\Lambda_+ = \operatorname{lcm}\{p+1 : p \in \mathcal{P}\} = 4\,\operatorname{lcm}\{(p+1)/4 : p \in \mathcal{P}\}.$$

The factors 2 and 4 belong outside the lcm of the *normalized odd parts*, not outside the lcm of $p-1$ and $p+1$.

Proposition 3.1. *Let* $S \subseteq \mathcal{P}$ *with* $|S|$ *odd,* $|S| \geq 3$*, and* $n = \prod_{p \in S} p$*. If*

$$n \equiv 1 \pmod{\Lambda_-} \qquad \text{and} \qquad n \equiv -1 \pmod{\Lambda_+}, \tag{2}$$

then n *satisfies conditions (a)–(d) of Theorem 2.1, hence is a counterexample to Conjecture 1.1.*

Proof. (a),(b) hold by construction of $\mathcal{P}$ and $|S|$. For (c): $p_i - 1 \mid \Lambda_-$ by definition of Λ_-, and $\Lambda_- \mid n-1$ by (2). Likewise (d) via $\Lambda_+ \mid n+1$. The elements of S are distinct primes, so n is squarefree with $k = |S| \geq 3$, hence composite. Disjointness of $Q_\pm$ and the valuations in Lemma 2.5 give $\gcd(\Lambda_-, \Lambda_+) = 2$. Thus the two target residues are CRT-compatible. This is a statement about the congruences, not a proof that a subset of the pool realizes them. □

Condition (2) is *stronger* than (c),(d) — it imposes the full lcm rather than the lcm over S. Before taking discrete logarithms, restrict to pool elements coprime to $\Lambda_-\Lambda_+$; selected elements must be units because the target residues are units. In the production design, pool primes exceeding every factor-base prime satisfy this restriction automatically. Taking discrete logarithms in each cyclic component of $(\mathbb{Z}/\Lambda_-)^\times \times (\mathbb{Z}/\Lambda_+)^\times$ (Pohlig–Hellman [11] is cheap, all component orders being smooth by design), (2) becomes a 0/1 subset-sum over $\bigoplus_i \mathbb{Z}/m_i$, with one extra $\mathbb{Z}/2$ coordinate enforcing $|S|$ odd. The heuristic solvability condition, under a sufficiently uniform subset-sum model, is

$$2^{|\mathcal{P}|} \;\gg\; |G|, \qquad G = (\mathbb{Z}/\Lambda_-)^\times \times (\mathbb{Z}/\Lambda_+)^\times, \tag{3}$$

This size inequality by itself guarantees collisions, not reachability of the specified target or an odd-cardinality solution. i.e. $|\mathcal{P}| \gtrsim \log_2 \Lambda_- + \log_2 \Lambda_+$ with slack for the solver. Candidate solvers,

in escalating order: staged CRT in the style of Löh–Niebuhr's Carmichael factories [10], Wagner's k-list generalized birthday algorithm [12] (designed for exactly this pool-rich regime), and lattice reduction only for terminal low-dimensional corrections. **[TODO: Phase 2: work out the staged-CRT schedule and the expected $|S|$.]**

4 The twin-smooth dictionary

Lemma 4.1 (Dictionary)**.** *Let $m \geq 1$ and $p = 2m + 1$. Then $p \in \mathcal{P}$ (for $Q_\pm$ large enough to contain the relevant primes) if and only if*

(i) m and $m + 1$ are both smooth with respect to $Q_- \cup Q_+ \cup \{2\}$, with the odd part of m being Q_--smooth and the odd part of $m + 1$ being Q_+-smooth;

(ii) $m \equiv 1 \pmod{40}$;

(iii) p is prime.

Moreover (ii) alone implies $p \equiv 3 \pmod{80}$, $v_2(p-1) = 1$, $v_2(p+1) = 2$, and $5 \nmid m(m+1)$.

Proof. $p - 1 = 2m$ and $p + 1 = 2(m + 1)$, so the smoothness conditions on $(p - 1)/2 = m$ and $(p + 1)/4 = (m + 1)/2$ are exactly (i) once the 2-adic valuations match. For (ii): $m \equiv 1 \pmod{40} \iff p = 2m + 1 \equiv 3 \pmod{80}$. Then m is odd (so $v_2(p - 1) = 1$), $m \equiv 1 \pmod 4$ gives $m + 1 \equiv 2 \pmod 4$ (so $v_2(p + 1) = 2$), and $m \equiv 1 \pmod 5$ gives $m \not\equiv 0$, $m + 1 \equiv 2 \not\equiv 0 \pmod 5$. □

Pairs of consecutive smooth integers $(m, m + 1)$ — *twin smooths* — with $2m + 1$ prime are precisely the parameters sought for the isogeny-based schemes B-SIDH [13] and SQIsign, and 2020–2025 produced an industrial literature on generating them:

- **PTE sieving** (Costello–Meyer–Naehrig [14]): Prouhet–Tarry–Escott solutions give degree-n polynomial pairs differing by a constant that split into linear factors; yielded 240–256-bit primes p with $p^2 - 1$ being 2^{15}-smooth, up to 512-bit at 2^{28}-smooth.
- **CHM** (Bruno et al. [15], optimizing Conrey–Holmström–McLaughlin [16]): iterative generation of *near-complete* sets of twin B-smooths. Headline data: $B = 547$ yields 82,026,426 pairs, the largest of 122 bits; implementation public.
- **Pell/Størmer** (Buzek–Hasan–Liu–Naehrig–Vigil [17]; classical [20, 21]): for fixed B the set of twin B-smooths is *finite*, computable via Pell equations $x^2 - 2Dy^2 = 1$; complete sets known provably for $B \leq 100$ and heuristically for $B \leq 113$ [19].
- **SVP records** (Mulder–Sterner–van Woerden [19]): shortest vectors in a prime-number lattice locate the *largest* twin for a given B (records: 196-bit at $B = 751$, 213-bit at $B = 997$), with an estimator for the largest twin as a function of B.
- **Boosting** (in [15, 18]): evaluate $p_n(x) = 2x^n - 1$ at twin-smooth arguments to reach cryptographic sizes.

By Lemma 4.1, a 1/40 congruence slice of *any* twin-smooth dataset, filtered for primality of $2m + 1$, is a ready-made pool $\mathcal{P}$ — before the one condition their literature does not consider: the global side-partition of Lemma 2.3.

5 Feasibility estimates

5.1 Supply and demand at $B = 547$

Demand. $\log_2 \Lambda_- + \log_2 \Lambda_+ \le \sum_{2<q\le B} \log_2 q^{e_q}$ where q^{e_q} is the largest power appearing in any used $p \mp 1$; the side-partition splits this sum but does not double it. With $\pi(547) = 101$ primes and exponents capped by observed usage, this is $\approx \theta(547)/\ln 2 \approx 750$ bits plus modest slack from powers of small primes: call it 1000–1500 bits, and less if the chosen subpool touches only a sub-universe of primes.

Supply. 82M pairs at $B = 547$, $\div 40$ for the congruence slice (Lemma 4.1(ii)), $\div \ln p \approx 20$–55 for primality of $2m+1$ at the dataset's typical sizes: roughly 5×10^4 pool candidates *from existing data*. **[TODO: Phase 1a: regenerate the dataset and replace these estimates with exact histograms.]**

The cushion between supply and demand is therefore a factor ≈ 25–40. Moreover it *grows* with B: demand grows linearly ($\approx 1.44\,B$ bits) while empirical twin counts grow like B^α with $\alpha \approx 2.5$–3 (from $B = 113 \to 547$). Størmer finiteness caps the supply at each fixed B, but the ceiling recedes superlinearly.

5.2 The coloring tax

The unmeasured factor is the side-partition. Each pool element consumes $\omega_-(p) + \omega_+(p)$ prime-side assignments (the number of distinct odd primes dividing $(p-1)/2$ and $(p+1)/4$), and all chosen elements must respect one partition $Q_- \sqcup Q_+$. Under a uniformly random balanced partition, an element survives with probability $\approx 2^{-(\omega_- + \omega_+)}$, so the expected number of survivors is $\sum_{p\in\mathcal{P}} 2^{-\omega(p)}$. At $B = 547$, bulk-of-distribution twins (30–50 bits) have $\omega_- + \omega_+ \approx 8$–$14$, predicting only *tens* of survivors from 5×10^4 candidates — two orders of magnitude short of the ∼1500–3000 needed.

The program therefore rests on the question:

Question 5.1. *By how much does an* optimized *partition beat a random one? Survivors correlate (elements sharing factor support survive together), supply is thickest exactly where ω is smallest, and demand shrinks with the effective prime universe. The complete Størmer sets ($B \le 113$) permit computing the exact optimum at small scale; the CHM set at $B = 547$ permits evaluating it at target scale.* ***[TODO: Phase 1a go/no-go: survival rate $\gtrsim 5\%$ means the direct route is live; $\lesssim 0.1\%$ kills it at fixed B.]***

5.3 Heuristic assumptions ledger

Statement	Status	Role
$\gcd(p-1, p+1) = 2$; Lemmas 2.2–4.1	theorems	structure
Størmer finiteness at fixed B	theorem	supply ceiling
Local densities $P(q \mid p \mp 1) = \frac{1}{q-1}$, exclusive	exact	constants only
Two-sided smoothness independence $(\rho(u_-)\rho(u_+))$	conjectural	steering only
One-sided density $\#\{p : p-1 \text{ smooth}\} \sim \rho(u)\pi(x)$	open (Erdős–Pomerance line)	steering only
Empirical supply curves (CHM, Størmer sets)	data	replaces both

6 An obstruction to boosting, and admissible local conditions

The boosting polynomials $p_n(x) = 2x^n - 1$ of [15, 18] are structurally attractive here: for $n = 2$, $(p+1)/2 = x^2$ and $(p-1)/2 = (x-1)(x+1)$ with $\gcd(x, x^2-1) = 1$, so the two sides have automatically disjoint prime support — Lemma 2.3 satisfied by construction — and the two-sided smoothness burden moves to (x, x^2-1) at *half* the bit-size, where Dickman densities are far kinder. Unfortunately:

Proposition 6.1. *For every $n \geq 2$ and every integer x: $2x^n - 1 \not\equiv 3 \pmod{16}$. In particular no value of any p_n, $n \geq 2$, satisfies condition (b) of Theorem 2.1. For $n = 2$ the obstruction is doubled: $x^2 \equiv 2 \pmod 5$ is also unsolvable, while for $n = 3$ the mod-5 condition is satisfiable ($x \equiv 3 \pmod 5$) and only the 2-adic obstruction remains.*

Proof. $2x^n \equiv 4 \pmod{16}$ requires $x^n \equiv 2 \pmod 8$. If x is even then $4 \mid x^n$ (as $n \geq 2$), so $x^n \equiv 0$ or $4 \pmod 8$; if x is odd then x^n is odd. In no case is $x^n \equiv 2 \pmod 8$. For the mod-5 claims: the squares modulo 5 are $\{0, 1, 4\} \not\ni 2$, while $3^3 = 27 \equiv 2 \pmod 5$. □

The obstruction is to condition (b) *as stated*, not (necessarily) to the theorem's mechanism. Condition (b) fixes one convenient cell of local conditions ($p \equiv 3 \bmod 16$ giving $v_2(p \mp 1) = (1, 2)$, and $\mathrm{ord}_5(p) = 4$); Váňa's $\rho(p)$ framework suggests other cells may be admissible, e.g. pools with a uniform but different 2-adic pattern $v_2(p+1) = v$ for $v > 2$, compensated in the mod-2^k component of the target congruence (2).

Question 6.2 (Admissible local cells)**.** *Determine the full set of local conditions (residues mod 2^k and mod 5, allowed v_2 patterns, and the required uniformity across the pool) under which the conclusion of Theorem 2.1 survives. In particular: is any admissible cell reachable by $p_n(x) = 2x^n - 1$ for some n, or by another side-separating polynomial family? A positive answer would collapse the coloring tax of §5.2 structurally and halve the effective smoothness size.*

For $n = 3$: $p = 2x^3 - 1$ has $v_2(p+1) = 1 + 3v_2(x)$, so $x \equiv 2 \pmod 4$ gives the uniform pattern $v_2(p+1) = 4$, $v_2(p-1) = 1$ — a natural first test case for Question 6.2. **[TODO: Phase 0.]**

7 Program

P0. Independent re-derivation of Theorem 2.1; resolution of Question 6.2; numerical unit tests of the congruence machinery; prior-art due diligence (in particular whether the L–P construction was ever run at scale).

P1a. Data-first go/no-go: regenerate the CHM $B = 547$ set, apply the dictionary filters, and answer Question 5.1 by direct optimization; calibrate against the complete $B \leq 113$ sets.

P1b. If needed, a native harvester (constructive enumeration of one side, trial division on the other) at larger B or targeted congruences.

P2. Subset-product solver (staged CRT, Wagner), with the parity coordinate.

P3. Assembly and dual independent verification of any find, including the direct computation of $T(-1, n, 5)$ and the Popovych congruence; write-up (a quantified negative result at fixed B is also publishable).

References

[1] M. Agrawal, N. Kayal, N. Saxena, *PRIMES is in P*, Ann. of Math. **160** (2004), 781–793.

[2] R. Bhattacharjee, P. Pandey, *Primality testing*, BTech thesis, IIT Kanpur, 2001.

[3] H. W. Lenstra, Jr., C. Pomerance, *Remarks on Agrawal's conjecture*, AIM workshop notes, 2003. https://aimath.org/WWN/primesinp/articles/html/50a/

[4] R. Popovych, *A note on Agrawal conjecture*, Cryptology ePrint 2009/008.

[5] T. Váňa, *Agrawal's conjecture and Carmichael numbers*, Comenius University Bratislava, 2009.

[6] A. Hegde, P. Devaraj, *Heuristics for the construction of counterexamples to the Agrawal conjecture*, Springer PROMS **344**, 2021.

[7] H. W. Lenstra, Jr., C. Pomerance, *Primality testing with Gaussian periods*, J. Eur. Math. Soc. **21** (2019), 1229–1269.

[8] R. G. E. Pinch, *The Carmichael numbers up to* 10^{18}, arXiv:math/0604376.

[9] W. R. Alford, A. Granville, C. Pomerance, *There are infinitely many Carmichael numbers*, Ann. of Math. **139** (1994), 703–722.

[10] G. Löh, W. Niebuhr, *A new algorithm for constructing large Carmichael numbers*, Math. Comp. **65** (1996), 823–836.

[11] S. Pohlig, M. Hellman, *An improved algorithm for computing logarithms over* $GF(p)$*...*, IEEE Trans. Inf. Theory **24** (1978), 106–110.

[12] D. Wagner, *A generalized birthday problem*, CRYPTO 2002.

[13] C. Costello, *B-SIDH: supersingular isogeny Diffie–Hellman using twisted torsion*, ASIACRYPT 2020.

[14] C. Costello, M. Meyer, M. Naehrig, *Sieving for twin smooth integers with solutions to the Prouhet–Tarry–Escott problem*, EUROCRYPT 2021; ePrint 2020/1283.

[15] G. Bruno, M. Corte-Real Santos, C. Costello, J. K. Eriksen, M. Meyer, M. Naehrig, B. Sterner, *Cryptographic smooth neighbors*, ASIACRYPT 2023; ePrint 2022/1439.

[16] J. B. Conrey, M. A. Holmström, T. L. McLaughlin, *Smooth neighbors*, Experiment. Math. **22** (2013), 195–202.

[17] J. Buzek, J. Hasan, J. Liu, M. Naehrig, A. Vigil, *Finding twin smooth integers by solving Pell equations*, arXiv:2211.04315.

[18] B. Sterner, *Towards optimally small smoothness bounds for cryptographic-sized twin smooth integers and their isogeny-based applications*, SAC 2024; ePrint 2023/1576.

[19] E. Mulder, B. Sterner, W. van Woerden, *Large smooth twins from short lattice vectors*, ANTS XVII.

[20] C. Størmer, *Quelques théorèmes sur l'équation de Pell* $x^2 - Dy^2 = \pm 1$ *et leurs applications*, Skr. Norske Vid. Akad., 1897.

[21] D. H. Lehmer, *On a problem of Størmer*, Illinois J. Math. **8** (1964), 57–79.