

Calculus 1B lecture 4

Tugce Akkaya

University of Twente

December 7, 2020

-Contents-

- Integrals

- Calculation techniques for integrals

- Power and Taylor series

- First order ODEs

- Complex numbers

- Second order ODEs

Differential equations – Introduction

Separable DEs (Thomas 7.2)

Direction Fields (Thomas 9.1)

Integrating factor (Thomas 9.2)

Definition

A first-order differential equation (DE) is an equation involving an unknown function together with its derivative¹.

¹First order: f' , second order f'' , ...

Definition

A first-order differential equation (DE) is an equation involving an unknown function together with its derivative¹.

- ▶ Example: $y' - y = x$. (1)

¹First order: f' , second order f'' , ...

Definition

A first-order differential equation (DE) is an equation involving an unknown function together with its derivative¹.

- ▶ Example: $y' - y = x$. (1)

Definition

A solution is a function that, whenever substituted, satisfies the equation.

¹First order: f' , second order f'' , ...

Definition

A first-order differential equation (DE) is an equation involving an unknown function together with its derivative¹.

- ▶ Example: $y' - y = x$. (1)

Definition

A solution is a function that, whenever substituted, satisfies the equation.

- ▶ The function $y(x) = e^x - x - 1$ satisfies (1):

$$y'(x) = e^x - 1,$$

consequently

$$y'(x) - y(x) = (e^x - 1) - (e^x - x - 1)$$

¹First order: f' , second order f'' , ...

Definition

A first-order differential equation (DE) is an equation involving an unknown function together with its derivative¹.

- ▶ Example: $y' - y = x.$ (1)

Definition

A solution is a function that, whenever substituted, satisfies the equation.

- ▶ The function $y(x) = e^x - x - 1$ satisfies (1):

$$y'(x) = e^x - 1,$$

consequently

$$\begin{aligned}y'(x) - y(x) &= (e^x - 1) - (e^x - x - 1) \\&= x.\end{aligned}$$

¹First order: f' , second order f'' , ...

Definition

An ordinary first-order differential equation (ODE) is a differential equation where y' can be expressed as a function of y and x .

$$y' = \text{cloud}$$

The cloud contains the letters y , x , and x arranged in a triangular pattern:

- y is at the top left.
- x is below y and to the right.
- x is below the first x and to the right.
- y is below the second x and to the right.

- ▶ The formal way to denote a normal equation is

$$y' = f(x, y).$$

Definition

An ordinary first-order differential equation (ODE) is a differential equation where y' can be expressed as a function of y and x .

$$y' = \begin{matrix} & y & x & x \\ & x & & y \end{matrix}$$

- ▶ The formal way to denote a normal equation is

$$y' = f(x, y).$$

Example

The equation $y' - y = x$ can be rewritten as a normal equation:


$$y' = y + x. \tag{2}$$

Hence we have: $f(x, y) = y + x.$

Differential equations - Introduction Conceived by?

4

Differential equations first appeared in the work of Newton and Leibniz, ±1672.




They were first constructed to model physics, but are nowadays used to model almost everything, stock prices, glucose concentration in the body, traffic flows, etc.

Differential equations - Introduction Conceived by?


4

Differential equations first appeared in the work of Newton and Leibniz, ±1672.




They were first constructed to model physics, but are nowadays used to model almost everything, stock prices, glucose concentration in the body, traffic flows, etc.

Next we derive a differential equation modelling the making of a mug of coffee via filtration.




Const	description	units
F_0	initial number of coffee particles in the filter	-
k	coffee filtration constant	s^{-1}
Var	description	units
t	time	s
$F(t)$	nr of coffee particles in filter	-
$N(t)$	nr of coffee particles in mug	-



- ▶ The number a of coffee particles that leaves the filter during a short period of time Δt is proportional to the number of coffee particles F in the filter and to the length of the period Δt :

$$a \propto F\Delta t.$$




- ▶ The number a of coffee particles that leaves the filter during a short period of time Δt is proportional to the number of coffee particles F in the filter and to the length of the period Δt :


$$a \propto F\Delta t.$$

- ▶ The coffee filtration constant k is defined as the fraction particles that trickle through the filter per second, hence


$$a = kF\Delta t.$$



- ▶ The number of coffee particles that enters the mug between time t and $t + \Delta t$ is $N(t + \Delta t) - N(t)$.



- ▶ The number of coffee particles that enters the mug between time t and $t + \Delta t$ is $N(t + \Delta t) - N(t)$.
- ▶ The number of coffee particles in the filter at time t is $F(t) = F_0 - N(t)$, so



- ▶ The number of coffee particles that enters the mug between time t and $t + \Delta t$ is $N(t + \Delta t) - N(t)$.
- ▶ The number of coffee particles in the filter at time t is $F(t) = F_0 - N(t)$, so
- ▶ The number a of coffee particles that leaves the filter during a period of Δt seconds is approximately $a = k(F_0 - N(t))\Delta t$.



- ▶ No coffee is wasted or added, so

$$N(t + \Delta t) - N(t) \approx k(F_0 - N(t))\Delta t.$$



- ▶ No coffee is wasted or added, so

$$N(t + \Delta t) - N(t) \approx k(F_0 - N(t))\Delta t.$$

- ▶ Divide left and right hand side by Δt :

$$\frac{N(t + \Delta t) - N(t)}{\Delta t} \approx k(F_0 - N(t)).$$



- ▶ No coffee is wasted or added, so

$$N(t + \Delta t) - N(t) \approx k(F_0 - N(t))\Delta t.$$

- ▶ Divide left and right hand side by Δt :

$$\frac{N(t + \Delta t) - N(t)}{\Delta t} \approx k(F_0 - N(t)).$$

- ▶ The approximation becomes better as Δt gets smaller:

$$N'(t) = \lim_{\Delta t \rightarrow 0} \frac{N(t + \Delta t) - N(t)}{\Delta t} = k(F_0 - N(t)).$$



- ▶ No coffee is wasted or added, so

$$N(t + \Delta t) - N(t) \approx k(F_0 - N(t))\Delta t.$$

- ▶ Divide left and right hand side by Δt :

$$\frac{N(t + \Delta t) - N(t)}{\Delta t} \approx k(F_0 - N(t)).$$

- ▶ The approximation becomes better as Δt gets smaller:

$$N'(t) = \lim_{\Delta t \rightarrow 0} \frac{N(t + \Delta t) - N(t)}{\Delta t} = k(F_0 - N(t)).$$

- ▶ The function N satisfies the differential equation

$$N' = k(F_0 - N)$$

Given a differential equation, a natural question is how to find a solution.

We will show three techniques:

- ▶ Separation of variables , a way of calculating the solution;
- ▶ Direction fields , which is graphical construction;
- ▶ Integrating factor , a general way for calculating the solution.

Definition

A quantity y depending on time grows exponentially if $y' \propto y$.

- If y grows exponentially, then y satisfies the differential equation

$$y' = ky \quad (1)$$

for some constant k .

Definition

A quantity y depending on time grows exponentially if $y' \propto y$.

- ▶ If y grows exponentially, then y satisfies the differential equation

$$y' = ky \quad (1)$$

for some constant k .

- ▶ If $k > 0$, then the quantity y grows, and if $k < 0$ then y decays.

Definition

A quantity y depending on time grows exponentially if $y' \propto y$.

- If y grows exponentially, then y satisfies the differential equation

$$y' = ky \quad (1)$$

for some constant k .

- If $k > 0$, then the quantity y grows, and if $k < 0$ then y decays.

Quiz A solution of equation (1) is given by

- (a) $y(x) = k e^x;$
- (b) $y(x) = -5e^{kx};$
- (c) $y(x) = 7e^{-kx};$
- (d) $y(x) = \ln(kx);$

Definition

A quantity y depending on time grows exponentially if $y' \propto y$.

- If y grows exponentially, then y satisfies the differential equation

$$y' = ky \quad (1)$$

for some constant k .

- If $k > 0$, then the quantity y grows, and if $k < 0$ then y decays.

Quiz A solution of equation (1) is given by

- (a) $y(x) = k e^x$;
- (b) $y(x) = -5e^{kx}$;
- (c) $y(x) = 7e^{-kx}$;
- (d) $y(x) = \ln(kx)$;

Answer (b) is correct. Why? Let us solve (1):

Solution

- ▶ $y' = k y \Leftrightarrow \frac{dy}{dx} = k y$

Solution

- ▶ $y' = k y \Leftrightarrow \frac{dy}{dx} = k y$
- ▶ $dy = k y dx \Leftrightarrow \frac{dy}{y} = k dx$

Solution

- ▶ $y' = k y \Leftrightarrow \frac{dy}{dx} = k y$
- ▶ $dy = k y dx \Leftrightarrow \frac{dy}{y} = k dx$
- ▶ $\int \frac{1}{y} dy = \int k dx$

Solution

- ▶ $y' = k y \Leftrightarrow \frac{dy}{dx} = k y$
- ▶ $dy = k y dx \Leftrightarrow \frac{dy}{y} = k dx$
- ▶ $\int \frac{1}{y} dy = \int k dx$
- ▶ $\ln |y| = k x + C_1$

Solution

- ▶ $y' = k y \Leftrightarrow \frac{dy}{dx} = k y$
- ▶ $dy = k y dx \Leftrightarrow \frac{dy}{y} = k dx$
- ▶ $\int \frac{1}{y} dy = \int k dx$
- ▶ $\ln |y| = k x + C_1$
- ▶ $|y| = e^{k x + C_1}$

Solution

- ▶ $y' = k y \Leftrightarrow \frac{dy}{dx} = k y$
- ▶ $dy = k y dx \Leftrightarrow \frac{dy}{y} = k dx$
- ▶ $\int \frac{1}{y} dy = \int k dx$
- ▶ $\ln |y| = k x + C_1$
- ▶ $|y| = e^{kx+C_1}$
- ▶ $y = \pm e^{C_1} e^{kx}$

Solution

$$\blacktriangleright y' = k y \Leftrightarrow \frac{dy}{dx} = k y$$

$$\blacktriangleright dy = k y dx \Leftrightarrow \frac{dy}{y} = k dx$$

$$\blacktriangleright \int \frac{1}{y} dy = \int k dx$$

$$\blacktriangleright \ln |y| = k x + C_1$$

$$\blacktriangleright |y| = e^{kx+C_1}$$

$$\blacktriangleright y = \pm e^{C_1} e^{kx}$$

$$\blacktriangleright y = C e^{kx} \text{ with } C \in \mathbb{R}$$

Indeed exponential change!

Examples of exponential growth are:

- the human population
(annual growth percentage is approximately 1.14%);
- duckweed growth in a pond;
- carbon dating
(the half-life of ^{14}C is approximately 5730 years);
- compound interest;
- Moore's law: the number of transistors on integrated circuits doubles approximately every two years.

The way of solving the differential equation for exponential change is an example of a more general method, applicable to differential equations that are **separable**:

Definition

A DE $y' = f(x, y)$ is **separable** if it can be written in the form

$$y' = g(x)h_1(y)$$

in which g is a function of x and h_1 is a function of y .

The solution of a **separable** differential equation can be found by the following steps:

- ▶ Let $y' = g(x)h_1(y)$ be written as $\frac{dy}{dx} = g(x)\frac{1}{h(y)}$.

The solution of a **separable** differential equation can be found by the following steps:

- ▶ Let $y' = g(x)h_1(y)$ be written as $\frac{dy}{dx} = g(x)\frac{1}{h(y)}$.
- ▶ Slightly rewriting: $\frac{dy}{dx} = \frac{g(x)}{h(y)}$.

The solution of a **separable** differential equation can be found by the following steps:

- ▶ Let $y' = g(x)h_1(y)$ be written as $\frac{dy}{dx} = g(x)\frac{1}{h(y)}$.
- ▶ Slightly rewriting: $\frac{dy}{dx} = \frac{g(x)}{h(y)}$.
- ▶ Then, cross multiplying gives $h(y) dy = g(x) dx$, and finally

The solution of a **separable** differential equation can be found by the following steps:

- ▶ Let $y' = g(x)h_1(y)$ be written as $\frac{dy}{dx} = g(x)\frac{1}{h(y)}$.
- ▶ Slightly rewriting: $\frac{dy}{dx} = \frac{g(x)}{h(y)}$.
- ▶ Then, cross multiplying gives $h(y) dy = g(x) dx$, and finally
- ▶ $\int h(y) dy = \int g(x) dx$.

The solution of a **separable** differential equation can be found by the following steps:

- ▶ Let $y' = g(x)h_1(y)$ be written as $\frac{dy}{dx} = g(x)\frac{1}{h(y)}$.
- ▶ Slightly rewriting: $\frac{dy}{dx} = \frac{g(x)}{h(y)}$.
- ▶ Then, cross multiplying gives $h(y) dy = g(x) dx$, and finally
- ▶ $\int h(y) dy = \int g(x) dx$.
- ▶ With anti-derivatives H and G , we have $H(y) = G(x) + C$, $C \in \mathbb{R}$.

The solution of a **separable** differential equation can be found by the following steps:

- ▶ Let $y' = g(x)h_1(y)$ be written as $\frac{dy}{dx} = g(x)\frac{1}{h(y)}$.
- ▶ Slightly rewriting: $\frac{dy}{dx} = \frac{g(x)}{h(y)}$.
- ▶ Then, cross multiplying gives $h(y) dy = g(x) dx$, and finally
- ▶ $\int h(y) dy = \int g(x) dx$.
- ▶ With anti-derivatives H and G , we have $H(y) = G(x) + C$, $C \in \mathbb{R}$.
- ▶ In which y is implicitly defined as a function of x . So

The solution of a **separable** differential equation can be found by the following steps:

- ▶ Let $y' = g(x)h_1(y)$ be written as $\frac{dy}{dx} = g(x)\frac{1}{h(y)}$.
- ▶ Slightly rewriting: $\frac{dy}{dx} = \frac{g(x)}{h(y)}$.
- ▶ Then, cross multiplying gives $h(y) dy = g(x) dx$, and finally
- ▶ $\int h(y) dy = \int g(x) dx$.
- ▶ With anti-derivatives H and G , we have $H(y) = G(x) + C$, $C \in \mathbb{R}$.
- ▶ In which y is implicitly defined as a function of x . So
- ▶ In case H is invertible $y = H^{-1}(G(x) + C)$.

Consider the differential equation²

$$\frac{dy}{dx} + P(x) y = 0$$

It can be solved applying the technique of the previous slide:

- ▶ Write as $\frac{dy}{dx} = -P(x) y$

²Thomas, end of page. 525.

Consider the differential equation²

$$\frac{dy}{dx} + P(x) y = 0$$

It can be solved applying the technique of the previous slide:

- ▶ Write as $\frac{dy}{dx} = -P(x) y$
- ▶ So $\frac{dy}{y} = -P(x) dx$

²Thomas, end of page. 525.

Consider the differential equation²

$$\frac{dy}{dx} + P(x) y = 0$$

It can be solved applying the technique of the previous slide:

- ▶ Write as $\frac{dy}{dx} = -P(x) y$
- ▶ So $\frac{dy}{y} = -P(x) dx$
- ▶ Integrating both sides gives $\ln |y| = - \int P(x) dx + C_1$

²Thomas, end of page. 525.

Consider the differential equation²

$$\frac{dy}{dx} + P(x) y = 0$$

It can be solved applying the technique of the previous slide:

- ▶ Write as $\frac{dy}{dx} = -P(x) y$
- ▶ So $\frac{dy}{y} = -P(x) dx$
- ▶ Integrating both sides gives $\ln |y| = - \int P(x) dx + C_1$
- ▶ $|y| = e^{- \int P(x) dx + C_1}$

²Thomas, end of page. 525.

Consider the differential equation²

$$\frac{dy}{dx} + P(x) y = 0$$

It can be solved applying the technique of the previous slide:

- ▶ Write as $\frac{dy}{dx} = -P(x) y$
- ▶ So $\frac{dy}{y} = -P(x) dx$
- ▶ Integrating both sides gives $\ln |y| = - \int P(x) dx + C_1$
- ▶ $|y| = e^{- \int P(x) dx + C_1}$
- ▶ $y = \pm e^{C_1} e^{- \int P(x) dx}$

²Thomas, end of page. 525.

Consider the differential equation²

$$\frac{dy}{dx} + P(x) y = 0$$

It can be solved applying the technique of the previous slide:



- ▶ Write as $\frac{dy}{dx} = -P(x) y$
- ▶ So $\frac{dy}{y} = -P(x) dx$
- ▶ Integrating both sides gives $\ln |y| = - \int P(x) dx + C_1$
- ▶ $|y| = e^{- \int P(x) dx + C_1}$
- ▶ $y = \pm e^{C_1} e^{- \int P(x) dx}$
- ▶ $y = C e^{- \int P(x) dx}$ with $C \in \mathbb{R}$.

Solved!

UNIVERSITY OF TWENTE.


²Thomas, end of page. 525.

Let's have a break!



Direction Fields (Thomas 9.1) General idea

16




$y(x)$ is solution of $y' = f(x, y)$
and $y_0 = y(x_0)$

TE.

Direction Fields (Thomas 9.1) General idea

16




$y(x)$ is solution of $y' = f(x, y)$
and $y_0 = y(x_0)$

slope of ℓ is $y'(x_0) = f(x_0, y_0)$

Direction Fields (Thomas 9.1) General idea

16




$y(x)$ is solution of $y' = f(x, y)$
and $y_0 = y(x_0)$

slope of ℓ is $y'(x_0) = f(x_0, y_0)$

Direction Fields (Thomas 9.1) General idea

16




$y(x)$ is solution of $y' = f(x, y)$
and $y_0 = y(x_0)$

slope of ℓ is $y'(x_0) = f(x_0, y_0)$

Direction Fields (Thomas 9.1) General idea

16




$y(x)$ is solution of $y' = f(x, y)$
and $y_0 = y(x_0)$

slope of ℓ is $y'(x_0) = f(x_0, y_0)$

TE.

Direction Fields (Thomas 9.1) General idea


17



$$y' = y + x$$

Direction Fields (Thomas 9.1) General idea

17




$$y' = y + x$$

$$y_1(0) = 1$$

Direction Fields (Thomas 9.1) General idea

17




$$y' = y + x$$

$$y_1(0) = 1$$


$$y_2(0) = .25$$

Direction Fields (Thomas 9.1) General idea

18




$$y' = y - x^2$$



$$y' = -\frac{2xy}{x^2 + 1}$$


Direction Fields (Thomas 9.1) General idea

18



$$y' = y - x^2$$

$$y(x) = x^2 + 2x + 2 + Ce^x$$



$$y' = -\frac{2xy}{x^2 + 1}$$




$$y(x) = \frac{C}{x^2 + 1}$$

- In general a differential equation has infinitely many solutions.

Definition


An initial value problem consists of a differential equation and an equation of the form $y(a) = b$.

- An initial value problem in general has one solution.




Direction Fields (Thomas 9.1) Exponential change

20



$$y' = ky$$

$k > 0$




$k < 0$

- Let $y(t)$ be a solution. Define $Y(t) = y(t)e^{-kt}$, then

$$Y' = y'e^{-kt} - kye^{-kt}$$


Direction Fields (Thomas 9.1) Exponential change

20



$$k > 0$$

$$y' = ky$$



$$k < 0$$


- Let $y(t)$ be a solution. Define $Y(t) = y(t)e^{-kt}$, then

$$\begin{aligned} Y' &= y'e^{-kt} - kye^{-kt} \\ &= (y' - ky)e^{-kt} = 0, \end{aligned}$$


hence $Y(t)$ is constant. So $Y(t) = y(t)e^{-kt} = C$, and ..

Direction Fields (Thomas 9.1) Exponential change

20



$$y' = ky$$

 $k > 0$  $k < 0$

- Let $y(t)$ be a solution. Define $Y(t) = y(t)e^{-kt}$, then


$$\begin{aligned} Y' &= y'e^{-kt} - kye^{-kt} \\ &= (y' - ky)e^{-kt} = 0, \end{aligned}$$

hence $Y(t)$ is constant. So $Y(t) = y(t)e^{-kt} = C$, and ..

- The general solution of $y' = ky$ is $y(t) = Ce^{kt}$ with $C \in \mathbb{R}$.

Direction Fields (Thomas 9.1) Exponential change

21



Exponential growth with an initial condition

The (unique) solution of the initial value problem

$$\begin{cases} y' = ky, \\ y(0) = y_0 \end{cases}$$

is $y(t) = y_0 e^{kt}$.

- We often write $y(t) = y(0)e^{kt}$.



- ▶ The coffee problem is described by the initial value problem

$$\begin{cases} N' = k(F_0 - N) \\ N(0) = 0. \end{cases}$$



- ▶ The coffee problem is described by the initial value problem

$$\begin{cases} N' = k(F_0 - N) \\ N(0) = 0. \end{cases}$$

- ▶ The number of coffee particles in the filter at time t is described by

$$F(t) = F_0 - N(t).$$



- ▶ The coffee problem is described by the initial value problem

$$\begin{cases} N' = k(F_0 - N) \\ N(0) = 0. \end{cases}$$

- ▶ The number of coffee particles in the filter at time t is described by

$$F(t) = F_0 - N(t).$$

- ▶ The quantity F is exponentially decaying:

$$\begin{cases} F' = -N' = -k(F_0 - N) = -kF \\ F(0) = F_0 - N(0) = F_0, \end{cases}$$

therefore $F(t) = F_0 e^{-kt}$.



- ▶ The coffee problem is described by the initial value problem

$$\begin{cases} N' = k(F_0 - N) \\ N(0) = 0. \end{cases}$$

- ▶ The number of coffee particles in the filter at time t is described by

$$F(t) = F_0 - N(t).$$


- ▶ The quantity F is exponentially decaying:

$$\begin{cases} F' = -N' = -k(F_0 - N) = -kF \\ F(0) = F_0 - N(0) = F_0, \end{cases}$$

therefore $F(t) = F_0 e^{-kt}$.

- ▶

$$N(t) = F_0(1 - e^{-kt})$$



- ▶ The number of coffee particles in the mug at time t is

$$N(t) = F_0(1 - e^{-kt}), \quad t \geq 0.$$

- ▶ For large t almost all coffee particles will be extracted from the filter into the brew.

Definition

A linear first-order equation is a differential equation of the form

$$y' + P(x)y = Q(x). \quad (*)$$

Definition

A linear first-order equation is a differential equation of the form

$$y' + P(x)y = Q(x). \quad (*)$$

- ▶ Multiply (*) with a positive function $v(x)$:

$$vy' + vP y = vQ$$

Definition

A linear first-order equation is a differential equation of the form

$$y' + P(x)y = Q(x). \quad (*)$$

- ▶ Multiply (*) with a positive function $v(x)$:

$$vy' + vP y = vQ$$

- ▶ Product rule: let $v(x)$ be an arbitrary function, then

$$(v y)' = vy' + v' y$$

Definition

A linear first-order equation is a differential equation of the form

$$y' + P(x)y = Q(x). \quad (*)$$

- Multiply (*) with a positive function $v(x)$:

$$vy' + vP y = vQ$$

- Product rule: let $v(x)$ be an arbitrary function, then

$$(v y)' = vy' + v' y$$

- If $v' = vP$ then $(vy)' = vQ$.

Consequently vy (and therefore y) can be obtained by integrating vQ .

- ▶ Rewrite $v' = vP$ to

$$\frac{v'}{v} = P(x).$$

³The constant of integration was taken 0, the anti-derivative is as simple as possible.

- ▶ Rewrite $v' = vP$ to

$$\frac{v'}{v} = P(x).$$

- ▶ We have

$$\ln v(x) = \int P(x) \, dx$$

where $\int P(x) \, dx$ is an antiderivative of $P(x)$.

³The constant of integration was taken 0, the anti-derivative is as simple as possible.

- ▶ Rewrite $v' = vP$ to

$$\frac{v'}{v} = P(x).$$

- ▶ We have

$$\ln v(x) = \int P(x) \, dx$$

where $\int P(x) \, dx$ is an antiderivative of $P(x)$.

- ▶

$$v(x) = e^{\int P(x) \, dx}.$$

The function $v(x)$ is called the **integrating factor**³.

³The constant of integration was taken 0, the anti-derivative is as simple as possible.

Integrating factor (Thomas 9.2) Example

Solve the differential equation $xy' - y = x^2 e^x$.

write in standard form	$y' + P y = Q$	
find the integrating factor $v(x)$	$v = e^{\int P(x) dx}$	
multiply with v	$vy' + vP y = vQ$	
use $v' = vP$	$vy' + v'y = vQ$	
product rule	$(vy)' = vQ$	
integrate	$vy = \int v(x)Q(x) dx$	
divide by v	$y = \frac{1}{v} \int v(x)Q(x) dx$	

Integrating factor (Thomas 9.2) Example

26

Solve the differential equation $xy' - y = x^2 e^x$.

write in standard form	$y' + Py = Q$	$y' - \frac{1}{x}y = x e^x$	$P(x) = -\frac{1}{x}$ $Q(x) = x e^x$
find the integrating factor $v(x)$	$v = e^{\int P(x) dx}$		
multiply with v	$vy' + vPy = vQ$		
use $v' = vP$	$vy' + v'y = vQ$		
product rule	$(vy)' = vQ$		
integrate	$vy = \int v(x)Q(x) dx$		
divide by v	$y = \frac{1}{v} \int v(x)Q(x) dx$		

Integrating factor (Thomas 9.2) Example

26

Solve the differential equation $xy' - y = x^2 e^x$.

write in standard form	$y' + Py = Q$	$y' - \frac{1}{x}y = x e^x$	$P(x) = -\frac{1}{x}$ $Q(x) = x e^x$
find the integrating factor $v(x)$	$v = e^{\int P(x) dx}$	$\int -\frac{1}{x} dx = -\ln x:$ $v(x) = e^{-\ln x} = x^{-1} = \frac{1}{x}$	
multiply with v	$vy' + vPy = vQ$		
use $v' = vP$	$vy' + v'y = vQ$		
product rule	$(vy)' = vQ$		
integrate	$vy = \int v(x)Q(x) dx$		
divide by v	$y = \frac{1}{v} \int v(x)Q(x) dx$		

Integrating factor (Thomas 9.2) Example

Solve the differential equation $xy' - y = x^2 e^x$.

write in standard form	$y' + Py = Q$	$y' - \frac{1}{x}y = x e^x$	$P(x) = -\frac{1}{x}$ $Q(x) = x e^x$
find the integrating factor $v(x)$	$v = e^{\int P(x) dx}$	$\int -\frac{1}{x} dx = -\ln x:$ $v(x) = e^{-\ln x} = x^{-1} = \frac{1}{x}$	
multiply with v	$vy' + vPy = vQ$	$\frac{1}{x}y' - \frac{1}{x^2}y = e^x$	
use $v' = vP$	$vy' + v'y = vQ$		
product rule	$(vy)' = vQ$		
integrate	$vy = \int v(x)Q(x) dx$		
divide by v	$y = \frac{1}{v} \int v(x)Q(x) dx$		

Integrating factor (Thomas 9.2) Example

Solve the differential equation $xy' - y = x^2 e^x$.

write in standard form	$y' + Py = Q$	$y' - \frac{1}{x}y = x e^x$	$P(x) = -\frac{1}{x}$ $Q(x) = x e^x$
find the integrating factor $v(x)$	$v = e^{\int P(x) dx}$	$\int -\frac{1}{x} dx = -\ln x:$ $v(x) = e^{-\ln x} = x^{-1} = \frac{1}{x}$	
multiply with v	$vy' + vPy = vQ$	$\frac{1}{x}y' - \frac{1}{x^2}y = e^x$	
use $v' = vP$	$vy' + v'y = vQ$	$\frac{1}{x}y' + \frac{d}{dx}\left(\frac{1}{x}\right)y = e^x$	
product rule	$(vy)' = vQ$		
integrate	$vy = \int v(x)Q(x) dx$		
divide by v	$y = \frac{1}{v} \int v(x)Q(x) dx$		

Integrating factor (Thomas 9.2) Example

Solve the differential equation $xy' - y = x^2 e^x$.

write in standard form	$y' + Py = Q$	$y' - \frac{1}{x}y = x e^x$	$P(x) = -\frac{1}{x}$ $Q(x) = x e^x$
find the integrating factor $v(x)$	$v = e^{\int P(x) dx}$	$\int -\frac{1}{x} dx = -\ln x:$ $v(x) = e^{-\ln x} = x^{-1} = \frac{1}{x}$	
multiply with v	$vy' + vPy = vQ$	$\frac{1}{x}y' - \frac{1}{x^2}y = e^x$	
use $v' = vP$	$vy' + v'y = vQ$	$\frac{1}{x}y' + \frac{d}{dx}\left(\frac{1}{x}\right)y = e^x$	
product rule	$(vy)' = vQ$	$\frac{d}{dx}\left(\frac{1}{x}y\right) = e^x$	
integrate	$vy = \int v(x)Q(x) dx$		
divide by v	$y = \frac{1}{v} \int v(x)Q(x) dx$		

Integrating factor (Thomas 9.2) Example

26

Solve the differential equation $xy' - y = x^2 e^x$.

write in standard form	$y' + Py = Q$	$y' - \frac{1}{x}y = x e^x$	$P(x) = -\frac{1}{x}$ $Q(x) = x e^x$
find the integrating factor $v(x)$	$v = e^{\int P(x) dx}$	$\int -\frac{1}{x} dx = -\ln x:$ $v(x) = e^{-\ln x} = x^{-1} = \frac{1}{x}$	
multiply with v	$vy' + vPy = vQ$	$\frac{1}{x}y' - \frac{1}{x^2}y = e^x$	
use $v' = vP$	$vy' + v'y = vQ$	$\frac{1}{x}y' + \frac{d}{dx}\left(\frac{1}{x}\right)y = e^x$	
product rule	$(vy)' = vQ$	$\frac{d}{dx}\left(\frac{1}{x}y\right) = e^x$	
integrate	$vy = \int v(x)Q(x) dx$	$\frac{1}{x}y = \int e^x dx = e^x + C$	
divide by v	$y = \frac{1}{v} \int v(x)Q(x) dx$		UNIVERSITY OF TWENTE.

Integrating factor (Thomas 9.2) Example

26

Solve the differential equation $xy' - y = x^2 e^x$.

write in standard form	$y' + Py = Q$	$y' - \frac{1}{x}y = x e^x$	$P(x) = -\frac{1}{x}$ $Q(x) = x e^x$
find the integrating factor $v(x)$	$v = e^{\int P(x) dx}$	$\int -\frac{1}{x} dx = -\ln x:$ $v(x) = e^{-\ln x} = x^{-1} = \frac{1}{x}$	
multiply with v	$vy' + vPy = vQ$	$\frac{1}{x}y' - \frac{1}{x^2}y = e^x$	
use $v' = vP$	$vy' + v'y = vQ$	$\frac{1}{x}y' + \frac{d}{dx}\left(\frac{1}{x}\right)y = e^x$	
product rule	$(vy)' = vQ$	$\frac{d}{dx}\left(\frac{1}{x}y\right) = e^x$	
integrate	$vy = \int v(x)Q(x) dx$	$\frac{1}{x}y = \int e^x dx = e^x + C$	
divide by v	$y = \frac{1}{v} \int v(x)Q(x) dx$	$y(x) = x(e^x + C)$	UNIVERSITY OF TWENTE.

Summarizing Exercises - 1

See slide #18: Given the DEs

$$y' = -\frac{2xy}{x^2 + 1} \qquad y' = y - x^2$$

Are they

- first order?
- Separable?
- Of the form $dy/dx + P(x)y = 0$?
- Of the form $dy/dx + P(x)y = Q(x)$?

Summarizing Exercises - 2

Solve

$$xy' + \frac{y}{x} = 0 \quad ; \quad y(1) = 1$$

Solve

$$xy' + \frac{y}{x} = e^{1/x}$$

-Contents-

- ☐ Integrals

- ☐ Calculation techniques for integrals

- ☐ Power and Taylor series

- ☐ First order ODEs

- ☐ Complex numbers

- ☐ Second order ODEs