Supporting information: recovery, screening and trend inference

Author information to be completed by the authors

1 Repetition and inference units

The earlier single-year and assembly benchmark fixes twenty integer seeds, 128 conditional trajectories, twenty multiple imputations and 200 trend degradations. The new whole-archive benchmark instead uses twenty paired archive masks per station and 256 joint draws; its separate results and calibration support are reported below. An outer region and every connected independence group are excluded before forming fitting roles. The remaining groups are assigned in shares 0.55/0.15/0.15/0.15 to estimation, interval calibration, failure learning and rule calibration; probability calibration is additionally separated inside failure learning. Bootstrap multiplicities stay inside roles. The source workflow requires at least four complete historical years. Case identity includes network, station, year, mask family, realization and reconstruction tier. The realization string alone is not unique.

Empirical repetition quantiles do not estimate uncertainty from new climates. The complete source data are the same in each repetition. Historical registered verdicts and all opened-data supplementary results retain their original status. No recalibration repairs a historical registered failure retrospectively.

There are 581 physical station identities but 602 network–station input instances, and 6,170 unique complete station-years versus 6,334 network–station–year tasks. Repeated physical records share their connected-group exclusion. Input identity therefore includes the network and its original time window. The observed-air control reconstructs those full-precision panels and original forcing, rather than using a clipped float32 archive proxy.

For AR(1) correlation ϕ and integer $m > 1$, the recovery model uses

$$v(m,\phi) = m^{-2}\{m + 2\sum_{j=1}^{m-1}(m-j)\phi^j\}, \qquad c(m,\phi) = (30m)^{-1}\sum_{i=1}^{m}\sum_{j=1}^{30}\phi^{|i-j|}.$$

The hidden interval in this covariance approximation is contiguous at one end of the window; correlations are clipped to $[-0.99, 0.999]$. The variance formula uses

rounded hidden-day counts, and zero hidden share has zero component mean and variance before the numerical scale floor. These approximations do not constitute a distribution-free recovery bound. The stored nominal mixture scale combines untruncated component variances with truncated means; it is a diagnostic scale proxy, not the exact truncated-mixture variance. The Student sensitivity must be interpreted accordingly.

2 Numerical scores and censoring

For ensemble $x_1, \ldots, x_M$ and outcome y, CRPS is

$$M^{-1} \sum_m |x_m - y| - (2M^2)^{-1} \sum_{m,l} |x_m - x_l|.$$

Threshold-weighted scoring replaces both arguments by their maximum with the training H30 threshold, without selecting outcomes. Seven quantile levels, 0.05/0.10/0.25/0.50/0.75/0.90/0.95, use midpoint quadrature with explicit constant outer tails. Residual-bridge distributions use the empirical ensemble directly. Classical quantile corrections receive the same disjoint calibration roles and weights as distribution regression. The 90% interval score adds twenty times the distance outside the interval to its width. Deterministic methods have no distribution score.

With an annual observation known only to exceed L, the Gaussian trend contribution is $\log \Phi\{(\mu - L)/\sigma\}$. It is a survival likelihood for a lower bound. Such bounds can be informative because the observation window and annual maximum are related; this working model is evaluated as a comparator, not assumed correct. Entirely censored series and optimizer failures are unavailable, not exact trends.

3 Comparisons, availability and subgroup reporting

Point and distribution scorecards cross empirical calendars and controlled gaps with decision-time peak overlap and tier. Air-indicated hot conditions use training-role air quantiles. True heat and true peak overlap appear only as evaluation diagnostics. Pairwise station and group effect summaries require common available predictions. Risk tables retain total, admitted and evaluable-group counts. Unavailable scores are rejected; no admitted observations imply undefined selected risk.

Three neural splits contain the same held-case masks across MSE boosting, LSTM, SAITS and CSDI. Neural inputs are water anomalies, air anomalies and seasonal sine/cosine; neighbor series are omitted. Models use a 30-epoch CPU cap and early stopping on at most three evenly spaced validation years per group. Full case-weighted development runs were stopped for computational cost before CSDI test prediction; all final arms were retrained using the fixed group-balanced rule. CSDI computes H30 separately for twenty sampled trajectories. Matched annual-model and classical-baseline files use exactly these case identities and validation groups. Availability, including missing discharge, is determined before computing failure rates. Medians across three splits are descriptive and cannot satisfy the 18-of-20 abstract rule.

Individual caps and batch-average rules answer different questions. With within-group admitted and failed fractions A_g, B_g, the bounded-group certificate targets $E(B_g)/E(A_g)$ for uniform-group sampling. It bounds $B_g + r(1 - A_g)$ at r using independent bounded group units and a fixed candidate family. It provides no guarantee for every group and does not target the case-weighted ratio reported in ordinary scorecards. The eight-threshold family uses a Bonferroni allocation of the error budget. Even zero observed bounded loss requires approximately 99 groups for a 5% cap and 5% total error; a single threshold requires about 59. Rejecting every value provides no nonempty selected-risk evidence.

Overlapping-group adjustments, random-effects intervals, EM prior changes, domain weights, ATC and prediction-powered mean-risk estimates have their own unverified assumptions. The O/E analysis uses a group-unit sampling-variance floor rather than treating dependent simulated masks as independent events. The PPI correction estimates a group-equal mean with an asymptotic interval; it is not a finite-sample semi-supervised selective-risk theorem. Synthetic label-shift controls change class priors while preserving class-conditional feature distributions; separate controls change gap, climate and conditional risk. The local-label curve reports nonempty certificates separately from realized risk.

4 Formal conditional and semi-supervised comparisons

The additional conditional-conformal implementation uses the authors' `conditionalconformal` 0.0.4 package and the binary LAC score $S(x, y) = |y - p(x)|$. Calibration samples one case uniformly per unique rule-calibration independence group, outside every held group. The basis contains overlapping macro-region, information-tier, peak-overlap, historical-heat, dam and zero-flow proxy indicators. The source rule-calibration role is distinct from the estimation, interval and failure-learning roles. Mask replicas are not additional calibration units.

Both the conservative and independently case-randomized versions are evaluated. The latter draws a test dual weight $u \sim \mathrm{Unif}(\tau - 1, \tau)$ separately for every case. The finite-basis boundary is computed by minimizing the calibration pinball loss tilted by $-u\Phi(x)^\top \beta$, then selecting the smallest target prediction among optimal fits. Mathematical unboundedness gives a full or empty prediction set according to the sign of the test weight. The boundary LP agrees with the authors' sensitivity algorithm in numerical tests, including random dual weights. Monotone cutoff caching retains casewise randomization and enlarges a set by at most the stated 10^{-8} numerical endpoint tolerance. Solver failures conservatively return the full binary set and are counted explicitly. An initial one-sided-residual development comparison is archived; the final binary-score results are separate.

For the nonrandomized version, every active binary group indicator requires at least $\tau/(1 - \tau)$ independent calibration representatives for a finite cutoff: nine at 90% and nineteen at 95%. This follows directly from the feasible range of the calibration dual sum for that group. It is a mathematical support limit, not a failed optimizer. The randomized version has a different support condition and is not suppressed by this nonrandomized count test. A previously unseen full region indicator permits only a full or empty randomized set, with no singleton release.

Group-conditional prediction-set coverage concerns $P(Y \in C(X) \mid G)$. Publishing only when $C = \{0\}$ therefore bounds false-admission frequency, not $P(Y = 1 \mid \text{published}, G)$. Both quantities are reported. The conditional theorem additionally assumes exchangeability of calibration and test representatives; source calibration and a held climatic region need not meet that condition.

The formal general-loss semi-supervised RCPS comparison applies the paper's Algorithms 1 and 4 to $L_g = B_g + r(1 - A_g)$. For distinct equally sized unlabeled blocks U_i,

$$W_i = |U_i|^{-1} \sum_{j \in U_i} \widetilde{L}_j + L_i - \widetilde{L}_i \in [-1, 2].$$

Finite-sample WSR bounds use predictable bets and stop a fixed testing sequence at its first failed candidate. Labeled, unlabeled-calibration and untouched test groups are disjoint. Too few distinct unlabeled groups is an explicit status. Labeled-only WSR and the earlier bounded-loss method remain separate comparators. Their target is a same-population group-equal selected-risk ratio under independent identically distributed group sampling, rather than every river's conditional risk.

5 Frozen-output deletion sensitivity and observed record checks

Group-deletion sensitivity freezes predictions and original admissions. Risk crossings reduce to additive integer margins $db_g - na_g$ for rational cap n/d. Retention reversals solve a binary integer program requiring that both remaining rules satisfy the same point-risk cap and the reference remains nonempty. The published comparator is selected once on opened full-data outcomes. A zero deletion count means no initial qualified advantage; impossible reversals are undefined. Source checks cover all twenty seeds and retain witness group lists. Historical N1 reproduces its original 7% point-risk endpoint, not the later proposed 6%/8% E1 criteria. Reapplying the batch rule at a frozen-admission witness is reported separately and has no claimed minimum-deletion optimality.

The observation-only record check fixes 1981–2025 and requires at least fifteen complete observed years at a station. It recomputes every complete reference H30 and every incomplete-year bound from the original daily inputs. A witness uses thirty consecutive finite observations, with dates and all thirty values retained. An observed-window mean above the maximum of all complete-year H30 values disproves the complete-only record attribution for the reported series. It does not require the candidate to be the unique hottest year, nor assert zero sensor error. A 5×10^{-6} degree numerical tolerance removes floating-point ties. Regional sensitivity separately fixes rosters and first-to-last-complete-year spans; common-year and expanded-year rankings distinguish added support from changes among already eligible years.

6 Supplementary evidence and remaining dependencies

The full machine-readable tables, seed-level results, deep training records, forcing provenance, 7DADM qualification, natural-reference exclusion reasons and New Zealand catalog request are repository artifacts. The natural search yields no qualified difficult annual peak-window outage among 180 candidate pair-years. A zero-failure binomial calculation concerns independent gaps, not dependent daily measurements. Twenty-three station-years have median air/donor residual correlation 0.505. Shared primary-sensor error can induce that correlation; it neither identifies separate sensor errors nor establishes three independent error channels. The reference-precision table assumes Gaussian AR1 errors and bounds linear window means; it does not measure logger accuracy.

The prior opened low-flow illustration is retained below. It concerns April–March positive-flow records and annual seven-day minimum discharge, reconstructed through log-flow regression and an annual distribution model. Its source files are `scripts/461_tr2_flow_demo.py` and `results/wrr_target_recovery_20261022/s1_flow`. A separate matched extension reproduces all 134,925 historical cases and adds best-donor MOVE.1, the same single-donor Ridge, and twenty Gaussian AR1 log-flow bridges. These working bridges use other-year in-sample residuals without independent interval calibration. In 33,609 controlled dangerous donor cases, log-scale MAE is 0.090 for single-donor Ridge and 0.106 for MOVE.1; the multidonor bridge gives 0.104 and historical D2 gives 0.075. The variance-preserving methods can change bias direction without improving MAE. Exponentiating a log-flow prediction differs from a conditional mean in original discharge units. This is descriptive prior evidence, not independent confirmation or a claim that every reconstructed extreme shares one empirical bias sign.

Cold-to-hot and calendar splits refit upstream quantities after removing all hot or future years, including from Air2stream, anomaly reference and donor regression training. The annual non-air stress is a deliberately incompatible diagnostic. The separate full daily perturbation adds 0.2°C per decade to all complete targets and donors and repeats upstream fitting under three seeds. These test different assumptions. Daily-year shifts are applied consistently to complete records and empirical mask calendars.

The Daymet comparison across all 319 USGS source sites changes both spatial product and daily averaging convention. Discharge uses the same original air features with log-flow lags; contradictory daily flow records remain unavailable. All 67,316 original POWER control identities and available points are reproduced exactly. Each alternative has its own shared-case reference. The earlier 35-site standardized-Ridge comparison remains a separate sensitivity. Lags now refer to actual calendar days, including missing leap-year dates, and every unavailable case remains explicit. The prior positional-lag results were reconstructed from the archived script and unchanged cached data; their original output bytes were not retained before overwrite. The European acquisition has 16 usable stations of 16 requested, and 4474 paired cases including 458 dangerous air-ridge cases. Nearby GHCN-Daily TAVG matching covers all 319 benchmarkable source USGS sites: 147 have a metadata-eligible station within 25 km, and 122 supply common valid cases. The exact original source control reproduces 67,316 case identities, availability and every available annual point. NOAA daily source S ends at 2400 UTC, while other source conventions vary; this is an observed-input sensitivity, not a perfect riverbank-air or clock-matched oracle. Missing inputs remain in all denominators. Daily metrics use identical dates in both arms. The 7DADM analysis uses approved maxima at 34 sites, without regulatory certification. Historical GAGES-II dam metadata are joined by exact station identifier; zero-flow fractions are proxies and lake-outlet classification is limited to one explicit agency designation. The distance screen fails to flag Big Sandy Creek; observed errors do not determine exclusion.

The selection exercise concerns an assembled panel rather than all USGS sites. Complete-year versus retained-year station eligibility is 98 versus 255 overall and 36 versus 163 in the USGS panel. Same-station slope contrasts, region-group bootstrap intervals, station BH adjustments, synchronized year permutations and circular shifts are reported separately. Estimated incomplete outcomes cannot prove the truth of a changed scientific conclusion. Hot-year hit rates, false alarms and SEDI are preserved even when no valid event denominator exists. The original specification table varies tolerances and decision caps on fixed predictions. A separate twenty-seed, 540-outer-fit H7 pipeline recomputes seven-day peak support, masks, histories, functional draws, MOVE augmentation and annual/risk calibration. Native H7 fields are adapted to the legacy fitting engine after removing the genuine

secondary H30 fields; exported native predictions restore H7 names. The daily-input audit reconstructs sixteen provider/tier combinations directly, and default H30 parity passes 240 cases and 201 fields. Deep models and all tolerance/model combinations are not refitted.

The Greater Wellington intake now contains actual native observations from all nine selected sites. Code 600 remains the specified accepted quality code; the returned legacy quality codes supply no eligible complete-reference hot years under that criterion. Consequently the external qualification denominator is nine and the recovery-accuracy denominator is zero. The study is unregistered. The source model and classical interval corrections were not adapted to obtain favorable target errors. Measurement-code compatibility must be resolved before treating these observations as reference truth.

The unflagged-exposure experiment uses the same source cases, masks and true annual targets in both conditions. Seven visible warm days are replaced by concurrent air readings, using air and visibility alone to locate the episode. About 97.6% of cases support such an episode, with an average 2.3°C daily air-minus-water shift on replaced days. This is an imposed stress severity, not an estimate of field exposure prevalence. Clean source weights and interval offsets remain fixed; the three seeds vary masks and trajectories. Table 27 separates missing-component and total trend coverage, and Table 28 reports unchanged-score release. Censored trends are omitted here because a contaminated visible window is no longer a valid physical lower bound. Unflagged exposure in an otherwise defined water target must not be interpreted as measurable water temperature in a dry bed.

Forcing-specific recovery probabilities are also recomputed from residual histories excluding both the target year and each residual-validation year. All 269,264 USGS arm/case points reproduce their existing controls, and 1,083 independent daily Ridge checks agree within numerical tolerance. In controlled dangerous air-tier cases, raw Gaussian recovery probabilities average 0.405 against observed failure 0.468 for POWER, and 0.365 against 0.457 for POWER plus flow. These have different available supports and failure targets; they do not establish paired calibration improvement or safe transfer. Complete scores and calibration bins are retained in CSV. Annual distribution and release-classifier weights remain unchanged.




Figure 1. Previously opened low-flow illustration, preserved as supplementary context. Its log-flow estimators and incomplete-year practices have their own assumptions and are not the new twenty-seed thermal benchmark.




Figure 2. Opened-source station failures and available metadata. Coordinates are available for all 581 stations. The map supports descriptive failure analysis, not causal river-type attribution.

The two stations specified in the review plan, USGS 13342500 and 06054500, each contribute twenty complete evaluation years; their median shares of random-missingness A1 squared error are 13.0 and 6.1 percent. Historical dam and zero-flow metadata do not identify thermal causation. Named European 2003, 2018 and 2022 examples use common finite cases across all four estimators and preserve event contingency counts, hit/false-alarm rates and undefined SEDI. The 2003 dangerous calendar sample has only 6–22 cases per repetition; median MAE is 0.913 degrees C for D2 and 0.595 for MOVE.1. US illustration years are selected using positive regional air anomalies with at least three complete-water stations, preserving unavailable regions. These opened examples do not establish universal estimator rankings.

The C1 stratification retains the original 336-USGS-site, 6,479-station-year missingness sample. Approved discharge yields 21 sites with recorded zeros, 266 without a recorded zero and 49 with unavailable flow. A station-conditional logistic model adjusts standardized summer air temperature for linear decade trend; 400 independence-group bootstrap repetitions give OR 1.208 [0.882, 1.564] for the zero-record group and 1.058 [0.988, 1.134] for the available-flow group without recorded zeros. Requiring thirty zero days gives 16 sites and OR 1.282 [0.876, 1.889]. Flow-adjusted specifications have smaller available samples and do not establish mediation. Zero flow is a limited historical proxy, and group exchangeability remains assumed.

The national extension fixes one canonical mean-temperature series per catalog site before numerical analysis, without combining sensors. All 1,273 candidates are attempted; 340 retain unresolved eligibility. National CSVs give eligibility bounds, one-degree station-grid occupancy, HUC2 counts, trends, basin-block stability, BH and synchronized field tests. Hot-year methods share a station threshold from the 90th percentile of at least five complete approved 1981–2010 H30 values; insufficient references remain unavailable. Same-roster/span and common-year comparisons separate added support from changed values. Observed windows provide hot-frequency lower bounds and record witnesses without imputation. Conditional imputation ranges are model sensitivities, not validated confidence coverage.

Table 1. scorecard; source tables contain all strata and empirical repetition intervals.

family	estimator	seeds	MAE	CRPS	coverage
calendar	H30 MOVE.1	20	0.992	–	–
calendar	Annual MOVE.1	20	0.550	–	–
calendar	Warm-season MOVE.1	20	0.509	–	–
calendar	Air2stream-3	20	0.568	–	–
calendar	Calibrated state	20	0.465	0.335	0.875
calendar	State simulation	20	0.465	0.337	0.803
calendar	Distribution regression	20	0.451	0.331	0.863
calendar	Functional debiasing	20	0.476	–	–
calendar	Lookup debiasing	20	0.553	–	–
calendar	Observed days	20	4.108	–	–
calendar	Daily plug-in	20	0.564	–	–
calendar	Stochastic regression	20	0.463	0.343	0.723
calendar	Calibrated stochastic	20	0.463	0.338	0.853
calendar	triage_hybrid	20	0.451	–	–
crossed	H30 MOVE.1	20	1.009	–	–
crossed	Annual MOVE.1	20	0.453	–	–
crossed	Warm-season MOVE.1	20	0.441	–	–
crossed	Air2stream-3	20	0.534	–	–
crossed	Calibrated state	20	0.363	0.264	0.898
crossed	State simulation	20	0.363	0.264	0.824

7 Whole-archive supplements

The full MSE cells, interactions, matched estimator contrasts, unavailable cases and task calibration sources are stored in results/whole_archive. Independent source groups, rather than mask repetitions, determine calibration support. Forward task calibration ends before 2010. The four error-direction illustration uses a single synthetic half-degree common shock on thirty annual values; it is an algebraic sensitivity, not new river evidence. The evaluation-only error covariance uses hidden reference values solely for diagnosis and is never a deployment estimator. Gaussian-tail extrapolation leaves the central stored quantiles unchanged but changes the marginal tails, and is reported separately from the fixed-marginal rank comparison.

8 Earlier diagnostic figures

Table 2. scorecard; source tables contain all strata and empirical repetition intervals.

family	estimator	seeds	MAE	CRPS	coverage
crossed	Distribution regression	20	0.337	0.248	0.888
crossed	Functional debiasing	20	0.355	–	–
crossed	Lookup debiasing	20	0.422	–	–
crossed	Observed days	20	1.427	–	–
crossed	Daily plug-in	20	0.428	–	–
crossed	Stochastic regression	20	0.336	0.248	0.753
crossed	Calibrated stochastic	20	0.337	0.247	0.892
crossed	triage_hybrid	20	0.337	–	–




Figure 3. Decision-time risk concentration across peak-overlap bins. Source tables preserve populations, layers and repetition variability.

Table 3. rules; source tables contain all strata and empirical repetition intervals.

family	delivery	rule	seeds	retention	risk	groups >10%
calendar	Distribution regression	all	20	1.000	0.030	0.071
calendar	Distribution regression	atc	20	0.965	0.014	0.006
calendar	Distribution regression	Batch average	20	0.994	0.027	0.033
calendar	Distribution regression	Batch + individual cap	20	0.966	0.014	0.011
calendar	Distribution regression	gap_49	20	0.806	0.005	0.000
calendar	Distribution regression	Group certificate	20	0.000	–	–
calendar	Distribution regression	individual_0.05	20	0.938	0.005	0.000
calendar	Distribution regression	Individual score ≤0.10	20	0.948	0.008	0.000
calendar	Distribution regression	individual_0.2	20	0.959	0.011	0.006
calendar	Distribution regression	individual_0.3	20	0.967	0.014	0.011
calendar	Distribution regression	Probability + group adjustment	20	0.921	0.003	0.000
calendar	Distribution regression	mondrian	20	0.966	0.014	0.011
calendar	Distribution regression	none	20	0.000	–	–
calendar	Distribution regression	one_feature_25	20	0.889	0.003	0.000
calendar	Distribution regression	Peak overlap <50%	20	0.916	0.005	0.000
calendar	Distribution regression	published_Beaufort_full	20	0.645	0.001	0.000
calendar	Distribution regression	published_Beaufort_full_as_written	20	0.641	0.001	0.000
calendar	Distribution regression	published_Dunham_full	20	0.654	0.002	0.000
calendar	Distribution regression	published_Dunham_full_as_written	20	0.655	0.002	0.000
calendar	Distribution regression	published_ETCCDI_15_3	20	0.350	0.000	0.000




Figure 4. Recovery-model discrimination and Brier skill against simple scores. Ranking does not imply calibration.

Table 4. rules; source tables contain all strata and empirical repetition intervals.

family	delivery	rule	seeds	retention	risk	groups >10%
calendar	Distribution regression	published_GSIM_350	20	0.590	0.000	0.000
calendar	Distribution regression	published_NorWest90	20	0.847	0.004	0.000
calendar	Distribution regression	published_NorWest90_as_written	20	0.848	0.004	0.000
calendar	Distribution regression	published_Oregon80	20	0.873	0.003	0.000
calendar	Distribution regression	published_Oregon80_as_written	20	0.875	0.003	0.000
calendar	Distribution regression	published_USGS_311_24	20	0.517	0.000	0.000
calendar	Distribution regression	published_USGS_strict7	20	0.167	0.000	0.000
calendar	Distribution regression	published_WMO_11_5	20	0.432	0.000	0.000
calendar	Distribution regression	published_complete_year	20	0.000	–	–
calendar	Distribution regression	published_gap14	20	0.648	0.001	0.000
calendar	Distribution regression	published_gap49	20	0.806	0.005	0.000
calendar	Distribution regression	published_warm_gap14	20	0.813	0.001	0.000
calendar	Distribution regression	published_warm_gap30	20	0.885	0.003	0.000
calendar	Distribution regression	published_warm_gap7	20	0.745	0.000	0.000
calendar	Distribution regression	random_effects	20	0.000	0.003	0.000
calendar	Distribution regression	90% warm-season coverage	20	0.789	0.000	0.000
calendar	Observed days	one_feature_25	20	0.889	0.009	0.000
calendar	Observed days	Peak overlap <50%	20	0.916	0.018	0.000
calendar	Observed days	published_Beaufort_full	20	0.645	0.001	0.000
calendar	Observed days	published_Beaufort_full_as_written	20	0.641	0.002	0.000




Figure 5. Decision cost sensitivity and station eligibility in the assembled archive.

Table 5. rules; source tables contain all strata and empirical repetition intervals.

family	delivery	rule	seeds	retention	risk	groups >10%
calendar	Observed days	published_Dunham_full	20	0.654	0.004	0.000
calendar	Observed days	published_Dunham_full_as_written	20	0.655	0.004	0.000
calendar	Observed days	published_ETCCDI_15_3	20	0.350	0.000	0.000
calendar	Observed days	published_GSIM_350	20	0.590	0.003	0.000
calendar	Observed days	published_NorWest90	20	0.847	0.011	0.000
calendar	Observed days	published_NorWest90_as_written	20	0.848	0.012	0.000
calendar	Observed days	published_Oregon80	20	0.873	0.010	0.000
calendar	Observed days	published_Oregon80_as_written	20	0.875	0.010	0.000
calendar	Observed days	published_USGS_311_24	20	0.517	0.001	0.000
calendar	Observed days	published_USGS_strict7	20	0.167	0.000	0.000
calendar	Observed days	published_WMO_11_5	20	0.432	0.000	0.000
calendar	Observed days	published_complete_year	20	0.000	–	–
calendar	Observed days	published_gap14	20	0.648	0.009	0.000
calendar	Observed days	published_gap49	20	0.806	0.025	0.000
calendar	Observed days	published_warm_gap14	20	0.813	0.008	0.000
calendar	Observed days	published_warm_gap30	20	0.885	0.018	0.000
calendar	Observed days	published_warm_gap7	20	0.745	0.002	0.000
calendar	Daily plug-in	one_feature_25	20	0.889	0.005	0.000
calendar	Daily plug-in	Peak overlap <50%	20	0.916	0.008	0.000
calendar	Daily plug-in	published_Beaufort_full	20	0.645	0.001	0.000

Table 6. rules; source tables contain all strata and empirical repetition intervals.

family	delivery	rule	seeds	retention	risk	groups >10%
calendar	Daily plug-in	published_Beaufort_full_as_written	20	0.641	0.002	0.000
calendar	Daily plug-in	published_Dunham_full	20	0.654	0.003	0.000
calendar	Daily plug-in	published_Dunham_full_as_written	20	0.655	0.003	0.000
calendar	Daily plug-in	published_ETCCDI_15_3	20	0.350	0.000	0.000
calendar	Daily plug-in	published_GSIM_350	20	0.590	0.001	0.000
calendar	Daily plug-in	published_NorWest90	20	0.847	0.006	0.000
calendar	Daily plug-in	published_NorWest90_as_written	20	0.848	0.007	0.000
calendar	Daily plug-in	published_Oregon80	20	0.873	0.005	0.000
calendar	Daily plug-in	published_Oregon80_as_written	20	0.875	0.005	0.000
calendar	Daily plug-in	published_USGS_311_24	20	0.517	0.001	0.000
calendar	Daily plug-in	published_USGS_strict7	20	0.167	0.000	0.000
calendar	Daily plug-in	published_WMO_11_5	20	0.432	0.000	0.000
calendar	Daily plug-in	published_complete_year	20	0.000	–	–
calendar	Daily plug-in	published_gap14	20	0.648	0.003	0.000
calendar	Daily plug-in	published_gap49	20	0.806	0.010	0.000
calendar	Daily plug-in	published_warm_gap14	20	0.813	0.003	0.000
calendar	Daily plug-in	published_warm_gap30	20	0.885	0.007	0.000
calendar	Daily plug-in	published_warm_gap7	20	0.745	0.001	0.000
crossed	Distribution regression	all	20	1.000	0.126	0.662
crossed	Distribution regression	atc	20	0.861	0.078	0.384




Figure 6. Average risk and the fraction of evaluable groups with risk above 10%, reported separately.

Table 7. rules; source tables contain all strata and empirical repetition intervals.

family	delivery	rule	seeds	retention	risk	groups >10%
crossed	Distribution regression	Batch average	20	0.858	0.078	0.368
crossed	Distribution regression	Batch + individual cap	20	0.847	0.074	0.346
crossed	Distribution regression	gap_49	20	0.333	0.082	0.400
crossed	Distribution regression	Group certificate	20	0.000	–	–
crossed	Distribution regression	individual_0.05	20	0.708	0.038	0.068
crossed	Distribution regression	Individual score ≤0.10	20	0.768	0.052	0.149
crossed	Distribution regression	individual_0.2	20	0.832	0.070	0.303
crossed	Distribution regression	individual_0.3	20	0.871	0.081	0.405
crossed	Distribution regression	Probability + group adjustment	20	0.618	0.021	0.027
crossed	Distribution regression	mondrian	20	0.841	0.072	0.311
crossed	Distribution regression	none	20	0.000	–	–
crossed	Distribution regression	one_feature_25	20	0.381	0.015	0.014
crossed	Distribution regression	Peak overlap <50%	20	0.492	0.035	0.043
crossed	Distribution regression	published_Beaufort_full	20	0.281	0.003	0.005
crossed	Distribution regression	published_Beaufort_full_as_written	20	0.291	0.013	0.030
crossed	Distribution regression	published_Dunham_full	20	0.281	0.007	0.006
crossed	Distribution regression	published_Dunham_full_as_written	20	0.292	0.011	0.011
crossed	Distribution regression	published_ETCCDI_15_3	20	0.000	–	–
crossed	Distribution regression	published_GSIM_350	20	0.000	–	–
crossed	Distribution regression	published_NorWest90	20	0.436	0.042	0.086

Table 8. rules; source tables contain all strata and empirical repetition intervals.

family	delivery	rule	seeds	retention	risk	groups >10%
crossed	Distribution regression	published_NorWest90_as_written	20	0.442	0.043	0.114
crossed	Distribution regression	published_Oregon80	20	0.285	0.006	0.005
crossed	Distribution regression	published_Oregon80_as_written	20	0.313	0.015	0.022
crossed	Distribution regression	published_USGS_311_24	20	0.000	–	–
crossed	Distribution regression	published_USGS_strict7	20	0.000	–	–
crossed	Distribution regression	published_WMO_11_5	20	0.000	–	–
crossed	Distribution regression	published_complete_year	20	0.000	–	–
crossed	Distribution regression	published_gap14	20	0.000	–	–
crossed	Distribution regression	published_gap49	20	0.333	0.082	0.400
crossed	Distribution regression	published_warm_gap14	20	0.181	0.001	0.006
crossed	Distribution regression	published_warm_gap30	20	0.478	0.058	0.238
crossed	Distribution regression	published_warm_gap7	20	0.159	0.001	0.000
crossed	Distribution regression	random_effects	20	0.000	0.000	0.000
crossed	Distribution regression	90% warm-season coverage	20	0.184	0.001	0.006
crossed	Observed days	one_feature_25	20	0.381	0.038	0.024
crossed	Observed days	Peak overlap <50%	20	0.492	0.097	0.397
crossed	Observed days	published_Beaufort_full	20	0.281	0.005	0.000
crossed	Observed days	published_Beaufort_full_as_written	20	0.291	0.022	0.044
crossed	Observed days	published_Dunham_full	20	0.281	0.013	0.000
crossed	Observed days	published_Dunham_full_as_written	20	0.292	0.024	0.039

Table 9. rules; source tables contain all strata and empirical repetition intervals.

family	delivery	rule	seeds	retention	risk	groups >10%
crossed	Observed days	published_ETCCDI_15_3	20	0.000	–	–
crossed	Observed days	published_GSIM_350	20	0.000	–	–
crossed	Observed days	published_NorWest90	20	0.436	0.114	0.557
crossed	Observed days	published_NorWest90_as_written	20	0.442	0.121	0.595
crossed	Observed days	published_Oregon80	20	0.285	0.013	0.000
crossed	Observed days	published_Oregon80_as_written	20	0.313	0.034	0.070
crossed	Observed days	published_USGS_311_24	20	0.000	–	–
crossed	Observed days	published_USGS_strict7	20	0.000	–	–
crossed	Observed days	published_WMO_11_5	20	0.000	–	–
crossed	Observed days	published_complete_year	20	0.000	–	–
crossed	Observed days	published_gap14	20	0.000	–	–
crossed	Observed days	published_gap49	20	0.333	0.305	0.984
crossed	Observed days	published_warm_gap14	20	0.181	0.000	0.003
crossed	Observed days	published_warm_gap30	20	0.478	0.213	0.962
crossed	Observed days	published_warm_gap7	20	0.159	0.000	0.000
crossed	Daily plug-in	one_feature_25	20	0.381	0.021	0.022
crossed	Daily plug-in	Peak overlap <50%	20	0.492	0.049	0.141
crossed	Daily plug-in	published_Beaufort_full	20	0.281	0.004	0.006
crossed	Daily plug-in	published_Beaufort_full_as_written	20	0.291	0.017	0.049
crossed	Daily plug-in	published_Dunham_full	20	0.281	0.008	0.008

Table 10. rules; source tables contain all strata and empirical repetition intervals.

family	delivery	rule	seeds	retention	risk	groups >10%
crossed	Daily plug-in	published_Dunham_full_as_written	20	0.292	0.015	0.033
crossed	Daily plug-in	published_ETCCDI_15_3	20	0.000	–	–
crossed	Daily plug-in	published_GSIM_350	20	0.000	–	–
crossed	Daily plug-in	published_NorWest90	20	0.436	0.057	0.200
crossed	Daily plug-in	published_NorWest90_as_written	20	0.442	0.059	0.232
crossed	Daily plug-in	published_Oregon80	20	0.285	0.008	0.011
crossed	Daily plug-in	published_Oregon80_as_written	20	0.313	0.020	0.043
crossed	Daily plug-in	published_USGS_311_24	20	0.000	–	–
crossed	Daily plug-in	published_USGS_strict7	20	0.000	–	–
crossed	Daily plug-in	published_WMO_11_5	20	0.000	–	–
crossed	Daily plug-in	published_complete_year	20	0.000	–	–
crossed	Daily plug-in	published_gap14	20	0.000	–	–
crossed	Daily plug-in	published_gap49	20	0.333	0.127	0.641
crossed	Daily plug-in	published_warm_gap14	20	0.181	0.001	0.006
crossed	Daily plug-in	published_warm_gap30	20	0.478	0.089	0.454
crossed	Daily plug-in	published_warm_gap7	20	0.159	0.001	0.000

Table 11. recovery; source tables contain all strata and empirical repetition intervals.

family	scope	model	seeds	auc median
calendar	all	lookup	20	0.921
calendar	all	one_feature	20	0.933
calendar	all	oracle_D2	20	1.000
calendar	all	oracle_plugin	20	1.000
calendar	all	recovery_model	20	0.972
calendar	all	recovery_t5	20	0.969
calendar	all	risk_classifier	20	0.978
calendar	danger	lookup	20	0.643
calendar	danger	one_feature	20	0.615
calendar	danger	oracle_D2	20	1.000
calendar	danger	oracle_plugin	20	1.000
calendar	danger	recovery_model	20	0.780
calendar	danger	recovery_t5	20	0.767
calendar	danger	risk_classifier	20	0.751
calendar	within_cell	recovery_model	20	0.873
crossed	all	lookup	20	0.785
crossed	all	one_feature	20	0.756
crossed	all	oracle_D2	20	1.000
crossed	all	oracle_plugin	20	1.000
crossed	all	recovery_model	20	0.867

Table 12. recovery; source tables contain all strata and empirical repetition intervals.

family	scope	model	seeds	auc median
crossed	all	recovery_t5	20	0.856
crossed	all	risk_classifier	20	0.866
crossed	danger	lookup	20	0.635
crossed	danger	one_feature	20	0.557
crossed	danger	oracle_D2	20	1.000
crossed	danger	oracle_plugin	20	1.000
crossed	danger	recovery_model	20	0.784
crossed	danger	recovery_t5	20	0.771
crossed	danger	risk_classifier	20	0.763
crossed	within_cell	recovery_model	20	0.806

Table 13. Matched deep-comparator crossed cases with dangerous gaps and air-ridge information tier; three seeds only. Neighbor data are absent from neural inputs.

model	seeds	MAE	bias median	availability median
boosting	3	0.556	0.008	1.000
Global lagged-air boosting	3	0.490	0.105	1.000
Local air boosting	3	0.447	0.097	1.000
Local flow boosting	3	0.398	0.129	0.600
Global flow boosting	3	0.579	0.257	0.600
CSDI	3	0.531	-0.010	1.000
distribution_regression_calibrated	3	0.333	0.155	1.000
distribution_regression_raw	3	0.333	0.155	1.000
Functional debiasing	3	0.338	0.133	1.000
LSTM	3	0.479	0.135	1.000
SAITS	3	0.460	-0.003	1.000
stochastic_regression_calibrated_local	3	0.261	-0.065	1.000
stochastic_regression_local	3	0.261	-0.065	1.000

Table 14. Matched deep-comparator calendar cases with dangerous gaps and air-ridge information tier; three seeds only. Neighbor data are absent from neural inputs.

model	seeds	MAE	bias median	availability median
boosting	3	0.793	-0.397	1.000
Global lagged-air boosting	3	0.699	-0.045	1.000
Local air boosting	3	0.643	-0.320	1.000
Local flow boosting	3	0.345	-0.065	0.429
Global flow boosting	3	0.620	-0.281	0.429
CSDI	3	0.689	-0.304	1.000
distribution_regression_calibrated	3	0.470	0.069	1.000
distribution_regression_raw	3	0.470	0.069	1.000
Functional debiasing	3	0.524	0.134	1.000
LSTM	3	0.697	-0.083	1.000
SAITS	3	0.825	-0.518	1.000
stochastic_regression_calibrated_local	3	0.894	-0.526	1.000
stochastic_regression_local	3	0.894	-0.526	1.000

Table 15. Nonempty bounded-group certification by local label budget. Undefined selected risks are not zero-risk successes.

family	labels (groups)	runs	nonempty certificate fraction	mean retention	risk among nonempty
calendar	2	40000	0.000	0.000	–
calendar	4	28000	0.000	0.000	–
calendar	8	18000	0.000	0.000	–
calendar	10	14000	0.000	0.000	–
calendar	16	6000	0.000	0.000	–
crossed	2	40000	0.000	0.000	–
crossed	4	28000	0.000	0.000	–
crossed	8	18000	0.000	0.000	–
crossed	10	14000	0.000	0.000	–
crossed	16	6000	0.000	0.000	–

Table 16. Poststratified outcomes per 1,000 historically supported incomplete station-years. Empirical repetition medians; unsupported strata stay in the denominator.

rule	used per 1000 median	failures per 1000 median	unsupported population per 1000 median
all	997.162	28.588	2.838
individual_0.05	933.980	3.997	2.838
Individual score ≤0.10	942.808	5.769	2.838
individual_0.2	953.531	9.070	2.838
individual_0.3	961.355	11.766	2.838
one_feature_25	893.668	2.913	2.838
Peak overlap <50%	924.923	4.721	2.838
warm90	799.642	0.583	2.838

Table 17. Exact frozen-admission deletion sensitivity. The best published comparator is fixed using opened full-data outcomes at a 5 percent point-risk cap. A zero is an initial failure; unavailable reversal minima remain undefined.

family	rule	repeats	finite minima	initial failures	median min. groups
calendar	Batch average	20	20	0	179.500
calendar	Batch + individual cap	20	20	0	168.500
calendar	Individual score ≤0.10	20	20	0	150.500
calendar	Probability + group adjustment	20	20	0	87.000
crossed	Batch average	20	20	20	0.000
crossed	Batch + individual cap	20	20	20	0.000
crossed	Individual score ≤0.10	20	19	15	0.000
crossed	Probability + group adjustment	20	20	0	114.000

Table 18. Historical opened N1 uses a 7 percent point-risk criterion. Fixed-admission minima and batch-rule reapplication at the witness answer different questions; historical verdicts do not change.

population	n groups	minimum groups	remaining risk	reapplied witness risk
deploy	40	10	0.071	0.056
mech	40	10	0.071	0.062

Table 19. Conditional conformal prediction-set comparators over twenty seeds. Conditional coverage targets false-admission frequency per eligible representative, not selected risk. Region means are descriptive under held-region shift; empty releases have undefined selected risk.

family	method	nominal coverage	seeds	retention	coverage	selected risk	LP fallback (%)
calendar	Coarse groups, conservative	0.900	20	0.000	1.000	–	0.000
calendar	Coarse groups, conservative	0.950	20	0.000	1.000	–	0.000
calendar	Coarse groups, randomized	0.900	20	0.156	0.900	0.005	1.754
calendar	Coarse groups, randomized	0.950	20	0.078	0.950	0.005	0.962
calendar	Full region, randomized	0.900	20	0.000	0.901	–	0.000
calendar	Full region, randomized	0.950	20	0.000	0.950	–	0.000
calendar	Full region, conservative	0.900	20	0.000	1.000	–	0.000
calendar	Full region, conservative	0.950	20	0.000	1.000	–	0.000
crossed	Coarse groups, conservative	0.900	20	0.000	1.000	–	0.000
crossed	Coarse groups, conservative	0.950	20	0.000	1.000	–	0.000
crossed	Coarse groups, randomized	0.900	20	0.111	0.900	0.083	0.824
crossed	Coarse groups, randomized	0.950	20	0.055	0.950	0.086	0.510
crossed	Full region, randomized	0.900	20	0.000	0.900	–	0.000
crossed	Full region, randomized	0.950	20	0.000	0.950	–	0.000
crossed	Full region, conservative	0.900	20	0.000	1.000	–	0.000
crossed	Full region, conservative	0.950	20	0.000	1.000	–	0.000

Table 20. Identical classifier and cap: isolate Platt probability calibration from its additional empirical overlapping-group factors. These are exploratory source repetition medians, not the formal conditional-conformal algorithm.

family	method	retention	risk	fraction groups above 10	worst group risk
calendar	Probability + group adjustment	0.921	0.003	0.000	0.045
calendar	Platt probability calibration	0.925	0.003	0.000	0.047
calendar	Raw classifier	0.948	0.008	0.000	0.080
crossed	Probability + group adjustment	0.618	0.021	0.027	0.159
crossed	Platt probability calibration	0.638	0.024	0.032	0.170
crossed	Raw classifier	0.768	0.052	0.149	0.231

Table 21. General-loss semi-supervised and labeled-only RCPS with WSR bounds. The unit is an independent group. Distinct unlabeled calibration groups and untouched test groups are required.

method	labels (groups)	runs	nonempty	unlabeled shortage
Labeled-only WSR	2	40000	0.000	0.000
Labeled-only WSR	4	28000	0.000	0.000
Labeled-only WSR	8	18000	0.000	0.000
Labeled-only WSR	10	14000	0.000	0.000
Labeled-only WSR	16	6000	0.000	0.000
Semi-supervised WSR	2	40000	0.000	0.400
Semi-supervised WSR	4	28000	0.000	0.643
Semi-supervised WSR	8	18000	0.000	1.000
Semi-supervised WSR	10	14000	0.000	1.000
Semi-supervised WSR	16	6000	0.000	1.000

Table 22. General-loss semi-supervised and labeled-only RCPS with WSR bounds. The unit is an independent group. Distinct unlabeled calibration groups and untouched test groups are required.

method	labels (groups)	runs	nonempty	unlabeled shortage
Labeled-only WSR	2	40000	0.000	0.000
Labeled-only WSR	4	28000	0.000	0.000
Labeled-only WSR	8	18000	0.000	0.000
Labeled-only WSR	10	14000	0.000	0.000
Labeled-only WSR	16	6000	0.000	0.000
Semi-supervised WSR	2	40000	0.000	0.400
Semi-supervised WSR	4	28000	0.000	0.643
Semi-supervised WSR	8	18000	0.000	1.000
Semi-supervised WSR	10	14000	0.000	1.000
Semi-supervised WSR	16	6000	0.000	1.000

Table 23. Same-roster and same-span regional sensitivity, 1981–2025. Annual means require 80 percent fixed-roster support. Common-year and expanded-year rankings differ. Regions with fewer than three independent groups have no bootstrap interval.

region	n stations	n groups	common years	top complete	top retained	trend difference
provider:chmi	11	4.000	18.000	2015.000	2015.000	0.000
provider:foen	11	2.000	25.000	2018.000	2018.000	0.009
provider:gkd_bayern	37	8.000	10.000	2022.000	2018.000	-0.009
provider:usgs	36	19.000	1.000	2020.000	2018.000	-0.046
region:usgs_huc02	5	2.000	10.000	2002.000	2002.000	0.111
region:usgs_huc03	5	4.000	11.000	2011.000	2010.000	-0.015
region:usgs_huc04	5	2.000	12.000	2020.000	2012.000	-0.178
region:usgs_huc17	15	7.000	8.000	2015.000	2015.000	0.027

Table 24. Observed-window record certificates among 96 stations with at least fifteen complete observed years in 1981–2025. Each incomplete-year lower bound exceeds every complete-year H30 at the same station; it need not be the unique full-record maximum.

margin c	n station years	n stations	n groups
0.000	35	20	15
0.500	12	6	6
1.000	3	2	2

Table 25. Source-only joint E1/E2 planning across all twenty algorithm and group-composition repetitions. Each budget has 40,000 empirical group simulations. Seed quantiles do not represent independent-climate uncertainty; E1 uses a planning delta approximation.

independent groups	n source seeds	simulation repetitions per seed	total simulations	joint E1 E2 probability	source seed lower	source seed upper
20	20	2000	40000	0.389	0.186	0.661
30	20	2000	40000	0.337	0.118	0.640
40	20	2000	40000	0.306	0.071	0.639
60	20	2000	40000	0.253	0.040	0.631
80	20	2000	40000	0.216	0.020	0.630
100	20	2000	40000	0.192	0.010	0.623
150	20	2000	40000	0.153	0.003	0.614
200	20	2000	40000	0.134	0.000	0.609

Table 26. Exploratory redesigned P4 analogues. Original numeric criteria require RMSE reduction at least 20 percent, bias no worse by over 0.01 degrees per decade, and missing-component coverage 90–97 percent. Total interval coverage cannot substitute for missing-component coverage. Historical P4 is unchanged.

scenario	repeats	passing	missing coverage median	total coverage median	rmse reduction median
mcar	20	0	0.792	1.000	0.695
non_air_trend	20	0	0.037	1.000	-0.602
strong_heat	20	0	0.796	1.000	0.696

Table 27. Unflagged visible-air exposure: medians of three mask/trajectory seeds with clean source-fit seed zero held fixed. Missing-component and total coverage refer to 95 percent trend intervals. Air replaces seven consecutive visible warm-day water readings; the underlying target remains defined. Random and strongly heat-associated missingness have matched marginals. Censored trends are omitted because contaminated windows are no longer valid lower bounds. This is synthetic measurement contamination, not dry-bed water truth or twenty-fit stability.

condition	scenario	pipeline	rmse	bias	missing coverage	total coverage
Clean	Random	90% warm-season rule	0.039	0.001	–	1.000
Clean	Random	Distribution-regression MI	0.011	-0.001	0.796	1.000
Clean	Random	Calibrated stochastic MI	0.012	-0.001	0.721	1.000
Clean	Heat	90% warm-season rule	0.045	-0.001	–	1.000
Clean	Heat	Distribution-regression MI	0.014	-0.000	0.799	1.000
Clean	Heat	Calibrated stochastic MI	0.014	-0.001	0.719	1.000
Air	Random	90% warm-season rule	0.085	-0.004	–	0.997
Air	Random	Distribution-regression MI	0.075	-0.004	0.169	0.998
Air	Random	Calibrated stochastic MI	0.075	-0.005	0.135	0.998
Air	Heat	90% warm-season rule	0.099	0.014	–	0.993
Air	Heat	Distribution-regression MI	0.091	0.017	0.161	0.995
Air	Heat	Calibrated stochastic MI	0.090	0.016	0.130	0.995

Table 28. The same clean source failure classifier and individual cap 0.1 under unflagged air exposure. Three mask/trajectory seeds share one source fit. Group risks require at least twenty admissions; no contaminated labels recalibrate the score.

condition	family	retention	risk	groups >10%	worst group risk
Clean	calendar	0.946	0.007	0.000	0.065
Clean	crossed	0.766	0.049	0.168	0.224
Air	calendar	0.950	0.369	0.818	1.000
Air	crossed	0.792	0.379	0.919	0.952

Table 29. Matched forcing sensitivities at all 319 benchmarkable USGS source sites. MAE is in degrees C. Exact original panels, forcing and daily Ridge code reproduce all 67,316 POWER control identities and available points. Each alternative has its own shared-case reference; different rows must not be compared as if they used one sample. Missing inputs stay in eligible denominators. Daymet uses a minimum/maximum midpoint; daily time conventions also vary. Nearby TAVG is not a perfect co-located or clock-matched oracle.

Input	Stratum	Eligible	Paired	Stations	POWER MAE	Alternative MAE
Daymet	All	67316	66133	319	0.317	0.325
Daymet	Danger	27530	27418	319	0.640	0.657
Daymet	Air, danger	9347	9271	308	0.661	0.682
POWER + flow	All	67316	51060	261	0.334	0.302
POWER + flow	Danger	27530	20972	259	0.683	0.612
POWER + flow	Air, danger	9347	7048	244	0.708	0.623
NOAA TAVG	All	67316	22028	122	0.318	0.312
NOAA TAVG	Danger	27530	8984	118	0.655	0.642
NOAA TAVG	Air, danger	9347	3010	112	0.678	0.671

Table 30. Native H7 source refits, 20 seeds and 540 outer-region fits. Seven-day peak supports, masks, history summaries, conditional functionals, MOVE augmentation, annual models and disjoint calibration are recomputed. MAE and proper scores are in degrees C; coverage refers to 90 percent intervals. These are dangerous decision-time cases, not the historical H30 endpoint; deep models are not refitted here.

family	estimator	seeds	MAE	CRPS	twCRPS	coverage
calendar	Distribution regression	20	0.494	0.365	0.065	0.874
calendar	Functional debiasing	20	0.522	–	–	–
calendar	Daily plug-in	20	0.638	–	–	–
calendar	Calibrated stochastic	20	0.516	0.380	0.068	0.858
crossed	Distribution regression	20	0.348	0.257	0.046	0.892
crossed	Functional debiasing	20	0.375	–	–	–
crossed	Daily plug-in	20	0.423	–	–	–
crossed	Calibrated stochastic	20	0.360	0.265	0.048	0.893

Table 31. Task calibration and independent-draw intervals on exactly the same supported cases. Coverage is a fraction; missing source support is reported separately.

Setting	Method	n	Original coverage	Task coverage	Width ratio
calendar / forward	D2	340	0.979	0.841	0.536
calendar / forward	Stochastic	332	0.991	0.873	0.524
calendar / offline	D2	840	0.935	0.969	1.274
calendar / offline	Stochastic	835	0.917	0.903	1.150
crossed / forward	D2	340	0.968	0.965	1.158
crossed / forward	Stochastic	331	0.961	0.976	1.297
crossed / offline	D2	840	0.918	0.971	1.478
crossed / offline	Stochastic	838	0.862	0.952	1.442

Table 32. Task calibration support. Empirical calibration from small source pools does not establish a finite-sample guarantee.

Family	Time setting	Estimator	Availability	Minimum source groups
calendar	forward	D2	0.415	1
calendar	forward	stochastic_calibrated	0.405	1
calendar	offline	D2	1.000	2
calendar	offline	stochastic_calibrated	0.994	2
crossed	forward	D2	0.415	1
crossed	forward	stochastic_calibrated	0.404	1
crossed	offline	D2	1.000	2
crossed	offline	stochastic_calibrated	0.998	2

Table 33. Additional observation-only witnesses from approved USGS daily means. Every candidate remains in the qualification table. Sparse complete-reference support precludes an incidence or transfer inference.

Station	Year	Complete n	Observed L	Complete max	Excess
08377900	2016	3	14.990	14.853	0.137
08377900	2018	3	15.537	14.853	0.683
08377900	2020	3	15.737	14.853	0.883
08405200	2013	2	28.693	28.327	0.367
08405200	2015	2	28.703	28.327	0.377
08405200	2016	2	28.537	28.327	0.210
08405200	2019	2	28.833	28.327	0.507
08405200	2020	2	28.970	28.327	0.643
07364133	2015	1	29.427	28.767	0.660




Figure 7. Trend RMSE, missing-component interval coverage and hot-year event skill. The shaded interval is the 90–97% coverage gate for nominal 95% missing-component intervals. Reduced RMSE does not establish calibrated functional uncertainty. The incompatible annual diagnostic differs from the separate full daily refit.