Supporting Information for "Retaining incomplete river-temperature records: temporal support and uncertainty in sustained-heat trends"

Author information to be completed by the authors

Contents of this file

1. Text S1–S40.

2. Tables S1–S68, including external-reference Table S46.

The following sections document source-comparison denominators, whole-archive and interval procedures, reference qualification, and sensitivity tables for the main article. Source data, model outputs and software require the separate access arrangements described in the article's Open Research section; this file does not substitute for those materials.

S1. Design and information separation

The source comparison uses twenty repetitions of masks and group-role assignment on the same assembled archive; these are not independent climates. Held test connected groups are excluded from estimator fitting, interval calibration, risk learning and rule calibration. In the original partitioned forward models, other training-only groups in a named target region can remain in source roles. The source-role audit and subsequent strict-region sensitivities distinguish this exception from the held target-group exclusion.

In each whole-archive setting, selected gaps in the target station are masked together before local daily fitting. Unmasked target observations and the designated donor observations remain available. The source annual model is fitted separately and held fixed within that archive reanalysis. The first seven years are observed burn-in; the forward evaluation masks only years from 2010 onward. Twenty repetitions at the same station are not twenty independent rivers. Initial seven-station stress pilots are followed by a 41-station forward sensitivity; the heat-associated extension fixes the number of missing years but can change their dates and the number of missing days.

Correction to historical design metadata

The 3,320 saved whole-archive `design.json` files are historical records and were not rewritten. Their field `annual_model` states that "held region and connected groups excluded." That statement is correct for the held test connected groups, but it is not an unconditional region exclusion. The original forward HUC17 estimator retained one other same-region training group, and NVE interval calibration retained two other same-region groups. The paper therefore distinguishes the held-test-group main analysis from the strict-region sensitivity. This correction applies to the historical metadata; it does not change any saved prediction. A correction manifest is available in `results/wrr_adversarial_review_20261027/metadata_correction/`.

S2. Observation-rule accounting

The rule table uses every eligible station-year, not only the selected witnesses. A reported window is not a complete-year H30, and an accepted incomplete year can contain an unobserved higher peak. The "new facts" column therefore compares reported windows

with the maximum reported window among the other accepted years; it does not prove a full-record ordering. The measurement-margin sensitivity treats a window ordering as unresolved when the reported difference is at most twice the assumed absolute error on each window. These margins are assumptions, not sensor-error estimates.

S3. Joint-rank numerical convergence

The convergence check fixes the annual quantiles, observed bounds and evaluation years and increases the stratified sample count through 256, 1,024 and 4,096 on the difficult forward controlled gaps. Three random seeds index the rank construction. The rank sequence is generated at 4,096 draws and truncated for smaller counts; annual marginals are regenerated separately for each sample size and remain specific to each estimator. Coverage, width and interval score are reported with the across-seed range. The range is not a confidence interval, and the check uses the same opened data as the main analysis. A 0.3-point coverage rule and 1% width and interval-score rules are applied to the three-seed mean. Five of six method–construction pairs meet the coverage rule on that mean. The largest three-seed-mean coverage change is 0.36 percentage points, and the largest single-seed change is 0.815 percentage points. Averaging seeds can cancel opposing changes, so these rules are not a resolution guarantee and the constructions are not ranked. At 4,096 samples, nominal 95% trend coverage remains 87.4–89.9%.

S4. Scoring and interval construction

For ensemble $x_1, \ldots, x_M$ and outcome y, the continuous ranked probability score is

$$M^{-1} \sum_{m} |x_m - y| - (2M^2)^{-1} \sum_{m,l} |x_m - x_l|.$$

Threshold-weighted scoring replaces samples and outcome by their maxima with the training H30 threshold. The 90% interval score adds twenty times the distance outside the interval to its width. A 95% trend interval score uses the corresponding factor forty. Missing predictions remain unavailable rather than being assigned zero error.

Both annual estimators use the observed-window support projection where a finite bound exists. The projection is applied to the annual distribution and then carried into the slope; a trend interval is not projected onto an annual lower bound. Seven stored quantiles do not determine their outer tails; the normal-spacing continuation and the alternative constant tail are assumptions. Within one estimator, cross-year rank constructions keep that estimator's annual marginal samples fixed. The two estimators have different annual marginals. The constructions are not a posterior for the fitted annual model. The source-only corrections and the post-hoc propagation reported below do not establish a finite-sample or independent-climate coverage guarantee.

In whole-archive annual coverage and the peak-visibility analysis, inclusive endpoint comparisons use an absolute $10^{-8}\,^{\circ}\mathrm{C}$ numerical tolerance. After detecting boundary sensitivity in the USGS exploration, we compared this convention with exact floating-point comparisons and a $10^{-10}\,^{\circ}\mathrm{C}$ tolerance on the same saved two-estimator common cases. For the forward empirical-calendar subset, D2 covers 3,610 of 3,674 repeated masked cases with either tolerance, but an exact comparison covers 2,037; all 1,573 changed comparisons lie in the reference-at-observed-bound stratum. Across all four family–mode combinations and both estimators, the two nonzero tolerances agree on every common-case classification. An independent scalar audit found the largest strict-comparison endpoint gap among changed method–case comparisons was $8.18 \times 10^{-14}\,^{\circ}\mathrm{C}$. These tolerances concern arithmetic on the

reported target, not physical measurement uncertainty, and masks from a station-year are not independent. The pooled common-case comparison differs in weighting and availability from the equal-archive means used elsewhere; no original saved coverage was replaced.

The historical single-year source comparison also uses an absolute $10^{-8}\,^{\circ}$C endpoint tolerance after support projection. A post-hoc check reproduced the published coverage for all 20 saved source repetitions, two methods, two gap families and both overlap strata. In the controlled-gap, high-overlap stratum, 10^{-10} and $10^{-8}\,^{\circ}$C give identical coverage classifications and median D2 and stochastic coverage remains 89.3% and 89.8%, respectively. Outside the high-overlap stratum there are 14 repeated method–case comparisons where the two tolerances differ, with endpoint gaps between 1.89×10^{-9} and $9.93 \times 10^{-9}\,^{\circ}$C. An independent scalar implementation reproduced the 160 method–family–stratum–seed counts. These are the same historical source predictions, not independent sensor validation or a population calibration guarantee.

For the exploratory release-then-trend analysis, an observed lower bound L contributes $\log \Phi\{(\mu - L)/\sigma\}$ to a Gaussian working likelihood. Released numerical estimates are treated as exact, unknown cases remain unavailable, and the working interval includes regression variability. It is not a calibrated missingness-only interval.

S5. Core source comparisons

S5.1. Additional observation-only corroboration

We obtained approved daily-mean USGS temperature records at ten additional metadata-selected stream stations. None was a source-model target. Seven candidates lie in HUC2 regions 13 or 20, absent from the source station regions; three are additional station checks within already represented HUC2 regions. The candidates and all qualification failures are

Table S1. Dangerous source-case summaries: the gap overlaps at least half the training-period peak window. Entries are medians across 20 repetitions sharing the source archive. Method-specific availability can differ; paired case comparisons are reported separately. Coverage and CRPS are not defined for point-only methods.

Family	Method	Availability	MAE	90% coverage	CRPS
calendar	D2	100.0%	0.451	86.7%	0.329
calendar	functional	100.0%	0.476	–	–
calendar	plug-in	100.0%	0.564	–	–
calendar	classical	99.9%	0.463	86.0%	0.336
crossed	D2	100.0%	0.337	89.3%	0.246
crossed	functional	100.0%	0.355	–	–
crossed	plug-in	100.0%	0.428	–	–
crossed	classical	99.7%	0.336	89.8%	0.245

Table S2. Held-region source-calibrated comparison of the three-output and score-only rules. The score cutoff is chosen from disjoint source rule-calibration groups. Retention differs in the evaluated target cases; errors are conditional on numerical publication.

Family	Method	Eligible	Published	Retention	Error rate
calendar	score only	31,672	26,896	84.9%	0.0%
calendar	three outputs	31,672	27,090	85.5%	0.7%
crossed	score only	114,012	80,980	71.0%	3.9%
crossed	three outputs	114,012	82,169	72.1%	4.9%

Table S3. Retrospective equal numerical-count comparison. The score threshold uses the evaluated feature distribution and the three-output count, so this is transductive. Equal numerical counts do not equate retained bounds or unknown states.

Condition	Family	Eligible	Published by each	Three failures	Score failures	Three / score error
air substitution	calendar	31,672	88	19	23	21.6% / 26.1%
clean	calendar	31,672	27,090	190	4	0.7% / 0.0%
clean	crossed	114,012	82,169	4026	3249	4.9% / 4.0%
future	crossed	27,846	17,583	1924	1220	10.9% / 6.9%

Table S4. Annual and trend coverage on the common archive cohort. Annual coverage weights cases by incomplete years. Tail and rank experiments use different random draws. Within one estimator the three rank constructions share that estimator's annual marginals; the two estimators do not share one annual distribution. Repetitions share the same source archive.

Family	Mode	Method	Cases	Groups	Annual 90%	Constant 95%	Normal 95%	Independent 95%	Shared 95%	Blocks 95%
calendar	forward	D2	818	17	98.2%	94.6%	95.7%	96.0%	96.1%	96.3%
calendar	forward	classical	699	17	98.2%	94.6%	95.3%	95.4%	95.4%	95.7%
calendar	offline	D2	840	18	98.5%	92.3%	94.9%	95.0%	95.1%	94.9%
calendar	offline	classical	835	18	97.4%	91.7%	93.3%	93.2%	93.3%	93.3%
crossed	forward	D2	818	17	90.1%	83.0%	88.8%	88.0%	87.9%	89.0%
crossed	forward	classical	736	17	88.1%	81.4%	86.3%	86.3%	86.1%	87.2%
crossed	offline	D2	840	18	94.1%	91.7%	94.2%	94.0%	94.5%	94.8%
crossed	offline	classical	838	18	92.6%	86.4%	90.7%	90.2%	91.6%	91.2%

Table S5. Evidence map. Each claim is paired with the comparison and metric used here and with the boundary that limits it. The table indexes the results; it is not an additional test.

Claim	Primary comparison	Metric	Boundary
Observed windows can overturn a complete-only record	Complete-year H30 vs. fully observed 30-day window L	Count of L above the complete-year maximum	Reported measurements only; not the unique warmest year
Target recovery beats a daily plug-in	D2 vs. plug-in, functional debiasing and calibrated stochastic regression	H30 MAE, CRPS and 90% coverage	No unique gain over calibrated stochastic regression
Retaining years improves finite-record trends	Warm-season rule vs. all supported years at the same estimator	Trend RMSE; arithmetic four-cell retention/estimator contrasts	Contrasts are setting-specific, not causal year-count shares
Annual and trend uncertainties need distinct evaluation	Bounded normal-tail vs. constant-tail marginals; three rank couplings	Coverage, width, interval score and availability	Tail, annual and dependence errors are not uniquely separated
Numerical release has a risk–retention trade-off	Fixed score cap vs. three-output rule	Selected risk, retention and group excess	A raw individual score is not a group-risk guarantee
Visible diagnostics do not restore extrapolation safety	Clean vs. hot-year, future-year and air-exposure stress	Selected risk, coverage and retention	Synthetic stress severity; not an archive prevalence

Table S6. Every eligible station-year under the compared completeness rules. "Number" is a complete-year H30; "bound" is an incomplete year with a fully observed 30-day window; "unknown" has neither. "New facts" counts admitted incomplete years whose observed window exceeds the maximum reported window among other accepted years.

Rule	Eligible	Admitted	Number	Bound	Unknown	New facts	Excluded facts
Complete year (100%)	2,700	2,139	2,139	0	0	0	35
Annual coverage ≥95%	2,700	2,591	2,139	452	0	18	3
Station warm-season ≥90%	2,700	2,634	2,139	495	0	19	2
Fixed warm-season ≥90%	2,700	2,635	2,139	496	0	19	2
July–August complete	2,700	2,509	2,139	370	0	18	3
Longest gap ≤49 d (context)	2,700	2,653	2,139	514	0	19	1

A reported window is not a complete-year H30. A rule-accepted incomplete year can contain a higher unobserved peak, so the last two columns compare reported windows only. The 49-day row is a contextual Johnson comparison, not a validated H30 rule.

Table S7. Measurement-margin sensitivity for the station-specific warm-season rule. A reported window ordering is counted as resolved only when the observed difference exceeds twice the assumed absolute error on each window. The margins are assumptions, not sensor-error estimates.

Assumed absolute error (°C)	Status	Facts	Resolved above 2ϵ	Not resolved
0.00	admitted	19	19	0
0.00	excluded	2	2	0
0.01	admitted	19	18	1
0.01	excluded	2	2	0
0.05	admitted	19	15	4
0.05	excluded	2	2	0
0.10	admitted	19	8	11
0.10	excluded	2	0	2
0.20	admitted	19	4	15
0.20	excluded	2	0	2
0.50	admitted	19	1	18
0.50	excluded	2	0	2

retained. This unregistered intake was not a random sample or a blinded confirmation, and the reference histories are shorter and sparser than the original long-record panel.

The unchanged four-complete-year reconstruction criterion yields zero eligible numerical-recovery references. The observation-only claim needs a different comparison: an incomplete year can contradict the maximum of the complete-year record whenever at least one approved-complete reference year exists. We retain every approved finite reference value, including reference years that an extra heuristic-QC filter might remove. Candidate windows must contain thirty consecutive approved, nonestimated daily values after source

QC, and the incomplete year must have an actual absent, null, contradictory or unapproved provider day. Heuristic censoring alone cannot create a witness.

There are four stations with a nonempty approved-complete reference, each with only one to three reference years. At 3 of these stations, 9 incomplete station-years contain verified windows exceeding every approved-complete H30 in the analyzed record (Figure 2 of the main article). For example, the Rio Mora 2020 window exceeds the complete-only record by approximately 0.883°C, and a Pecos River 2020 window by 0.643°C. These facts corroborate record omission without fitting any imputer. They do not estimate an omission incidence, verify a unique warmest year, establish external reconstruction accuracy, or validate trend or risk guarantees. Every witness includes the thirty original approved values; agency reporting and measurement validity remain assumptions.

Table S8. Numerical convergence of the three fixed-marginal joint-rank constructions on the difficult forward controlled gaps. Annual quantiles, bounds and evaluation years are fixed. The range is across three random seeds; it is not a confidence interval.

Method	Construction	Draws	Coverage (%)	Width (°C/dec)	Interval score
D2	independent	256	88.3 [88.0, 88.5]	0.450 [0.449, 0.451]	0.855 [0.849, 0.860]
D2	independent	1024	89.2 [89.0, 89.4]	0.457 [0.457, 0.458]	0.845 [0.842, 0.847]
D2	independent	4096	89.2 [89.1, 89.2]	0.459 [0.459, 0.459]	0.842 [0.839, 0.844]
D2	residual blocks	256	89.2 [88.6, 89.6]	0.451 [0.450, 0.451]	0.840 [0.828, 0.851]
D2	residual blocks	1024	89.8 [89.7, 89.9]	0.456 [0.456, 0.456]	0.828 [0.825, 0.830]
D2	residual blocks	4096	89.7 [89.6, 89.9]	0.457 [0.457, 0.458]	0.825 [0.825, 0.826]
D2	shared parameters	256	88.4 [88.3, 88.6]	0.444 [0.444, 0.445]	0.855 [0.845, 0.863]
D2	shared parameters	1024	88.7 [88.3, 88.9]	0.450 [0.449, 0.451]	0.840 [0.835, 0.846]
D2	shared parameters	4096	88.7 [88.5, 88.9]	0.452 [0.451, 0.452]	0.835 [0.834, 0.835]
Classical	independent	256	86.3 [86.0, 86.7]	0.384 [0.383, 0.384]	0.856 [0.849, 0.866]
Classical	independent	1024	87.5 [87.2, 87.6]	0.390 [0.390, 0.390]	0.853 [0.852, 0.853]
Classical	independent	4096	87.6 [87.5, 87.6]	0.391 [0.391, 0.391]	0.851 [0.851, 0.852]
Classical	residual blocks	256	86.9 [86.5, 87.1]	0.384 [0.383, 0.385]	0.840 [0.837, 0.845]
Classical	residual blocks	1024	87.6 [87.4, 87.9]	0.389 [0.389, 0.390]	0.837 [0.835, 0.839]
Classical	residual blocks	4096	88.0 [87.8, 88.2]	0.390 [0.390, 0.391]	0.837 [0.836, 0.839]
Classical	shared parameters	256	86.3 [85.9, 86.5]	0.380 [0.380, 0.381]	0.867 [0.864, 0.868]
Classical	shared parameters	1024	87.3 [87.0, 87.5]	0.385 [0.385, 0.385]	0.856 [0.854, 0.859]
Classical	shared parameters	4096	87.5 [87.4, 87.6]	0.386 [0.386, 0.386]	0.854 [0.852, 0.856]

S6. Whole-archive supplements

Full paired MSE cells, group and region sensitivities, unavailable cases and source calibration support are in the review-response results. Forward task calibration ends before 2010. Normal-spacing tail extrapolation is compared to constant tails under the same observed-window constraint; its outer distribution is assumed, not identified by the seven stored quantiles. Auxiliary rank coupling is not a posterior for the annual estimator.

S7. Matched numerical release within each archive

After the original release and trend results were inspected, we fixed a further descriptive comparison on the same 3,320 archive settings. In each archive the three-output rule's number of numerical releases was matched exactly by choosing the lowest available raw classifier scores; equal scores were broken by year, and the existing detected-quality exclusion was applied to both rules. The choice uses the target archive's score distribution and its original release count, but no reference outcomes. It is therefore transductive and not a prospective release threshold. The same observed years and D2 estimates are used, while the selected years can differ. Nonselected years retain a visible lower bound if allowed by the same quality flag, and otherwise remain unknown. An unassessed quality diagnostic does not count as a detected flag in this analysis, as in the original pipeline.

Table S15 gives annual errors and working-likelihood trend RMSE; lower-bound counts are not an assertion that the same years or bound values were retained. The number of numerical outputs is matched within each archive, but equal counts do not establish equal total information or comparable prospective risks. Repeated masks are not independent climates. A connected-group bootstrap of paired squared-error differences for forward

Table S9. The same 35 observation-only complete-year witnesses under each screening rule. The accepted-record comparison uses observed windows, not unknown complete H30 values. The 49-day and July–August rows are contextual proxies, not published H30 rules.

Rule	Witnesses	Admitted	Excluded	Above accepted
Complete	35	0	35	35
Annual 95	35	31	4	3
Station warm 90	35	32	3	2
Fixed warm 90	35	32	3	2
July–August	35	30	5	3
Gap $\leq$49 d	35	34	1	1

Table S10. Seven-station stress comparisons on common available station/repetition rows. Heat-associated masks and numbers of missing years differ between arms; unavailable cases remain in the eligible denominator. Coverage and width are supplied in the result table.

Scenario	Eligible	Available	Base RMSE	Stress RMSE
Donors / forward	30	30	0.079	0.116
Donors / offline	35	35	0.037	0.047
Heat / forward	30	30	0.079	0.088
Heat / offline	35	30	0.035	0.058

Table S11. Release followed by finite-record trend inference for controlled gaps. The values-and-bounds interval is a Gaussian working-likelihood interval, including regression variability; it is not calibrated missing-data uncertainty.

Mode	Released data	Eligible	Available	RMSE	Working width
forward	Values + bounds	820	820	0.202	3.262
forward	Values	820	820	0.210	–
offline	Values + bounds	840	840	0.040	1.986
offline	Values	840	840	0.043	–

Table S12. Task calibration and independent-draw intervals on exactly the same supported cases. Coverage is a fraction; missing source support is reported separately.

Setting	Method	n	Original coverage	Task coverage	Width ratio
calendar / forward	D2	40	0.925	0.975	0.947
calendar / forward	Stochastic	38	1.000	1.000	0.773
calendar / offline	D2	95	0.937	0.958	1.167
calendar / offline	Stochastic	88	0.977	0.989	1.225
crossed / forward	D2	39	1.000	1.000	1.112
crossed / forward	Stochastic	38	1.000	1.000	1.015
crossed / offline	D2	301	0.977	0.980	1.236
crossed / offline	Stochastic	307	0.948	0.951	1.108

Table S13. Task calibration support. Empirical calibration from small source pools does not establish a finite-sample guarantee.

Family	Time setting	Estimator	Availability	Minimum source groups
calendar	forward	D2	0.049	3
calendar	forward	stochastic_calibrated	0.046	3
calendar	offline	D2	0.113	3
calendar	offline	stochastic_calibrated	0.105	3
crossed	forward	D2	0.048	3
crossed	forward	stochastic_calibrated	0.046	3
crossed	offline	D2	0.358	3
crossed	offline	stochastic_calibrated	0.365	3

Table S14. Additional observation-only witnesses from approved USGS daily means. Every candidate remains in the qualification table. Sparse complete-reference support precludes an incidence or transfer inference.

Station	Year	Complete n	Observed L	Complete max	Excess
08377900	2016	3	14.990	14.853	0.137
08377900	2018	3	15.537	14.853	0.683
08377900	2020	3	15.737	14.853	0.883
08405200	2013	2	28.693	28.327	0.367
08405200	2015	2	28.703	28.327	0.377
08405200	2016	2	28.537	28.327	0.210
08405200	2019	2	28.833	28.327	0.507
08405200	2020	2	28.970	28.327	0.643
07364133	2015	1	29.427	28.767	0.660

controlled gaps gives [-0.0017, 0.0072] (°C/decade)^2; this is descriptive and spans zero. The likelihood treats released predictions as exact, censoring may be informative, and its intervals are not calibrated missingness-only uncertainty. No model was refitted for this comparison.

Table S15. Per-archive matched numerical output. Annual errors exceed 0.5 °C. Counts and RMSE are original three-output rule / score-only comparator; RMSE uses common converged release-plus-bound working-likelihood fits and units °C per decade.

Scenario	Numeric	Errors	Bounds	Fits	RMSE
Calendar, offline	6,345	11 / 9	590 / 590	840	0.0198 / 0.0237
Calendar, forward	3,435	43 / 32	371 / 370	819	0.1240 / 0.1220
Controlled, offline	5,543	242 / 229	1378 / 1378	840	0.0405 / 0.0407
Controlled, forward	3,174	397 / 370	603 / 603	820	0.2024 / 0.1959

The audit separately reconstructed the selection masks for all 3,320 archives using Python ordering, recomputed 3,319 available score-comparator numerical-only OLS slopes and checked all eight summary cells. A second implementation refitted all 6,640 rule-by-archive censored likelihoods using normal log-density and survival functions. Convergence status agreed for every fit; 6,638 converged in both implementations, and their largest slope difference was approximately 1.2×10^{-6} °C per decade. This numerical check used the same observations, selected cases and working likelihood; it did not independently reconstruct source sensors or models, establish likelihood validity or verify interval standard errors.

S8. Expanded missingness-mechanism sensitivities

After the earlier results and seven-site pilot had been inspected, we fixed an expansion to the first five masks for all 41 forward-eligible original archive stations in 17 connected groups. Only controlled gaps and one seed-zero source-model assignment are studied. For the donor sensitivity, the target masks are unchanged and those same dates are hidden in every other panel column before local fitting. All 205 settings have at least one finite donor-panel observation removed; the total of 427,375 finite donor-panel cells counts repeated masks and may include columns not ultimately selected by the local estimator. It is not a count of independent outages or necessarily of observations the estimator would have used.

For the heat-associated sensitivity, the original 35% Bernoulli draw determines the number of masked years in each station–repetition archive. An independent seeded exponential-race sample chooses exactly that number of eligible years, with weight five for years whose annual air H30 is at least the first-seven-year 80th percentile, and weight one otherwise. Air information determines a synthetic test mask; it is not a hidden water-temperature outcome. The full cohort has 943 masked years in either design, of which 511 and 700 are classified as air-hot. However, the selected years have different existing within-year gap shapes and positions, so the total hidden target days change from 55,950 to 57,540. These paired scenarios cannot isolate the causal effect of heat association alone or represent a measured outage mechanism.

Table S16 compares fixed-prediction cohorts for both estimators, with availability reported on the common subset. Nominal trend interval coverage, widths and scores use independent annual ranks and the original normal-spacing tails; the intervals are not

validated climate-transfer guarantees. Connected-group resampling of paired D2 squared-error differences gives [0.0030, 0.0434] for synchronized donors and [-0.0040, 0.0091] for heat-associated masking, in $(°C/decade)^2$. These describe a selected archive and depend on group exchangeability. Removing Bavaria without refitting changes baseline and donor RMSE to 0.0829 and 0.1287, and baseline and heat RMSE to 0.0829 and 0.0816. Other fixed-prediction leave-region-out summaries are supplied in the aggregate files. Repeated masks are not independent climates; original forward source roles are not uniformly strict named-region holdouts.

Table S16. Expanded paired controlled-gap stresses, original / stress. RMSE and interval widths are °C per decade. Coverage is for nominal 95% trend intervals; scores are also in °C per decade. Common fitted cohorts differ between estimators and scenarios.

Scenario	Estimator	Cases	RMSE	Coverage	Width	Score
Donor gaps	D2	205	0.1555 / 0.2024	92.2 / 86.8	0.465 / 0.479	0.756 / 1.039
Donor gaps	Stochastic	190	0.1386 / 0.1877	87.4 / 76.3	0.404 / 0.418	0.808 / 1.257
Heat, fixed years	D2	205	0.1555 / 0.1655	92.2 / 88.8	0.465 / 0.456	0.756 / 0.806
Heat, fixed years	Stochastic	187	0.1370 / 0.1552	87.7 / 87.2	0.403 / 0.383	0.776 / 0.884

An independent audit reconstructs all 205 baseline masks, 205 synchronized masks and 205 heat-associated masks, and recomputes eight paired RMSE and mean-squared-error cells. It uses the same source observations, random seeds and saved fitted predictions; it does not independently refit the models, check bootstrap intervals or establish sensor quality.

S9. Heat-associated years with matched gap templates

After examining earlier results and the available gap templates, we repeated the forward controlled-gap sensitivity for 41 stations and five archive masks per station. The selected heat-associated years are the same as in the prior heat sensitivity. For years masked in both the baseline and heat design, the original gap is retained; remaining baseline offset–length templates are transferred to the newly selected heat years. Every archive retains its original masked-year count, multiset of templates and actual number of hidden days. Total hidden days are 55,950 in either design, and air-hot masked years increase from 511 to 700; repeated masks are not independent observations. Offsets are relative to the year, and the relationship of a transferred gap to a particular seasonal peak can change.

The matched-gap plan incorrectly described the first-seven-year air baseline as entirely pre-2010. In 4 of the 41 stations, one of those seven reference-year labels is 2010 or later. The saved implementation uses the first seven reference years; no predictions were changed by this clarification. Air information constructs a synthetic retrospective mask and is not a hidden water-temperature outcome, but the mechanism is not a strictly prior-vintage or prospective outage generator.

After viewing the pairwise results, we additionally restricted all three designs and both methods to the same 183 available station–replicate settings from 41 stations and 17 connected groups (Table S17). Group-resampled differences in paired MSE for matched minus baseline are [-0.0096, 0.0039] for D2 and [-0.0074, 0.0081] for the stochastic comparator, in $(°C/\text{decade})^2$. The shared cohort is post-hoc, and these differences change in group-equal weighting. The fixed heat-year count and gap-template distribution do not isolate year,

position and peak alignment, establish a natural-outage mechanism, or prove climate transfer.

Table S17. Three-way common cohort, baseline / previous heat / matched heat, with independent annual ranks. RMSE, widths and interval scores are °C per decade; coverage is for nominal 95% trend intervals.

Method	RMSE	Coverage (%)	Width	Interval score
D2	0.153 / 0.162 / 0.142	92.3 / 90.2 / 91.3	0.464 / 0.445 / 0.445	0.743 / 0.770 / 0.696
Stochastic	0.138 / 0.155 / 0.136	88.0 / 87.4 / 85.8	0.408 / 0.385 / 0.391	0.789 / 0.878 / 0.865

Separate checks reconstruct all 205 masks from the same underlying panels and seeded selection and independently recompute 32 paired summary rows, 96 fixed-prediction leave-region-out rows, 24 three-way summary rows and 16 three-way contrasts, including seeded group resampling. The underlying observations, trained source models and local daily refits are reused; these checks do not constitute independent physical or climate validation.

S10. Source-model sensitivity of matched heat-gap archives

After observing the earlier source-role and matched-gap results, we held each target mask and daily feature record fixed and re-applied three source-model choices. The original version uses the original source model in every region. The strict-original version replaces only the affected HUC17 model with its earlier refit excluding all HUC17 source groups; elsewhere it is identical to the original. The candidate-provider version uses the earlier target-region-excluded models, which also change the provider set, sample size and role allocation. In all, 205 candidate and 15 strict-original matched-gap settings were

re-scored. The source models and daily feature fits were not independently retrained for this comparison.

Table S18 uses the same 184 archives at 41 stations and 17 connected groups, where all six source-version by baseline/stress results are available for both methods. The nominal 95% trend intervals use independent annual ranks. Group resamples concern repeated masks and depend on the sampled groups; all displayed paired-MSE intervals cross zero. Changing source versions also changes group-equal estimates, so the apparent average direction is not a general effect of heat-associated outage. The four stations whose seven-year air reference includes an evaluation-period year, template transfer and retrospective scenario selection remain limitations of the same stress design.

Table S18. Matched-gap source sensitivity, baseline / stress on the same two-method, six-way common cohort. RMSE is °C per decade; group-equal paired MSE difference and group-resampled intervals are $(°C/\text{decade})^2$.

Method	Source	RMSE	Coverage (%)	Group-equal MSE	Paired MSE interval
D2	Original	0.152 / 0.142	92.4 / 91.3	0.0013	[-0.0092, 0.0039]
D2	Strict original	0.153 / 0.142	92.4 / 90.8	0.0008	[-0.0092, 0.0036]
D2	Candidate providers	0.153 / 0.134	88.0 / 90.8	-0.0013	[-0.0130, 0.0026]
Stochastic	Original	0.138 / 0.136	88.0 / 85.9	0.0057	[-0.0071, 0.0081]
Stochastic	Strict original	0.138 / 0.136	88.0 / 85.9	0.0057	[-0.0070, 0.0087]
Stochastic	Candidate providers	0.145 / 0.140	87.0 / 85.9	0.0042	[-0.0084, 0.0067]

After inspecting the source-specific results, we also computed a directly paired difference between the stress-minus-baseline MSE changes under the candidate and original sources. For D2 with independent ranks, this difference is -0.0024 with group-resampled interval

[-0.0052, -0.0001] (°C/decade)^2. The interval does not cross zero but its upper limit is close to zero, whereas each within-source interval in Table S18 crosses zero; paired contrasts account for covariance and are not equivalent to comparing two separate intervals. The preliminary calculation used separate 4,000-draw streams across rank representations that shared the same point predictions, and its limits differed near zero. After inspecting those results, we retained them and used 40,000 shared group resamples per method and source comparison. In a further post-hoc fixed-prediction check, deleting one connected group and recomputing the interval makes it cross or touch zero for 6 of 17 possible deletions. This is one of 16 reported post-hoc cells, with no multiplicity adjustment; rank representations share point estimates and are not independent tests. Candidate source membership, role assignments and sample size change together. It is evidence of sensitivity in these selected archives, not a causal provider or climate-transfer effect.

Independent checks compare every non-source-model column in 220 replay files (1,003 annual rows), confirm that strict and original predictions coincide outside HUC17, and reconstruct 48 summary cells, 144 fixed-prediction leave-region-out cells, all 16 paired source interactions and 68 leave-one-group resamples from saved predictions. These are numerical checks on shared observations and fitted models, not an independent climate, sensor or legal-rights evaluation. The first attempt to distribute tasks to eight processes failed to deserialize a task; the same plan completed serially with receipt checks.

S11. Source-model sensitivity of expanded stresses

Following inspection of the original source-role audit and expanded stress results, we replayed the stressed daily features and masks through two already fitted alternative source-model sets. The strict-original version replaces only the forward HUC17 model,

the sole affected target region in this cohort, with the model refitted after excluding every HUC17 connected group from its source roles; other source assignments remain as before. This removes source cases and changes source sample size. The candidate version replaces the models for all six forward target regions with the separately refitted candidate-provider models, which exclude all target-region groups but also change source providers, source sample size and role assignments. Source-model outputs and calibration offsets are recalculated; the previously fitted local daily features and target masks remain fixed. Neither comparison is prospective or a clean causal experiment for a single change.

Table S19 uses, separately for each estimator and stress, only archives available under both baseline and stress in all three source versions. The independent annual-rank intervals use the same six-way common interval cohort, with the same normal-spacing tail assumptions. D2 has 205 common archives; the stochastic method has 190 for donor gaps and 187 for the fixed-count heat stress. Repeated masks are not independent climates. Descriptive connected-group bootstrap and fixed-prediction region deletion results accompany the aggregate files. In the heat scenario the number of masked years is fixed within archive, but the number and placement of hidden days can change; in the donor scenario target masks are identical. These simulations do not verify a natural outage mechanism or transfer to new climates.

The source-role audit checked the positive-weight roles of the original and alternative models. All 440 alternative stress reanalyses preserved the target masks, hidden-value checks, plug-in values and observed lower bounds. The audit checked all 205 candidate baseline receipts against their source models, calibration, original features and saved outputs, and replayed the 15 affected strict-original baselines, including 60 annual predictions

Table S19. Source-model sensitivity on six-way common archives, baseline / stress. RMSE, width and score use °C per decade; coverage is the nominal 95% trend interval percentage. Strict changes only the affected HUC17 source model; candidate changes source providers and assignments across all regions.

Scenario	Estimator	Source	Cases	RMSE	Coverage	Width	Score
Donor gaps	D2	Original	205	0.1555 / 0.2024	92.2 / 86.8	0.465 / 0.479	0.756 / 1.039
Donor gaps	D2	Strict	205	0.1561 / 0.2027	92.2 / 86.8	0.467 / 0.475	0.758 / 1.050
Donor gaps	D2	Candidate	205	0.1514 / 0.2068	88.8 / 82.4	0.425 / 0.502	0.738 / 1.117
Donor gaps	Stochastic	Original	190	0.1386 / 0.1877	87.4 / 76.3	0.404 / 0.418	0.808 / 1.257
Donor gaps	Stochastic	Strict	190	0.1386 / 0.1877	87.4 / 76.3	0.404 / 0.418	0.808 / 1.257
Donor gaps	Stochastic	Candidate	190	0.1455 / 0.1969	86.8 / 76.3	0.392 / 0.461	0.799 / 1.364
Heat, fixed years	D2	Original	205	0.1555 / 0.1655	92.2 / 88.8	0.465 / 0.456	0.756 / 0.806
Heat, fixed years	D2	Strict	205	0.1561 / 0.1661	92.2 / 88.3	0.467 / 0.457	0.758 / 0.812
Heat, fixed years	D2	Candidate	205	0.1514 / 0.1642	88.8 / 86.8	0.425 / 0.410	0.738 / 0.877
Heat, fixed years	Stochastic	Original	187	0.1370 / 0.1552	87.7 / 87.2	0.403 / 0.383	0.776 / 0.884
Heat, fixed years	Stochastic	Strict	187	0.1370 / 0.1552	87.7 / 87.2	0.403 / 0.383	0.776 / 0.884
Heat, fixed years	Stochastic	Candidate	187	0.1443 / 0.1677	87.7 / 82.9	0.390 / 0.373	0.764 / 0.947

and 180 trend rows. A second implementation independently recomputed the six-way common cohorts, 24 aggregate cells, 144 region-deletion cells and 4,000 group bootstrap draws per cell from 9,840 saved case rows. These checks did not independently refit source models, check sensor accuracy or validate climate transfer.

S12. Exploratory projected-point failure score

After inspecting prior projection and risk results, we fitted six region-specific binary classifiers to identify absolute D2 point error above 0.5 °C after the observed-bound

projection. For each target region, the source estimator, interval-calibration, risk-learning and rule-calibration connected-group roles are disjoint, all groups appearing in the target region are excluded, and forward source cases precede 2010. Training uses the assigned risk-learning source groups, weighted equally by connected group. The new classifier uses the existing decision-time feature matrix, the projected 90% interval width and the projected point's distance above the visible bound; neither the old risk score nor the held reference is a predictor. Each model uses 120 low-leaf-count boosted trees with the fixed settings documented in the analysis plan. Six to eight source training groups contribute per model.

For each region and mask family, separate source rule-calibration groups select an old-score and a new-score cutoff to approximate 80% group-equal-weight numerical retention on valid source cases, without using failure labels to choose the cutoff. Both cutoffs are fixed before target scoring. Target numerical retention can differ. A second, transductive comparison ranks the new target scores and matches the old-score numerical count separately within each target region and family; ties are broken deterministically without target outcomes. It is not a prospective cutoff, and equal counts need not retain the same years or bound values.

The candidate annual caches contained no reference H30 values. Therefore source models, source cutoffs, target scores and selections were saved first; the target reference was then joined by station, masked year, family and replicate from the separately saved complete-archive record. A second comparison against the original source station–year–family H30 verified all 7,583 reference values, and the saved selections did not change. These target

archives and source data had nevertheless been inspected in earlier analyses; this is not an unopened reference experiment.

Table S20 gives pooled forward results. Source thresholds in the controlled-gap family give old/new numerical retention 89.6% / 85.3% and selected failure 10.72% / 9.50%. At the matched 3,385 outputs, the original and projected-risk scores have 363 and 383 failures, respectively; the connected-group bootstrap interval for the difference in selected failure rates (new minus old) is [-0.73, 1.86] percentage points. Group resampling is descriptive and repeated masks are not independent climates. The old classifier targeted an unprojected point; the new classifier targets the projected point, but a better-aligned training label does not itself imply better deployment decisions. The source masks degrade one year at a time, while target archives are jointly degraded across years. These results do not establish calibrated probabilities, group-level risk control, natural-outage performance or independent-climate transfer.

Table S20. Forward annual failure comparison, old raw score / projected-point score. An error exceeds 0.5 °C. Source-threshold counts can differ; the target count match is transductive and uses no target outcomes.

Mask	Selection	Cases	Numerical	Failures	Failure (%)
Calendar	Source threshold	3,806	3,431 / 3,112	6 / 6	0.17 / 0.19
Calendar	Target count match	3,806	3,431 / 3,431	6 / 6	0.17 / 0.17
Controlled	Source threshold	3,777	3,385 / 3,222	363 / 306	10.72 / 9.50
Controlled	Target count match	3,777	3,385 / 3,385	363 / 383	10.72 / 11.31

The independent audit refits the six classifiers using the same source records and feature implementation, rechecks the disjoint roles, reproduces the twelve family-specific source threshold pairs, reconstructs all target scores and label-free selections, and recomputes group bootstrap and region-deletion summaries. It also checks reference attachment against both saved reference sources. It does not independently reconstruct physical observations, establish source permissions or verify the classifier's prospective reliability.

S13. Seasonal-window and screening-rule diagnostics

The previously selected source sample contains 96 eligible long-record stations; this calculation uses the 95-station project-processed subset with a documented candidate redistribution path. None of the 35 recorded witnesses is at the excluded LUBW station, although it remains part of the original eligible-station denominator. The subset comprises 2,122 complete station-years at 95 stations and 33 connected groups. For each complete year, we compare the full-year reported H30 with the largest 30-day mean whose every day lies from 1 July through 31 August. Of those years, 779 exceed the summer-window value by more than the numerical tolerance 5×10^{-6} °C, and 304 exceed it by more than 0.5 °C. The mean difference is 0.255 °C. These are complete recorded years, not simulated reconstruction errors; neither the tolerance nor the 0.5 °C threshold is an estimate of sensor error.

The comparison was specified after the original witnesses and literature had been examined. It is not a reproduction of Beaufort et al.'s multiyear thermal peak or their summer air-temperature climate correction; that study also excluded sites with strong human influence. The present sample contains regulated and incompletely characterized

sites, covers different locations and years, and the fully-contained July–August window differs from a peak-center criterion. Consequently our fraction must not be compared directly with their reported fraction. The incomplete-year comparison is also conditional on the 35 already selected record witnesses: all have a fully observed July–August 30-day window, and 9 have a larger fully observed 30-day window elsewhere in the year. This does not establish the full H30 of an incomplete year or a population prevalence.

Table S21 counts previously eligible station-years rejected by four explicit study rules, including how many still supply at least one fully observed 30-day window. A lower bound is information about the reported record, not a complete annual estimate or a guarantee of better trends. The 90% station-specific warm-season rule uses a leave-target-year-out climatology; the complete July–August row is a contextual comparator rather than the full published climate-correction procedure. The records are not an independent validation sample and no sensor error has been estimated.

Table S21. Study screening rules on 2,656 eligible station-years in the 95-station candidate subset. Rejected-with-bound means the rejected incomplete year contains at least one complete observed 30-day window.

Rule	Accepted	Rejected	Unknown	Rejected with bound	Stations
Complete year	2,122	534	0	518	70
Annual 95%	2,550	106	0	90	48
Station warm season 90%	2,590	66	0	50	30
Complete July–August	2,472	184	0	168	58

After inspecting the largest annual gaps, we separately summarized existing project regulation labels and removed the two sites with the largest individual gaps (Table S22). The 2010 USGS analysis identified Blue River and Fall Creek as dam-affected monitoring sites in 2001–2002; it does not establish the cause or measurement quality of every year examined here. The existing *unmatched gages* label does not mean unregulated, and labels and sites were not randomized. Removing these two post-hoc sites leaves 749 of 2,086 complete station-years above the numerical tolerance, with mean difference 0.197 °C. These summaries do not identify causal dam effects or transfer to streams absent from the sample.

Table S22. Post-hoc seasonal-window influence by existing project label and after removing two selected sites. "Above" is greater than 5×10^{-6} °C; the second count exceeds 0.5 °C. Labels are observational metadata, not independently verified treatments.

Cohort	Stations	Years	Above	> 0.5	Mean (°C)
All stations	95	2,122	779	304	0.255
Exclude two selected sites	93	2,086	749	279	0.197
Label: regulated	23	397	173	103	0.725
Label: unmatched gages	61	1,542	569	196	0.158
Label: unknown	11	183	37	5	0.049

The comparison and screening totals were independently reconstructed using pandas rolling windows on the same snapshot. The post-hoc influence totals and 36 selected-site

peak-window dates were independently checked using NumPy convolution; the stored regulation labels were checked against the project's source metadata. Neither audit verifies the original provider's measurements or whether each project regulation label is accurate. The supplied 95-station snapshot permits the window and screening arithmetic to be rerun, subject to its attribution and reuse conditions.

S14. Literature-inspired interannual thermal-peak comparison

After inspecting earlier results and data availability, we selected all 22 of the previously eligible 95 stations with complete reported water years from 2009 through 2017. They belong to 10 connected groups; project labels mark one as regulated and leave the others unmatched to the regulation inventory, not necessarily natural. Following the structure described by Beaufort et al. (`https://doi.org/10.5194/hess-26-3477-2022`), we examine a nine-year mean of annual 30-day peaks and fill missing summer daily water with station-specific regressions on right-aligned 2–10-day air-temperature means. In each synthetic experiment, we hide 1–8 whole years in every possible combination, select the lag with the highest in-sample R^2 using retained summers only, and fit the regression without withheld-year water. We calculate the 30-day maximum wholly contained in July–August for observed and predicted years. All nine years hidden is unavailable because no local fitting data remain. The 11,220 station–mask cases reuse only 22 stations and are not independent climate replicates.

The primary reference is the mean of the nine observed July–August 30-day maxima. A separately reported reference is the mean of the nine full-calendar-year H30 values; this changes the target, so any summer-only estimator may omit a peak outside July–August. The unfilled comparison averages only the retained years; the filled comparison averages

retained observed peaks and reconstructed missing-year summer peaks. Both methods are evaluated on the same cases. These are point estimates of interannual averages, not annual H30 recovery, trend, interval calibration or release-rule validation. The observed-only mean bias across the exhaustive symmetric masks is zero for the summer reference by construction; this does not establish unbiasedness under real missingness.

Table S23 reports station-equal MAE and 4,000 connected-group bootstrap intervals for the filled-minus-observed absolute-error difference, with one row per station in the group resample. The comparison is post-hoc and uses ten project connected groups; the resampling interval is descriptive and depends on those groups. Other missing-year counts and leave-one-provider-out results are in the aggregate output. Removing a provider does not refit models or establish independent climates.

This is an adaptation, not a reproduction of the French study: it uses a different station population, processed water records, NASA POWER-based daily air instead of SAFRAN, explicit train-only fitting, all combinations of hidden years, and a July–August-contained peak definition. The published article does not establish all of our implementation choices. NASA POWER's gridded local-solar air days may not align with provider water days; the original API version and raw response for these 22 cached stations were not retained. After viewing the proxy results, we requested current NASA POWER responses for 15 representatives of distinct stored air series. All 9,585 comparable summer and lag-input days matched within 10^{-5} °C. This does not establish historical version, unrequested stations, air or water accuracy, clock alignment or independent-climate generality. A separate implementation recomputed 11,220 masks and all 16 summary cells from the same processed inputs; it did not reproduce the original authors' datasets or code.

Table S23. Adapted summer-regression proxy against two distinct nine-year targets. MAE and filled-minus-observed MAE differences are °C; cases are repeated year masks.

Reference	Hidden years	Cases	Observed MAE	Filled MAE	Difference	Group interval
Summer	1	198	0.100	0.069	-0.031	[-0.051, -0.020]
Summer	2	792	0.156	0.112	-0.043	[-0.070, -0.028]
Summer	4	2772	0.266	0.189	-0.078	[-0.114, -0.057]
Summer	6	1848	0.418	0.285	-0.133	[-0.184, -0.103]
Summer	8	198	0.803	0.530	-0.273	[-0.416, -0.192]
Full-year	1	198	0.190	0.192	0.002	[-0.016, 0.012]
Full-year	2	792	0.222	0.223	0.001	[-0.030, 0.018]
Full-year	4	2772	0.305	0.288	-0.017	[-0.068, 0.012]
Full-year	6	1848	0.437	0.370	-0.067	[-0.143, -0.023]
Full-year	8	198	0.805	0.581	-0.224	[-0.390, -0.127]

S15. Direct archive comparison with a literature-inspired summer air proxy

After viewing the original archive and interannual-proxy results, we evaluated a different, deterministic adaptation on the same 3,320 jointly masked archives. For each target year, all target-archive gaps are hidden before fitting, the entire target year is excluded from the local summer water–air fit, and forward fits exclude water from 2010 onward. We require at least three complete eligible training years, 30 usable summer days from at least three years, and common finite predictors for all 2–10-day right-aligned air averages; the lag maximizing in-sample R^2 is selected. We fill missing July–August water days and take the maximum of the reconstructed summer 30-day peak and the fully observed full-year

30-day bound. When summer contains no gaps, the result is the observed full-year bound alone. Missing air and unsupported fits remain unavailable. This differs in data, target and implementation from Beaufort et al. (https://doi.org/10.5194/hess-26-3477-2022); it is not their code or a test of their original multiyear thermal-peak estimator.

There are 21,446 masked station-year records in the repeated archive settings; 21,278 proxy outputs are available. Of all records, 11,334 have no summer gaps and return only the observed bound; 14,784 available outputs equal that bound, including cases where a regression was fitted. A bound can remain below the true annual maximum when the missing peak is outside July–August. These are repeated synthetic masks, not independent site-years or climates. The source roles, pre-existing fixed models, air-day alignment and restricted-data access limitations remain. The proxy supplies no predictive interval. Annual retrospective strata and pairwise availability are reported in the aggregate output; the bound-equals-reference stratum uses the hidden reference and cannot be selected at deployment.

Table S24 uses cases where all four point methods are finite. Trend rows exclude archives with no masked year and use the same finite-record OLS target. Differences are descriptive after selecting this analysis; resampling connected groups and fixed-prediction leave-one-region-out summaries do not establish out-of-region prediction or causal superiority. Independent least-squares and rolling-window code checked 52 selected archives in seven regions, 310 masked years and target-year fit perturbations, plus the reported aggregate cells. It used the same processed sources and fixed baseline predictions.

Table S24. Common-cohort annual and nontrivial-archive point comparisons. Annual MAE is °C; trend RMSE is °C per decade. Counts are repeated mask records or archives.

Setting	Annual n	D2 MAE	Proxy MAE	Archive n	D2 RMSE	Proxy RMSE
Calendar, offline	6,891	0.018	0.037	822	0.009	0.018
Calendar, forward	3,676	0.029	0.047	699	0.062	0.083
Long gap, offline	6,786	0.165	0.294	818	0.028	0.068
Long gap, forward	3,683	0.225	0.361	736	0.164	0.237

S16. Past-only warm-season sensitivity

The original station-specific warm season leaves out the evaluated year but can use observations from later years. After inspecting that comparison and the 35 selected witnesses, we recomputed the same 15-day-smoothed day-of-year climatology and five-month selection with only daily values from years strictly before each evaluated year. The source is the current processed archive, not a historically timestamped data vintage. The 95 stations and 35 witnesses were themselves selected using records available after some evaluated years, so this is a retrospective information-separation sensitivity, not a prospective site recruitment or field performance test.

Two past-only variants are reported. One uses any earlier finite water observation, which can amount to only 1 day for an incomplete year; climatological interpolation with so little data is not evidence of adequate support. The other requires at least four fully observed prior calendar years before classifying an incomplete year. Complete years are accepted even if the climatology is unavailable, because every possible warm season is fully observed. The four-year threshold is an exploratory study choice, not a published

standard. Unsupported incomplete years remain unknown rather than being counted as rejections. On 2,656 eligible station-years, the past-only variant without a minimum changes 1 decision relative to the original leave-year-out rule; the four-year version leaves 127 years unknown, 122 of which still have an observed 30-day lower bound.

Table S25 also compares the reported observed-window maximum in each rejected year to the greatest reported window from earlier years accepted by that same rule. A rejected year contributes only if both its observed bound and an earlier accepted reported window exist. Earlier accepted incomplete years can have an unobserved higher H30, and these are not comparisons with physical sensor truth. The leave-year-out rule's earlier classifications may use years later than their evaluation year; only the past-only rules enforce the stated temporal information boundary. Among the 35 retrospectively selected witnesses, the four-prior-complete-years variant accepts 27, rejects 2 and leaves 6 unknown; 2 rejected witnesses exceed the earlier accepted reported-window maximum. These counts are not an incidence estimate.

Table S25. Retrospective screening in the 95-station candidate subset. "Above prior" counts rejected years whose observed 30-day window exceeds all earlier accepted reported windows. Witness column gives accepted / rejected / unknown among 35 previously selected incomplete years.

Rule	Accepted	Rejected	Unknown	Rejected with bound	Above prior	Witnesses
Complete year	2,122	534	0	518	86	0 / 35 / 0
Annual 95%	2,550	106	0	90	11	31 / 4 / 0
Warm 90%, leave year out	2,590	66	0	50	9	32 / 3 / 0
Warm 90%, past data	2,589	67	0	51	9	32 / 3 / 0
Warm 90%, past + 4 complete	2,487	42	127	31	3	27 / 2 / 6

An independent implementation rebuilt all 2,656 earlier-only climatologies, 13,280 rule states and prior-record comparisons on the same processed daily records. In 35 witness-year probes, shifting the target and all later observations by 5 °C did not change the earlier-only warm season. These checks do not verify historical data vintages, station selection, provider quality control or physical measurement accuracy.

S17. Conservative source-group maximum sensitivity

After earlier annual, trend and classifier results had been inspected, we combined the candidate source model's assigned risk-learning and rule-calibration connected groups for each target region. These groups are disjoint from the numeric quantile estimator and its interval-offset groups and exclude groups in the held target region. The risk-learning groups trained a separate failure classifier, but no classifier score or selection is used in this interval analysis. Replacing the stored classifier with a constant score in 300 source-case probes left both methods' seven annual quantiles unchanged. Source cases are before 2010 and use single-year masks. The source and target observations were already opened; this is a post-hoc sensitivity, not a new prospective experiment.

For each valid source group, we calculate the maximum of the nonnegative distances by which a reference falls outside its projected 90% interval. With G groups, the expansion is the $\lceil 0.9(G+1) \rceil$-th ordered group maximum if that rank exists and is positive; otherwise expansion is unsupported. The six forward regions have 12–16 contributing groups per method and family, so the chosen rank is the maximum among the available group maxima. This finite order statistic does not establish a 90% group-coverage guarantee: exchangeability is not demonstrated, source cases use individual-year masks, target archives have joint multiyear gaps, and groups have different sizes and climates. Both methods'

lower endpoints are constrained to the visible observed window. Table S26 compares the same target cases; the hidden-peak subset uses the complete reference and cannot be selected at decision time. All evaluated cases in each of the 17 target groups happen to be covered after expansion in the controlled-gap cohort, but those groups are not independent-climate replications.

Table S26. Nominal 90% annual intervals in common forward controlled-gap cases, baseline / source-group-maximum expansion. Width and interval score are °C; lower score is better.

Method	Stratum	Cases	Coverage (%)	Width	Score
D2	All	3,683	89.2 / 100.0	0.734 / 4.690	1.442 / 4.690
D2	Hidden peak	2,146	81.9 / 100.0	1.161 / 5.429	2.349 / 5.429
Stochastic	All	3,683	87.3 / 100.0	0.650 / 5.793	1.476 / 5.793
Stochastic	Hidden peak	2,146	78.2 / 100.0	1.078 / 6.525	2.496 / 6.525

We then propagated the same fixed source radii through the original masked archives and three rank constructions. Of 820 settings, 82 lack at least one annual distribution and 2 have no masked year; 736 nontrivial settings in 17 connected groups support both methods and all rank constructions. Two hypothetical quantile representations share the same expanded 5th and 95th percentiles: one changes only endpoints, and the other also tapers changes through interior quantiles. They preserve the same point median and use identical ranks, but their interpolated interior and normal-spacing tails differ. Table S27 shows empirical nominal 95% trend coverage, width and interval score. On the 685 archives

with the earlier narrow source correction and all ranks, the independent-rank tapered D2 interval width is 0.403 with that earlier correction and 2.212 °C per decade with group maxima; the corresponding stochastic widths are 0.376 and 2.747. Matching the annual endpoints does not identify the trend distribution, and observed coverage on these selected masks does not certify coverage for future archives.

Table S27. Nominal 95% forward trend intervals on 736 common nontrivial archives. Each entry is baseline / endpoint-only / tapered; widths and interval scores are °C per decade.

Method	Ranks	Coverage (%)	Width	Score
D2	Independent	88.6 / 100.0 / 100.0	0.399 / 3.102 / 2.222	0.735 / 3.102 / 2.222
D2	Shared parameter	88.7 / 100.0 / 100.0	0.395 / 3.035 / 2.178	0.732 / 3.035 / 2.178
D2	Residual blocks	88.9 / 100.0 / 100.0	0.400 / 3.022 / 2.229	0.714 / 3.022 / 2.229
Stochastic	Independent	84.5 / 100.0 / 100.0	0.366 / 3.968 / 2.751	0.896 / 3.968 / 2.751
Stochastic	Shared parameter	83.4 / 100.0 / 100.0	0.362 / 3.894 / 2.700	0.902 / 3.894 / 2.700
Stochastic	Residual blocks	83.8 / 100.0 / 100.0	0.369 / 3.881 / 2.781	0.882 / 3.881 / 2.781

The independent annual audit reconstructed 88,053 source case–method predictions, 352 source group maxima, 14,714 target case–method scores and the annual summary cells. An isolation check identified a previously implicit pre-2010 bridge-repair table needed for the stochastic source cases. The original source plan did not list this file: applying the saved table reproduces all source group maxima, while omitting it changes source-case availability and three of the twelve stochastic region–family radii; D2 radii are unchanged. The audit now requires the table explicitly and fails if it is absent or changed. A separate calculation reconstructs 13,248 trend treatment rows from saved quantiles and rank arrays

and checks all 685 overlapping earlier rank snapshots. These numerical audits use the same source and target observations and do not validate exchangeability, physical sensors, provider rights or an independent climate.

Further post-hoc replay identified bridge-repair dependencies in the twenty historical source-seed exports and in the candidate-provider archive calibration. Reapplying the twenty saved repairs reproduced the stored D2 and stochastic quantiles on all 2,914,470 seed–case records. Without those repairs, 1,081,252 records lost complete stochastic quantiles and 1,018,696 records with complete quantiles in both replays had at least one changed quantile. These counts include repeated masks and are not independent station-years. In the archive calibration, applying the two saved repairs reproduced all 26 mode–region–method offset sets; omitting repairs changed offsets in all 13 stochastic sets and none of the 13 state sets. The original candidate-provider archive plan did not list these repairs; the later single-year plan did record these repairs. These dependency checks reuse saved cases and model roles and were assembled after the earlier analyses, not prospectively registered.

S18. Annual support and hidden-peak diagnostics

The projection-stage and peak-visibility tables use common finite cases. The peak-visibility stratum depends on the withheld complete reference and is unavailable for an operational decision. Repeated masks can place the same physical station-year in both strata.

The failure-strata table reports case means. The saved summary also contains group-equal means and group-bootstrap intervals, and the leave-one-region-out table repeats the D2-minus-classical comparison after removing one region at a time. These are descriptive summaries of the sampled archive, not independent-climate guarantees. The peak-visibility stratum uses the complete reference and is unavailable at decision time; the gap, overlap and donor bins were chosen after inspection.

Table S28. Replay of annual interval stages on common finite-bound cases. Raw is the original model output; stochastic raw quantiles use a finite set of imputations that retain observed days. First applies the first explicit bound projection; Pre-final applies the fixed projection-aware offsets and sorts; Final applies the bound again. New is the count first becoming zero-width at the final projection. Zero widths use 10^{-8} °C tolerance. These are repeated mask cases, and no alternative calibration was fitted.

Family	Setting	Estimator	n	Raw	First	Pre-final	Final	New
calendar	forward	D2	3674	0	2	1096	1476	380
calendar	forward	Stochastic	3674	3090	3090	2977	2977	0
calendar	offline	D2	6919	0	0	4402	4538	136
calendar	offline	Stochastic	6919	5818	5818	5482	5482	0
crossed	forward	D2	3683	0	0	227	323	96
crossed	forward	Stochastic	3683	1362	1362	1091	1091	0
crossed	offline	D2	6919	0	1	1376	1391	15
crossed	offline	Stochastic	6919	2491	2491	2030	2030	0

Table S29. Retrospective annual performance on the two-estimator common finite-bound cohort, stratified using the complete reference. Match means that the observed lower bound equals the complete reported H30 within 10^{-8} °C; excess means the reference exceeds that bound. This outcome-defined stratum is unavailable at deployment. A station-year can occur in both strata under different masks. Intervals in brackets resample connected groups with 2,000 repetitions and are descriptive under the sampled archive, not independent-climate guarantees.

Family	Setting	Stratum	Method	n	Groups	MAE	Coverage	Group interval
calendar	forward	Match	D2	3125	17	0.000	100.0%	[100.0, 100.0]
calendar	forward	Match	Stochastic	3125	17	0.000	99.9%	[99.6, 100.0]
calendar	forward	Excess	D2	549	15	0.191	88.3%	[84.6, 93.2]
calendar	forward	Excess	Stochastic	549	15	0.151	87.8%	[84.1, 93.5]
calendar	offline	Match	D2	5891	18	0.000	99.9%	[99.6, 100.0]
calendar	offline	Match	Stochastic	5891	18	0.000	99.9%	[99.8, 100.0]
calendar	offline	Excess	D2	1028	16	0.122	90.6%	[86.8, 92.9]
calendar	offline	Excess	Stochastic	1028	16	0.120	83.1%	[79.8, 88.7]
crossed	forward	Match	D2	1537	17	0.012	99.3%	[98.9, 99.8]
crossed	forward	Match	Stochastic	1537	17	0.025	94.4%	[89.1, 99.3]
crossed	forward	Excess	D2	2146	17	0.378	83.2%	[72.6, 93.0]
crossed	forward	Excess	Stochastic	2146	17	0.345	81.6%	[72.2, 91.7]
crossed	offline	Match	D2	2787	18	0.008	99.3%	[98.9, 99.6]
crossed	offline	Match	Stochastic	2787	18	0.005	99.7%	[99.5, 99.9]
crossed	offline	Excess	D2	4132	18	0.285	90.7%	[88.5, 93.6]
crossed	offline	Excess	Stochastic	4132	18	0.289	87.9%	[84.6, 91.3]

Table S30. Descriptive failure strata for the crossed forward controlled gaps. Peak visibility uses the withheld complete reference. "At least half" includes overlap equal to 0.5. Repeated masks are not independent rivers or climates, and small regional samples do not identify a climate mechanism.

Stratum	Method	Cases	Stations	Groups	MAE (°C)	90% coverage	Width (°C)
reference peak: reference peak above bound	D2	2,146	41	17	0.378	83.2%	1.30
reference peak: reference peak above bound	Classical	2,146	41	17	0.345	81.6%	1.12
reference peak: reference peak at bound	D2	1,537	41	17	0.012	99.3%	0.17
reference peak: reference peak at bound	Classical	1,537	41	17	0.025	94.4%	0.10
longest gap: 30 d	D2	1,234	41	17	0.157	92.6%	0.70
longest gap: 30 d	Classical	1,234	41	17	0.145	88.2%	0.58
longest gap: 31–60 d	D2	1,262	41	17	0.225	90.3%	0.84
longest gap: 31–60 d	Classical	1,262	41	17	0.213	88.5%	0.71
longest gap: 61–90 d	D2	1,187	41	17	0.295	86.9%	0.94
longest gap: 61–90 d	Classical	1,187	41	17	0.279	84.1%	0.80
training peak overlap: at least half	D2	1,874	41	17	0.350	87.1%	1.36
training peak overlap: at least half	Classical	1,874	41	17	0.332	82.2%	1.17
training peak overlap: no overlap	D2	1,232	41	17	0.028	97.1%	0.06
training peak overlap: no overlap	Classical	1,232	41	17	0.025	96.5%	0.04
training peak overlap: partial	D2	577	41	17	0.241	83.9%	0.70
training peak overlap: partial	Classical	577	41	17	0.218	82.1%	0.55

Table S31. Descriptive failure strata for the crossed forward controlled gaps. Peak visibility uses the withheld complete reference. "At least half" includes overlap equal to 0.5. Repeated masks are not independent rivers or climates, and small regional samples do not identify a climate mechanism. Continued: visible donors and region.

Stratum	Method	Cases	Stations	Groups	MAE (°C)	90% coverage	Width (°C)
visible donors: 0	D2	14	2	2	0.270	92.9%	0.94
visible donors: 0	Classical	14	2	2	0.327	85.7%	0.82
visible donors: 1–3	D2	1,012	17	13	0.194	89.4%	0.68
visible donors: 1–3	Classical	1,012	17	13	0.172	89.2%	0.65
visible donors: 4+	D2	2,657	29	8	0.236	90.1%	0.88
visible donors: 4+	Classical	2,657	29	8	0.226	86.1%	0.71
region: Switzerland (FOEN)	D2	1,119	10	2	0.100	98.7%	0.77
region: Switzerland (FOEN)	Classical	1,119	10	2	0.103	98.1%	0.62
region: Bavaria	D2	1,983	24	8	0.315	82.9%	0.85
region: Bavaria	Classical	1,983	24	8	0.279	77.9%	0.69
region: USGS HUC03	D2	92	1	1	0.072	100.0%	0.53
region: USGS HUC03	Classical	92	1	1	0.063	100.0%	0.56
region: USGS HUC04	D2	81	1	1	0.075	100.0%	0.98
region: USGS HUC04	Classical	81	1	1	0.098	100.0%	0.91
region: USGS HUC10	D2	180	2	2	0.233	94.4%	1.10
region: USGS HUC10	Classical	180	2	2	0.359	93.3%	1.09
region: USGS HUC17	D2	228	3	3	0.161	97.4%	0.75
region: USGS HUC17	Classical	228	3	3	0.140	96.5%	0.72

Table S32. Exploratory source-only annual interval expansion on two-method common cases. Supported cases have at least three disjoint source groups and a positive empirical correction; unavailable and zero-correction regions remain in the denominator. Before and after use exactly the same supported cases. The source-group-weighted empirical correction has no finite-sample or climate-transfer guarantee; it was explored after inspecting the principal results. Coverage is percent, width and interval score are in °C.

Family	Setting	Method	Eligible	Supported	Coverage	Width	Interval score
calendar	forward	D2	3676	0	–	–	–
calendar	forward	Stochastic	3676	0	–	–	–
calendar	offline	D2	6926	0	–	–	–
calendar	offline	Stochastic	6926	0	–	–	–
crossed	forward	D2	3683	92	100.0→100.0	0.532→0.543	0.532→0.543
crossed	forward	Stochastic	3683	92	100.0→100.0	0.555→0.567	0.555→0.567
crossed	offline	D2	6919	0	–	–	–
crossed	offline	Stochastic	6919	0	–	–	–

S19. Source role audit for partitioned forward models

We re-derived the seed-zero bootstrap weights and checked them against the roles saved in 98 model bundles for the four evaluated source splits. The selected held test stations and their recorded connected groups did not enter any source role. However, in the before-2010 split, the role function defines held groups from test-period cases; training-only groups in the same named region may therefore remain in the source. The table reports positive-weight same-region source cases, not held-target observations. This is not a uniformly held-region design; the distinction affects the interpretation of forward transfer. It does not establish leakage of target-station hidden values or physical independence across groups.

S20. Post-hoc strict-region sensitivity

We refitted the two affected before-2010 seed-zero models after removing every source connected group that appears in the named target region. The original assignment and bootstrap multiplicities of other groups were retained; no replacement groups were drawn. Source interval and release thresholds were recalculated from the filtered roles. In the whole-archive experiment, only the three affected HUC17 stations were reanalysed, on the same 120 original masks; all other original results were retained. Each strict run checked the original mask and the absence of hidden target values. Both estimators were compared on the same station–repetition cases available in both versions. For forward controlled gaps this common cohort comprises 738 cases at 41 stations. Original and strict D2 trend RMSE are 0.1638 and 0.1645; the corresponding stochastic RMSE values are 0.1466 and 0.1466 °C per decade.

The future-year release analysis replaces predictions in both affected regions while preserving all target cases and the other regional models. Across 27,846 forward controlled-

gap cases, the three-output rule publishes 63.1% and 64.5% in the original and strict comparisons, with 10.9% and 11.3% observed failures among the published values. In HUC17, publication changes from 62.7% to 71.6%, and the selected error rate from 6.0% to 8.2%. The selected sets differ, so the conditional rates cannot be interpreted without retention. The strict refits use fewer source observations; these paired comparisons do not identify the causal effect of an individual group and do not establish external validity or safety. They do not recompute every analysis in this paper under strict regional exclusion. Case counts contain repeated masks and are not independent climates.

The source-role comparison additionally checked held test station identities across all source regions, the positive bootstrap weights and the effective split of classifier and probability-calibration groups. Only the two before-2010 models changed positive source weights; zero-weight role membership still matters for downstream group-list use, so this does not prove every downstream result unchanged.

The release-then-trend pipeline was replayed for all 120 HUC17 settings with the original masks, source-specific strict limits and newly fitted model. Original annual states reproduced for 1,440 station–year records. Across the affected records, 39 observed-bound outputs become numerical estimates; the other observed-bound and estimate states remain unchanged. The original and strict pooled controlled-gap comparisons comprise 820 common station–repetition cases and give released-values-plus-bounds working-likelihood trend RMSE 0.2024 and 0.1986 °C per decade, respectively, with mean working interval widths 3.26 and 3.29. These working intervals include regression variation and do not establish calibrated missingness-only coverage.

In the outcome-defined stratum with an unobserved peak, the 2,146 common forward controlled-gap annual cases have D2 nominal 90% coverage 83.2% originally and 82.9% under the stricter source. This retrospective stratum is unavailable to a decision maker and repeated masks are not independent climate realizations. The affected HUC17 source rule-calibration pool contains at most one eligible connected group before model prediction, below the minimum of three for both annual expansion and task calibration; strict refitting cannot change that source-group upper bound, although the corresponding radius and errors were not recalculated.

For forward controlled gaps, the same 818 station–repetition cases from 17 connected groups have every annual, tail and rank-coupling output in both source versions. Under constant tails, nominal 95% D2 trend coverage is 83.0% originally and 83.3% strictly; normal-spacing tails yield 88.8% and 88.9%, with widths 0.451 and 0.454 and interval scores 0.860 and 0.862. In the separate normal-tailed rank experiment, independent ranks yield 88.0% and 87.9% coverage and residual blocks yield 89.0% and 89.0%. Original tail results reproduced on all 480 affected tail rows. These comparisons use the same random ranks within each tail or coupling experiment, but the tail and coupling experiments use different draws; they do not identify additive causes of undercoverage. Descriptive connected-group bootstrap intervals for strict-minus-original metrics accompany the aggregate results.

For the forward controlled-gap D2 versus stochastic comparison, 738 cases at 17 connected groups are available under both source versions and both estimators. The pooled D2-minus-stochastic MSE difference is +0.005350 originally and +0.005559 strictly; the descriptive connected-group bootstrap interval for the strict result is [+0.000957, +0.010431] $(^\circ\mathrm{C/decade})^2$. The same resampled groups are used for both source versions. Removing

Bavaria from these fixed predictions leaves 331 cases in 9 groups and changes the difference from -0.000143 originally to +0.000323 strictly. For empirical calendars the corresponding original and strict differences are -0.000213 and -0.000159 on 332 cases. These small changes can alter the sign of the regional comparison; the leave-one-region-out exercise does not refit source models and does not establish independent climates or significance.

The small controlled-gap stress comparison reanalyses 20 settings for one HUC17 station while retaining other stations' original results. Among 30 common baseline/stress station–repetition cases, original and strict baseline D2 RMSE are 0.0794 and 0.0803; for synchronized donor gaps they are 0.1164 and 0.1159, and for heat-associated gaps 0.0885 and 0.0902 °C per decade. Original scenario metrics and masks reproduce. The heat-associated scenario changes both the identity and potentially the number of masked years, so these are exploratory paired scenario comparisons, not isolated causal effects or independent climate tests.

A separate recalculation of the transductive, same-number numerical-output comparator under the strict source model publishes 17,961 controlled-gap values: three-output and score-only failure rates are 11.26% and 7.15%, respectively. The score threshold uses the target feature distribution and the three-output count; counts and selected cases differ between source versions, and lower-bound information is not held equal.

Table S33. Seed-zero before-2010 model roles with positive-weight source cases in the named target region. Bootstrap-weighted counts include repeated role draws and do not represent independent samples.

Region	Role	Groups	Stations	Cases	Weighted cases
nve_hydapi	interval_calibration	2	3	431	695
usgs_huc17	estimator	1	1	140	280

S21. Source-provider dependence of whole-archive models

The 42 whole-archive target stations belong to providers with recorded reuse paths, but source-model training has additional provenance. Recomputing the source role assignments and checking them against all 13 saved seed-zero model bundles shows that 13 estimator roles and 6 interval-calibration roles have positive-weight LUBW source cases. Source cases are repeated masks; their counts are not independent observations. Positive source weights establish model dependence but do not resolve the legal status of any particular derivative. Individual source, model, target, mask and reviewer-access rights still require confirmation. A post-hoc candidate-provider strict-region refit is reported separately and does not replace the main results. A subsequent post-hoc audit reconstructed donor visibility for all 194,370 eligible candidate source-case rows using the saved masks and processed panels, and checked the counts against the stored cases. In 39 of 52 mode–region–role cells, some positive-weight rows had a finite donor-regression estimate while a visible donor was absent from the original three local catalogs. These are repeated rows, not independent sites or evidence of unauthorized use. The 73 such USGS panel-column occurrences correspond to 69 identifiers, all returned by the USGS site metadata service; one GKD column has a matching project-generated network-QC record. The independent check re-evaluated 156 selected source cases and all 52 role cells. Site-number existence and project QC metadata do not establish the original provenance of cached temperature values, complete processing lineage, or permission to redistribute derived models. The public site metadata snapshot can be replayed separately; source panel mappings remain local. After the metadata audit, we retrieved current USGS daily-mean water temperatures for those 69 sites. Of 293,851 finite local panel-column days, 292,285 had a single distinct finite USGS value and all

agreed within 10^{-6} °C. Another 1,525 had multiple distinct USGS values; every local value matched at least one alternative, but that post-hoc match does not determine which series was used historically. Their median within-day range was 0.2 °C; 41 local days had no finite returned value. These are repeated panel-column days, not independent site-days. A separate implementation recomputed all comparisons. The current public responses are included for source-only replay; they do not establish historic data vintage, sensor quality or derivative permissions.

S22. Candidate-provider, strict-region source sensitivity

After observing the original results and LUBW source dependence, we refitted all 13 whole-archive source models on seven candidate providers with recorded reuse paths. Role assignments and bootstrap draws were recomputed using the filtered data, and every group occurring in the target region was excluded from source roles. An independent role recount found no positive-weight target-region source case or post-2009 forward source case. Original target masks and daily features were reused. All 3,320 archive settings, masks and the unchanged plug-in baselines were checked. Removing providers, changing source assignments and strict regional exclusion jointly change the analysis; the result is not a causal attribution to one of them. The filtered inputs remain processed study data, not provider-native raw records. This after-the-fact sensitivity neither establishes legal redistribution rights nor independent climate transfer, and it does not replace the original analysis. A descriptive connected-group bootstrap resamples the same groups for both source versions and both estimators. It does not establish independent climates or a population transfer guarantee. The corresponding paired MSE differences and bootstrap intervals are supplied with the candidate-source summary.

Table S34. LUBW cases with positive source-role weights, not independent station counts. Groups with bootstrap multiplicity zero do not contribute.

Mode	Region	Estimator case rows	Interval calibration case rows
forward	foen	383	0
forward	gkd_bayern	287	0
forward	usgs_huc03	383	0
forward	usgs_huc04	96	0
forward	usgs_huc10	191	96
forward	usgs_huc17	191	0
offline	foen	288	311
offline	gkd_bayern	862	2732
offline	syke	1030	311
offline	usgs_huc03	3475	0
offline	usgs_huc04	1461	168
offline	usgs_huc10	1150	0
offline	usgs_huc17	1246	2732

Table S35. Trend RMSE in °C per decade on the four-way common cases: both methods and both source-model versions must be available. Repetitions are not independent climates.

Family	Mode	Cases	Groups	D2 orig.	Class. orig.	D2 cand.	Class. cand.
calendar	forward	701	17	0.06178	0.05134	0.06033	0.05822
calendar	offline	835	18	0.00873	0.01124	0.00890	0.01119
crossed	forward	738	17	0.16381	0.14658	0.15806	0.15600
crossed	offline	838	18	0.02780	0.03208	0.02868	0.03195

Table S36. Nominal 95% D2 trend-interval results on cases where both versions are available. Width is in °C per decade; this comparison is descriptive and the selected cases may differ from the four-way point cohort.

Family	Mode	Cases	Cover orig.	Cover cand.	Width orig.	Width cand.
calendar	forward	820	96.0%	95.7%	0.116	0.115
calendar	offline	840	95.0%	95.2%	0.018	0.018
crossed	forward	820	88.0%	87.6%	0.450	0.408
crossed	offline	840	94.0%	92.9%	0.089	0.086

Table S37. Symmetric MSE decomposition on cases with all four retention-by-estimator cells available under both versions. Fractions divide the mean retention contribution by mean total improvement; they are not causal or universal.

Family	Mode	Estimator	Cases	Groups	Original fraction	Candidate fraction
calendar	forward	D2	820	17	0.799	0.783
calendar	forward	classical	701	17	0.744	0.767
calendar	offline	D2	840	18	0.887	0.891
calendar	offline	classical	835	18	0.931	0.930
crossed	forward	D2	807	17	0.939	0.937
crossed	forward	classical	725	17	0.925	0.931
crossed	offline	D2	840	18	0.930	0.933
crossed	offline	classical	838	18	0.942	0.942

S23. Annual uncertainty under candidate-source refits

The follow-up compares the same target cases where both methods and both source versions have finite seven-quantile distributions and a finite observed bound. The original 24 annual summary cells were independently reproduced on their original cohorts. Target identities, bounds and connected groups agree between versions. The decomposition by whether the complete reference exceeds the visible bound uses the withheld reference and is unavailable at decision time. Candidate-versus-original differences are post hoc; changed source providers, role assignments and strict region exclusion cannot be separated. The quoted CRPS uses the stated normal-spacing tail continuation, and no new calibration or independent climate validation is established. For forward controlled gaps with hidden peaks, the paired candidate-minus-original classical coverage difference is -3.4 percentage points with descriptive connected-group bootstrap interval [-8.6, 2.0]. Removing FOEN or Bavaria from the fixed evaluation cases changes that difference to +0.3 or -8.9 percentage points, respectively. Regions were not refitted in this deletion, and connected groups are not proven independent climates.

S24. Single-year source expansion for annual intervals

After reviewing prior annual and trend results, we used single-year masked source cases from the candidate models' rule-calibration groups. These groups do not enter the estimators, original interval calibration or risk-learning roles; every target-region group is excluded. Some groups contributed to the separate risk-rule component; the pool is not an unopened independent dataset. Forward source cases are from before 2010. For each family, mode, region and estimator we estimate an equal-group-weighted empirical 90th percentile of nonnegative interval misses. A correction is applied only with at least three

Table S38. Annual 90% interval comparison on the same four-way common cases. Hidden peak means that the complete reported H30 exceeds the observed bound; this is an outcome-defined stratum. Widths are in °C. Repeated masks are not independent climates.

Family / mode	Stratum	Method	Cases	Groups	Cover old	Cover new	Width old	Width new
calendar / forward	all	D2	3674	17	98.3%	97.9%	0.118	0.135
calendar / forward	all	classical	3674	17	98.1%	97.9%	0.088	0.093
calendar / forward	hidden	D2	549	15	88.3%	86.3%	0.690	0.680
calendar / forward	hidden	classical	549	15	87.8%	86.7%	0.566	0.610
calendar / offline	all	D2	6919	18	98.5%	98.5%	0.095	0.093
calendar / offline	all	classical	6919	18	97.4%	97.7%	0.066	0.069
calendar / offline	hidden	D2	1028	16	90.6%	90.8%	0.573	0.578
calendar / offline	hidden	classical	1028	16	83.1%	85.1%	0.427	0.434
crossed / forward	all	D2	3683	17	90.0%	89.2%	0.826	0.734
crossed / forward	all	classical	3683	17	87.0%	87.3%	0.693	0.650
crossed / forward	hidden	D2	2146	17	83.2%	81.9%	1.299	1.161
crossed / forward	hidden	classical	2146	17	81.6%	78.2%	1.121	1.078
crossed / offline	all	D2	6919	18	94.1%	92.8%	0.730	0.701
crossed / offline	all	classical	6919	18	92.7%	92.4%	0.660	0.639
crossed / offline	hidden	D2	4132	18	90.7%	88.6%	1.171	1.127
crossed / offline	hidden	classical	4132	18	87.9%	87.4%	1.068	1.037

available source groups and a positive radius. This rule is exploratory, and the source cases hide a single year rather than an entire degraded archive. It neither establishes group exchangeability nor identifies a calibrated joint trend distribution. At most eight forward groups are assigned to this role, below the nine needed for a finite one-maximum-per-group 90% rank; the group-maximum diagnostic was not applied. Re-prediction of the 52 source support cells, verification of individual target intervals and a separate two-method support-intersection check passed. Group-resampling and leave-region-out summaries are supplied; they do not prove independent climates. Unsupported cells and unchanged all-case baselines remain visible. The source group-mass empirical percentile is not a finite-sample coverage guarantee.

S25. Trend sensitivity to annual interval expansion

After examining the source-only annual correction and the earlier trend results, we propagated it through the original 256-draw normal-spacing quantile representation. The endpoint-only scheme moves the 5th and 95th quantiles while holding the five interior quantiles fixed. The tapered scheme also shifts the 10th, 25th, 75th and 90th quantiles by 0.8, 0.4, 0.4 and 0.8 times the radius, respectively; the median and complete-year values are unchanged. Both schemes use the same adjusted 90% endpoints and respect observed lower bounds. Their interior and normal-spacing outer tails differ. The same independent, auxiliary shared-parameter and residual-block rank matrices are reused within each scheme; no calibrated joint distribution is implied. Among 820 forward controlled-gap station–repetition settings, 760 belong to regions with source support for both methods. Seventy-three of these lack at least one annual distribution, and two have no incomplete year, leaving 685 common nontrivial settings. Original saved trend intervals are reproduced

Table S39. Exploratory interval expansion: same target cases must be supported for both methods. Baseline and expanded scores use the identical supported subset; all-case baselines are retained separately in the supplied CSV. Hidden-peak strata depend on the complete reference. Widths are in °C.

Family / mode	Stratum	Method	Eligible	Supported	Groups	Cover before	Cover after	Width before	Width after
calendar / forward	all	D2	3674	0	0	–	–	–	–
calendar / forward	all	classical	3674	0	0	–	–	–	–
calendar / forward	hidden	D2	549	0	0	–	–	–	–
calendar / forward	hidden	classical	549	0	0	–	–	–	–
calendar / offline	all	D2	6919	0	0	–	–	–	–
calendar / offline	all	classical	6919	0	0	–	–	–	–
calendar / offline	hidden	D2	1028	0	0	–	–	–	–
calendar / offline	hidden	classical	1028	0	0	–	–	–	–
crossed / forward	all	D2	3683	3411	14	88.7%	89.9%	0.705	0.755
crossed / forward	all	classical	3683	3411	14	86.6%	89.1%	0.608	0.696
crossed / forward	hidden	D2	2146	1982	14	81.0%	82.9%	1.110	1.171
crossed / forward	hidden	classical	2146	1982	14	76.9%	81.3%	1.009	1.115
crossed / offline	all	D2	6919	0	0	–	–	–	–
crossed / offline	all	classical	6919	0	0	–	–	–	–
crossed / offline	hidden	D2	4132	0	0	–	–	–	–
crossed / offline	hidden	classical	4132	0	0	–	–	–	–

before adjustment. The target selection and calibration remain retrospective. Connected-group paired bootstrap and fixed-prediction leave-one-region-out summaries are supplied. They are descriptive, use only 14 target groups and do not establish independent climates. Annual 90% endpoint correction does not identify the inner or outer distribution, cure annual undercoverage, guarantee trend coverage, or resolve source rights. The underlying case records and rank arrays remain local pending permission.

Table S40. Nominal 95% finite-record trend interval comparison on the same 685 station–repetition cases, 38 stations and 14 connected groups. Width and interval score are in °C per decade; repeated masks are not independent climates.

Method	Rank construction	Marginal scheme	Coverage	Width	Interval score
D2	independent	baseline	87.9%	0.382	0.742
D2	independent	endpoints	89.3%	0.409	0.736
D2	independent	tapered	88.9%	0.403	0.735
D2	shared parameter	baseline	88.0%	0.379	0.741
D2	shared parameter	endpoints	89.2%	0.405	0.730
D2	shared parameter	tapered	89.2%	0.399	0.733
D2	residual blocks	baseline	88.2%	0.385	0.721
D2	residual blocks	endpoints	89.8%	0.411	0.708
D2	residual blocks	tapered	89.5%	0.406	0.707
classical	independent	baseline	83.4%	0.344	0.912
classical	independent	endpoints	86.9%	0.387	0.856
classical	independent	tapered	85.5%	0.376	0.866
classical	shared parameter	baseline	82.2%	0.342	0.922
classical	shared parameter	endpoints	86.3%	0.383	0.863
classical	shared parameter	tapered	85.0%	0.373	0.872
classical	residual blocks	baseline	82.6%	0.349	0.901
classical	residual blocks	endpoints	86.4%	0.391	0.845
classical	residual blocks	tapered	85.7%	0.381	0.853

S26. External reference qualification

The Greater Wellington intake returned records for nine selected sites. Under the originally specified quality-code-600 criterion no complete reference hot year was qualified; the provider describes code 200 as unverified, while other returned legacy code meanings remain unresolved. A NEON PRT archive supplied half-hourly records at 24 river or stream sites. With one fixed location code per site, no site supplied four complete years under either 48 quality-passing intervals per UTC day or the 46-of-48 sensitivity. The latter is an observed-interval average rather than a complete physical day; a location code does not establish unchanged hardware.

After the original intake and quality screen, we separately tested an optimistic time-support bound on the same nine cached Greater Wellington records: all quality flags were ignored, while the original fixed-NZST cadence and fully observed day rule were retained and finite values required. Across these records there were 10 potentially complete July–June years, with zero to two per site; no site reached four. This post-hoc feasibility screen is conditional on the cached series and these completeness rules, not a quality assessment, reference qualification, or recovery score. The annual counts were recomputed independently from the same cached XML using a separate day-count implementation; the XML parser and source observations were shared.

An initial 15-station USGS intake had already been processed in the national panel, and ten stations had been scored in an earlier external comparison. The subsequent eight-station roster excluded these and other identified source, evaluation and cached-value contacts; no station passed the four-year reference requirement after the documented quality checks. A further five metadata candidates with prior network screening had no

four years of complete approved daily counts in 2010–2024; counts can include estimated values and do not establish subdaily quality. Five selected Réunion Hub'Eau sites provided records with quality code 1, but no site supplied four complete July–June years under the 24-distinct-valid-hour rule; the provider-day time zone is unresolved. The same intake results do not describe the fraction of eligible stations in any complete network.

At ECCC Red River at Emerson, the USGS alias and an upstream source station place the record on a previously encountered river. The files use the word VALIDATED and, for some rows, a station identifier different from the catalog; the mapping to literal V and numeric quality grades and the daylight-saving convention are unresolved. Under the fixed literal V and grade 1/2 rule, zero complete local-clock years qualified. These exclusions are not an assertion that the rejected observations are erroneous; no numerical recovery score was calculated for this site. Data-source citations appear in the main article.

S27. Prior Australian reference exposure

An earlier Australian recruitment stage identified 80 observation-supported candidate station-years from 52 stations. The historical intake record states that its designated F2 formal test was closed; this audit does not run that model or calculate new F2 errors. Subsequently, a separate, earlier-method Australian evaluation used calendar-year cases from the same acquired source archive. All 80 of the candidate station-years appear in that case file; 79 had at least one case in the old evaluation's eligible case set, including all 10 of the 10 originally designated F2 test-role station-years. The prior evaluation contained 30,728 eligible masked cases across its wider roster. These are previously evaluated records for a different pipeline, not unopened target station-years for a new independent test.

A later Australian development analysis used July–June rather than calendar-year windows. Counting physical time overlap rather than matching the year labels alone, 74 of the original 80 calendar station-years intersect a saved development window, of which 73 intersect at least one eligible development case. A separate Arrow/Python recount reproduced these counts. Matching station and interval does not prove that masked days are identical, that the F2 model was evaluated, or that data leaked into source-model training. The original native-sampling criterion allowed 20 observed hours per day and does not establish a complete 24-hour physical reference. These previously opened Australian records cannot supply a fresh independent confirmation of the current pipeline.

S28. Exploratory British Columbia daily-reference comparison

All 23 author-selected published stations were retained for qualification. Four stations each have four complete calendar years in the published daily series. The original study accepted daily means based on at least 20 hourly values after researcher quality checks; hourly counts, original flags and the day convention are not available in this derivative. The source-water cleaning used water–air comparisons. A complete published year is not equivalent to 24 independently quality-checked hours per day. The 23-station author cohort was selected for data availability and these results are unregistered. The fixed global source model and independent source-role recalibration are used; local reconstruction models are refit after holding out the scored year. Each year is masked separately, with 30-, 60- and 90-day controlled gaps. NASA POWER air forcing uses coordinates rounded to 0.01 degree from the source article; there are no same-panel donor stations. The score treats the published daily mean as reference, not as error-free physical truth. Repeated gaps in a station-year and the four stations are dependent; three stations lie in the Fraser River

basin (Nechako, Thompson and Fraser). These are descriptive results, without population uncertainty or a natural-outage validation. Across 372 masked cases, independently

Table S41. Controlled-gap annual H30 mean absolute errors in degrees Celsius. Each row has four published-daily reference years; mask counts are not independent observations.

Station	Masks	Plug-in	D2	Stochastic
08FF003	72	0.833	0.475	0.520
08JA017	72	0.241	0.243	0.203
08LF051	72	0.383	0.207	0.220
08MF040	72	0.605	0.298	0.265

recomputed 30-day reference maxima, observed-window bounds, missing counts, and 52 point and interval summary metrics agree with stored results within numerical tolerance. As a limited noninterference probe, one 90-day case per station was regenerated after adding 5 degrees C to hidden target values. All 4 probes retained the same decision-time model features and numerical predictions, while the evaluation reference changed. These probes do not establish noninterference for every case, model path, or experiment. The attribution, candidate roster, year-level qualification, individual station summaries and code are included in the local evidence package. Prior-identifier and geographic proximity checks are best effort and cannot prove independence from every previously opened record.

S29. Nova Scotia verified workbooks and open-portal comparison

Four river locations were fixed from the provider's historical workbook list before opening their temperature sequences. The workbooks are labelled verified, but their inspected headers do not identify per-hour temperature quality or estimated-value flags; one header states UTC–04:00 and the remaining headers do not establish a time basis. We count

complete reported clock-label years, requiring 24 unique hourly labels with finite numeric cells per calendar day, without claiming these labels correspond to 24 independent physical hours or nonestimated values. The openly licensed portal was accessed only after the workbook completeness screen; it is another release of the same monitoring programme, not an independent sensor. Of 14 station-years with complete labels in both releases, 12 have every numeric value equal within the originally specified 10^{-6} °C tolerance. These include at least four such years at 2 stations. Across 262,667 shared unique numeric labels, 12,810 differ at that tolerance; 1,086 exceed 0.5 °C in a post-hoc diagnostic, with a maximum 4.493 °C. These comparisons do not identify which version is correct, and clock-label matching does not resolve physical time. The original reference screen did not score a reconstruction method; a later, explicitly exploratory amendment is reported separately. The portal source is Government of Nova Scotia, dataset bkfi-mjgw, Open Government Licence – Nova Scotia; legacy workbook reuse remains unresolved.

S30. Nova Scotia reported-clock exploratory masking

The preceding qualification did not establish strict reference quality. After inspecting availability and two-release agreement, a separate unregistered exploratory amendment fixed two sites and eight station-years before any model prediction errors were computed. Each selected year has 24 finite reported hourly labels per day and agrees numerically across the two releases of the same monitoring programme. Time zone, estimated-value status and point-level quality remain unresolved. The reported daily value is the mean of the 24 reported hourly values; it is not guaranteed to represent a physical civil day or an error-free temperature. The fixed global source model and source-only interval calibration were reused. Each year was masked separately with 30-, 60- and 90-day controlled gaps,

using the other three selected years for local history; no empirical-calendar family or whole-archive degradation was evaluated. NASA POWER forcing uses local solar time and coordinates rounded to 0.01 degree. The adjacent-normal tail continuation is an assumption for CRPS. The 144 cases are repeated masks at two sites and eight station-years, not 144 independent watersheds. An independent arithmetic audit recomputed reference H30, bounds, masks and reported scores. In two post-hoc hidden-value probes, adding 5 °C to hidden days changed the true H30 but did not change decision features or predictions when identically serialized controls were compared. Those two probes are not an exhaustive leakage proof.

S31. Nova Scotia reported-day boundary sensitivity

After inspecting the exploratory results, we tested hypothetical daily bins shifted by -1, 0 and $+1$ hour relative to the recorded clock labels. We required all original eight station-years and the adjacent boundary hours to be unique, finite and equal within 10^{-6} °C across the two releases; missing or disputed hours were not filled. Daily water inputs and local leave-one-year-out models were rebuilt for each shift, with the original masks, global source model, source calibration and NASA POWER local-solar-time forcing held fixed. All 144 masks are available across all three shifts for both methods; the zero-shift replay agrees with the original outputs. Across the eight reported reference years the largest H30 change is 0.0023 °C, while a post-hoc casewise diagnostic finds a largest D2 point-prediction change of 0.094 °C. This does not identify the true timezone or explain the cause of prediction changes; hourly values, quality and meteorological day alignment remain uncertain. The two sites and repeated masks do not justify a population-level ranking or independent-climate inference.

Table S42. Retrospective 2009–2017 reported-label completeness and release concordance. Equal means that every hourly numeric value in the common complete-label year matches to 10^{-6} °C at the same reported clock label. Quality, estimated-value status, time basis and climate independence are not established.

Station	Workbook	Portal	Both	Fully equal
NS01DL0009	4	5	4	4
NS01ED9111	2	2	1	1
NS01EF0002	4	4	3	3
NS01FB0001	6	6	6	4

Table S43. Reported-clock exploratory controlled-gap scores on identical available cases. Coverage is percent, and MAE, width and interval score are in °C. These are two sites, not independent climatic replications; source quality and time basis remain unresolved.

Method	Cases	MAE	90% coverage	90% width	Interval score
D2	144	0.311	88.2	1.000	1.572
Stochastic	144	0.313	93.1	1.281	1.706

Table S44. Post-hoc descriptive paired results by gap length and site. Counts are repeated masks; group differences are not inferential tests.

Axis	Level	Cases	D2 MAE	Stochastic MAE	D2 coverage	Stochastic coverage
Gap days	30	48	0.247	0.231	89.6	93.8
Gap days	60	48	0.297	0.319	93.8	95.8
Gap days	90	48	0.387	0.388	81.2	89.6
Station	NS01DL0009	72	0.400	0.393	81.9	93.1
Station	NS01FB0001	72	0.221	0.232	94.4	93.1

S32. Nova Scotia field values and reported hourly ranges

After the portal comparisons and model errors were inspected, we obtained a public field-sample dataset from the same provincial monitoring programme (`https://doi.org/10.25976/sfmj-kf56`). For the same two rivers and 2009–2017 there are 146 water-temperature rows on 73 reported station-dates: each date has two rows with identical metadata and numerical values. We do not know whether they represent duplicate records or separate sampling events. All are labelled preliminary and have no recorded sample time or time zone. Dataset metadata describes co-location with automated probes, but does not establish separate instruments, physical clock alignment or point-level quality.

We compare each field value with the minimum and maximum of the 24 distinct finite portal hours bearing the same reported date, without filling incomplete hours. On 67 station-dates with an available range, values lie inside it on 66. In the previously selected eight model-evaluation station-years, all 33 field dates have a 24-hour reported-label range and 32 lie inside; the largest excess is 0.01 °C. This is not a synchronous comparison: an untimed field value can fall outside the sampled hourly extrema, and an unknown day boundary can change date matching. Pooling the preceding, same and following reported dates gives a much wider, three-day range, not a physical-day uncertainty interval.

After viewing the range result, we also searched all hourly values on each reported date: on 12 of the 33 selected station-dates the field value equals at least one hourly value within 10^{-6} °C. Searching 24 values gives many opportunities for accidental agreement and does not show simultaneous sampling or common/separate sensors. A separate CSV/datetime implementation reproduces all 18 range-summary and 18 proximity-summary cells using the same source files. The field records' preliminary status, absent clock, possible equipment

dependence and provider quality procedures remain unresolved. None of these counts upgrades the two sites to strict independent references, establishes physical sensor accuracy, or validates H30 recovery.

S33. Truckee River reported-clock exploratory evaluation

After the original reference attempts, we accessed the California Department of Water Resources 12-station Truckee water-quality dataset through the public DataONE resolver (`https://doi.org/10.6073/pasta/aefcf302d088aeacc40459091d7e990c`). EML documents provider quality checks and labels the observation clock PST; individual estimated or corrected values are not flagged. Earlier project inventories contain downstream USGS observations from the same watershed. On the named coordinate inventory, the nearest previously known station to each DWR station is more than 10 km away, but coordinate separation does not establish distinct rivers, independent source-model groups or climates. The 12 sites are in one managed watershed.

The station list reports an impossible start after its end for one station; we retained that site for the preplanned timestamp screen instead of interpreting the inconsistency as zero observations. All 12 files have at least four years with a complete grid of unique quarter-hour labels, but this is an upper bound: after requiring all 96 reported water values per day to be finite, only 18 of 171 station-years are complete, and only one station has four such years. The selected years are 2011, 2013, 2014 and 2020 at TRB8. The finite reference consists of each reported-day arithmetic mean of 96 values; it is not a verified nonestimated, physically error-free reference. A second CSV/set implementation recomputed every day and year. The EDI EML declares CC0 while the California discovery catalog lists Creative Commons

Attribution; source attribution is retained and the difference requires reconciliation before permanent release.

After observing availability and numerical reference completeness, but before calculating H30 or target prediction errors, we fixed the one-station cohort and source model. We hold each target year out, use the other three selected years as local history, hide the target gap and use no donor stations. Controlled 30-, 60- and 90-day gaps are crossed with a training-period climatological peak window, with two realizations per cell; there are 72 repeated masks, not 72 independent stations or years. The fixed global source model and source-only calibration are reused. The complete-record reference is subject to sensor and provider-processing uncertainty. No trend is estimated from four years.

After viewing these results, we tested the current NASA POWER hourly UTC and LST products at the same rounded coordinates. For January 1 through December 30 of each reference year, all 34,968 finite LST values equal UTC values eight hours later within 10^{-6} °C; 24-hour LST means differ from the originally retrieved daily product by at most 0.005 °C over 1,457 days. Hourly and daily API versions are v2.10.3 and v2.10.1 respectively. If the provider's PST labels denote fixed UTC-8, the two day boundaries agree; the actual water sensor's daylight handling or other clock shifts remain unverified. These numerical checks do not validate either product's physical accuracy or historical version.

Table S46 summarizes the same 72 available cases. D2 coverage is 16/18 in 2011 and 18/18 in each of the other three years; pooling repeated masks does not establish interval calibration or generality. Independent arithmetic reconstructed all 72 references, bounds and score cells. A single post-hoc 90-day hidden-peak perturbation left model inputs and

predictions unchanged while altering the hidden reference; it is a diagnostic, not exhaustive proof against leakage. No group-bootstrap interval is reported for one watershed.

In a further post-hoc stratification, the complete reference equals the observed bound in 35 masks and exceeds it in 37. In the latter, D2 / calibrated-stochastic MAE is 0.246 / 0.356 °C; their 90% coverage is 36/37 / 37/37 and mean width is 2.046 / 2.740 °C. This stratification uses the hidden complete reference and is unavailable as a decision-time rule. All masks still reuse only four years and one station; the selected comparison and apparent coverage cannot establish population or climate-transfer performance.

After the model results were viewed, a Water Quality Portal search identified a field-measurement record at the same DWR station identifier (`https://www.waterqualitydata.us/provider/STORET/CALWR_WQX/CALWR_WQX-G7166310/`). The published coordinates differ by approximately 66 m; the actual sampling positions are unknown. The query returned 60 accepted, actual field-temperature observations for 2011, 2013 and 2014, and no records for 2020. DWR metadata records an EXO2 instrument start on 13 June 2019; these field records therefore do not examine the 2020 instrument period. All carry PST labels and EPA thermometric method 170.1, whose description permits different thermometer types (`https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=30000Q10.TXT`). Matching each to the nearest scheduled 15-minute logger time gives 59/60 differences within 0.5 °C; the maximum time offset is six minutes. A further post-hoc timing stratification finds a mean absolute difference of 0.185 °C and a field-minus-logger mean difference of +0.179 °C among 39 exactly coincident labels. The records are from the same agency and station, and do not identify whether a separate instrument was used. Field visits are sparse and may coincide with servicing; these instantaneous differences cannot be converted directly

into uncertainty in a 30-day H30 maximum. The comparison assumes both PST labels refer to the same fixed clock and is not an independent-climate test.

The local candidate includes the public DWR source files, provider metadata, reconstructed daily values, a reduced WQP field snapshot and the retrieved POWER responses; offline replays recalculate the timestamp, finite-value, field-comparison and forcing checks. The state catalog describes a DWR discrete dataset as public domain, but a record-level crosswalk to this WQP feed was not independently established. The package does not reproduce model scores without the restricted global source-model weights and calibration. A request for access to those materials or a permitted replacement is still needed.

S34. Additional reservoir-network reference qualification

After the preceding external searches, we examined a 2026 HydroShare collection of continuous water-quality observations above and below reservoirs in Colorado and New Mexico (Van Horn et al., `https://www.hydroshare.org/resource/430ee557554948b5811830c1e2ec41aa/`). One of eleven resources named a previously contacted USGS gage; ten remaining candidate resources were retained for screening, including those with possible network overlap. For the eleven referenced gage proxies, 200-km upstream and downstream NLDI searches found previously named sites for six gages and source-panel sites for two. Gages are proxies rather than necessarily the sensor locations, and a bounded search without a hit does not establish independence.

The ten published level-1 files contained 69 station-years. Reported fixed Mountain Standard Time labels allowed 29 potentially complete calendar years before checking numerical values. Requiring exactly 96 unique quarter-hour values per day, all finite and in the broad -5 to 50 °C sensor-range guard, yielded 16,153 complete days but no complete

calendar year. This is a screen of these ten files under this stringent rule, not a statement about all years in the network. The published level-1 process can correct measurements, while explicit per-value correction or estimation flags were not found; finite values do not establish independent, uncorrected ground truth. No annual H30, model error, or trend was scored. The candidate package includes attributed time-and-temperature column extracts, station-year counts, and an independent replay. USGS series metadata for four of eleven gages show some water-temperature records, including instantaneous records extending into the study period for two Arkansas gages, but contemporaneous coverage and co-location remain unverified and both gages have known source-panel network connections.

After inspecting the completeness results, we bounded H30 in every 2021–2025 station-year, assuming each unobserved or ambiguous quarter-hour lies between -5 and $50\,^{\circ}\mathrm{C}$ and treating finite, uniquely reported values as fixed. The range is an assumption about unobserved water, not an established physical or sensor-error bound. Each daily bound is the mean after setting all unknown slots to the relevant endpoint; the annual bound is the maximum of the resulting 30-day moving averages wholly inside the calendar year. The construction encloses the reported-series target only if the assumptions hold. Four of 50 intervals have width at most 0.01 °C and 13 at most 0.1 °C, whereas the median width is 25.18 °C. The ten sites are not independent climates and the narrow intervals were selected after the availability study; no H30 prediction was scored.

All 50 years contain a fully observed 30-day window. A generic 95% complete-day rule accepts 11 years and rejects 39 with visible windows; the 100% rule accepts none. Among rejected years with at least one previously accepted year during 2021–2023, one has a visible window exceeding every previously accepted annual upper bound. At Rio Chama

below Abiquiu, 2024 has 346/366 complete days, and its visible 30-day mean is 19.386 °C. The only earlier accepted year in 2021–2023 is 2022, with 357/365 complete days and conditional H30 interval [18.461, 18.461] °C; the 2024 window is 0.925 °C higher. The 2024 conditional upper bound is 28.190 °C, and rejected 2021 or 2023 may also have higher H30. This is not a record against all earlier years, a physical measurement-error guarantee, or a prevalence estimate. Source quality corrections and hydrologic dependence remain unresolved. An independent implementation recomputes all 50 bounds and both screening rules; the candidate package replays them from attributed source extracts.

S34.1. Partial-history exploratory model comparison

The original model pipeline requires at least four complete daily training years, which none of these sites supplies. After examining completeness and H30 bounds, we fixed an exploratory adaptation using all candidate target years with at least 30 complete days and at least three other history years satisfying 75% complete days, observed endpoints, and no missing run longer than 45 days. Missing days in each selected history year were linearly interpolated within that year; up to 58 days in one history year were interpolated across these cases. The target year was excluded from fitting, and its naturally incomplete days and additional controlled 30/60/90-day masks were hidden together. We supplied the actual count of complete training years, zero, to the fixed global source model and source-only calibration. This is outside the original complete-history regime, not a strict replay of it. NASA POWER local-solar-time air forcing may not align with the reported fixed-MST water days; no neighboring water donor was used. Prior network exposure is known for some gage proxies, while no 200-km hit for others does not prove independence.

All 306 masks in 17 target station-years at four sites produced all three point estimates. For a point p and a conditional reference interval $[L, U]$, the absolute error is bounded by $\max(L - p, 0, p - U)$ and $\max(|p - L|, |p - U|)$. Predictive coverage is definite only if the interval contains $[L, U]$, and possible when the intervals intersect. On the post-hoc subset of 180 masks in ten station-years at three sites with reference width at most 0.1 °C, mean absolute error is bounded by [0.299, 0.325] for plug-in, [0.348, 0.374] for D2 and [0.318, 0.343] °C for the calibrated stochastic comparator. The paired D2-minus-plug-in mean absolute-error difference is bounded by [0.045, 0.052] °C; the D2-minus-stochastic difference is [0.026, 0.033] °C. These average per-case extrema are conservative because masks within a station-year share an unknown H30. D2 nominal 90% coverage is bounded between 80.0% and 88.9%, and stochastic coverage between 83.9% and 95.6%, on that same selected set. The full 306-case cohort has mean absolute-error upper bounds exceeding 11 °C for all methods, so its rankings remain weakly identified. These are repeated masks, not independent sites, and no general ranking or coverage guarantee follows.

An independent scalar implementation recomputed all bounded scores. In one selected mask per target year, paired same-process perturbations of all hidden observed daily values and of natural-gap placeholders preserved all 68 model features and 17 outputs, within 10^{-10} tolerance. An earlier comparison against saved outputs failed in four of 17 masks under the audit's default numerical-thread environment, with maximum drift 0.0061 °C; setting OMP, OpenBLAS, MKL and NumExpr threads to one before numerical imports yielded agreement in those 17 checks. An additional single-thread replay reproduced all 306 saved masks, features and predictions using the same code and source model. This does not

prove that every mask is free of leakage, quantify physical uncertainty, or independently refit the restricted model.

S34.2. Training-feature support and the numerical gate

After the score analysis, we reconstructed the global model's seed-zero role assignment and bootstrap weights. Its quantile regressors fitted 43,804 distinct positive-weight source cases from 68 recorded groups, with total bootstrap weight 68,028. One-dimensional finite training minima and maxima were compared with all 68 decision-time input features. All 306 reservoir cases have zero complete training years, below the source estimator range of 3–41. Excluding this feature, every case is still outside a training extremum: 270 have a training-anomaly lag-one correlation below the estimator minimum and the remaining 36 have a reconstruction-residual lag-one correlation above its maximum. These are repeated masks in 17 station-years. For comparison, 1,931 of 30,384 crossed-mask cases in positive-weight non-estimator source roles are outside a marginal training range excluding complete-year count; those source cases were used in other model components and are not a new independent test set. Ranges are marginal, not a multivariate support guarantee, and this comparison does not establish a failure probability.

For the 17 reservoir station-years, the anomaly correlation calculated from interpolated historical days is outside the source range in 15 years. Rebuilding the historical climatology using only fully observed days, while retaining the interpolated version's warm months, leaves 14 of those 17 correlations outside the source range. This changes both the climatology and finite lag pairs, so it does not isolate a causal interpolation effect; it also does not recompute the residual correlation or prediction accuracy. An independent day-of-year and scalar correlation implementation reproduced these results.

For a separate diagnostic, we counterfactually clipped complete-year count, anomaly correlation, and residual correlation to their respective source training extrema, individually and jointly. All 306 cases retained the same leaf IDs in all seven quantile regressors and the classifier, and hence the same raw quantile outputs and classifier scores. Sorting, source offset and observed-window projection were independently reproduced. This does not mean the features have no influence inside the training range, and clipping an actually incomplete history is not a valid fix. The original three-output rule additionally requires at least four complete history years, so it cannot release a numerical estimate for any of these cases regardless of the individual score. If quality flags were all absent and optional limits were omitted, 304 cases would produce an observed bound and two would be unknown; actual source quality flags are unresolved, so those are conditional states, not actual releases. An individual score at or below 0.1 occurred in 158 cases, but it is not a group-risk guarantee.

S34.3. Source evaluation of a marginal-range warning

After inspecting external cases and source outcomes, we examined whether the marginal-range warning identifies errors in crossed-mask cases from the global model's positive-weight rule-calibration role. That role is group-disjoint from the quantile estimator, interval calibration and risk learning roles, but was used in other rule analyses. We retained 8,010 cases from 19 groups with finite D2, reference and individual score, a finite observed bound and at least four complete historical years; 648 of 8,658 source role cases did not meet these criteria. Quality flags and the original optional range gates were not applied, so this is not the complete three-output policy. An error is an absolute D2 error above 0.5 °C against the reported complete reference.

The warning flags an input outside any estimator-role marginal minimum/maximum, excluding complete-year count, or a nonfinite category absent in training. It flags 484 cases, including 42 of 868 errors: 4.8% recall and 6.2% false-positive rate among non-errors. Score ≤ 0.1 alone retains 6,007 cases with 288 errors. Requiring both score ≤ 0.1 and no warning retains 5,596 with 268 errors. To compare equal numerical retention, an outcome-free but transductive ranking of the same cases by score retains 5,596 with 211 errors. The case-weighted difference in conditional failure is +1.02 percentage points; the 19-group bootstrap interval is [+0.68, +1.41] points. Selected years can differ, and the score-only cutoff is selected from this evaluated cohort's score distribution, so neither the interval nor the matched comparison is an external safety guarantee.

As a post-hoc numerical-tolerance sensitivity, we enlarged each finite training range by 0.000001 or 0.01 of that feature's weighted 1st–99th percentile span, falling back to its full range or one for zero span. The smaller margin leaves the counts unchanged. At the 0.01 margin, the warning flags 430 source cases and 38 errors (4.4% recall); the same-count score-plus-warning and score-only policies retain 268 and 216 errors in 5,620 cases, respectively. It flags 270 of the 306 reservoir masks, rather than all 306. Margins were chosen after earlier results, source cases are repeated masks, and one-dimensional ranges do not establish multivariate support. Independent Booster, decision, group and bootstrap implementations reproduced both comparisons.

S34.4. Source-history degradation and count-feature sensitivity

After inspecting the reservoir results, we selected all 15 Swiss and Bavarian stations from the earlier processed-data cohort with complete daily years 2009–2017: 11 Swiss and four Bavarian stations, representing six recorded connected groups. The target is

the reported 2017 calendar-year H30. Its nine fixed target masks cross 30/60/90-day lengths with three calendar positions and remain identical across all interventions. In the full-history condition, 2009–2016 daily observations are retained. In the degraded-history condition, July 1–30 is hidden in each of those eight training years and linearly interpolated within that year before fitting the local daily model. The fixed source annual models and calibration exclude target-region groups; no water donor is used.

The D2 complete-history-count feature is crossed with history condition at values 0 and 8. A count of 8 is truthful only for the full history; zero is truthful only for the degraded history. The two other combinations deliberately misstate history completeness and are diagnostic counterfactuals, not deployment rules. In the 135 cases common to all four D2 combinations and both plug-in and stochastic history versions, D2 MAE is 0.158 °C for full/count 8 and 0.188 °C for degraded/count 0. Averaging over the two count settings gives a case-equal history effect of +0.027 °C; averaging over histories gives a count-8-minus-count-0 effect of −0.002 °C. The corresponding six-group-equal estimates are +0.042 °C and approximately zero, with 4,000 group-bootstrap intervals [−0.009, 0.092] and [−0.0015, 0.0014] °C. D2 nominal 90% coverage changes from 91.1% for full/count 8 to 88.9% for degraded/count 0, while mean interval score changes from 0.787 to 0.963 °C. Repeated masks and clustered stations do not provide 135 independent tests.

The selected years, data vintage and source results were previously examined; a single fixed warm-season degradation and one target year do not characterize real missingness. They also cannot isolate why the reservoir adaptation differed from its comparators. Independent arithmetic recomputed the 15 target H30 values, all 540 factorial rows and six-group summaries; one post-hoc perturbation of 30 hidden target days and 240 hidden

historical days left the corresponding model inputs and predictions unchanged. A single-thread replay in a separately provisioned Python environment reproduced all 540 saved masks, features and predictions using the same code and local restricted models. The open snapshot replay checks source water-year completeness and air inputs, but the source model and case predictions remain subject to separate access conditions.

S35. Previously opened USGS daily targets with continuous support

After the preceding studies, we reexamined all ten previously opened USGS daily-value candidates (`https://api.waterdata.usgs.gov/`). Eight complete calendar years at four sites had finite approved daily means, no ambiguity, and no estimated qualifier recognized by the existing parser; every target had at least three other years with at least 75% usable days, finite endpoints and no missing run longer than 45 days. Missing historical daily values were linearly interpolated within each history year, while the actual complete history count was zero to two. Target years were excluded from local fitting, but later history was allowed. These inputs fall outside the global model's estimator training range of three to 41 complete history years. The global source model and source-only calibration were held fixed, without a target water donor; this is not the original four-complete-history protocol or a prospective forecast.

A post-hoc audit checked all 2,922 raw published daily records in the eight target years: each was approved with an empty qualifier and matched the processed target value. Empty qualifiers do not prove absence of unrecorded corrections or physical measurement error. All four daily series have a uniquely matched continuous parent ID in the USGS metadata. We obtained the continuous parent observations for all eight target years and required finite, approved, unique, quarter-hour-grid readings with no nonempty qualifier. Under

fixed local standard time, three years have at least one accepted reading in each of the 24 physical hours of every day, and one of those years has all 15-minute slots; all three years are at Rio Mora near Terrero, New Mexico (USGS 08377900). Under local civil time, daylight-saving transition days have 23 or 25 physical hours; an independent scalar audit verified 5,844 day-by-clock records across both conventions. One of the four target station IDs is present in a source panel; this alone does not prove positive-weight source-model use. A bounded 200-km network search found no named source or other prior station for Rio Mora, but a non-hit does not prove an independent watershed or climate. Daily records and some target H30 outcomes had already been examined; selection is post-hoc.

The reported daily H30 in Rio Mora in 2017 is 14.1100 °C. Recomputing from the fully supported 15-minute series gives 14.1066 °C using fixed standard time and 14.1055 °C using civil time. Close annual values do not establish the actual reporting clock, physical sensor accuracy, unrecorded corrections, or USGS daily preprocessing. The station's earlier quality report also flagged possible ice effects and long constant runs. NASA POWER air forcing uses local solar time and need not align with either water-day convention.

All 144 artificial masks have finite plug-in, D2 and calibrated stochastic point estimates. Against the reported complete daily reference, their MAEs are 0.304, 0.251 and 0.300 °C, respectively; nominal 90% coverage is 121/144 for D2 and 123/144 for the stochastic method. In the post-hoc fixed-standard hourly-supported subset, all 54 masks are drawn from three years at the single Rio Mora station: MAEs are 0.266, 0.180 and 0.174 °C, with nominal interval coverage 52/54 and 54/54. Repeated masks do not make these independent stations or climates; no regional accuracy, trend or risk guarantee follows.

A subsequent audit found that the existing daily parser recognized estimated values but retained other nonempty qualifiers. In Rio Mora, 2015 and 2016 contain 57 and 88 raw daily ICE qualifiers, respectively; ICE does not itself prove the values incorrect. Conservatively removing every nonempty daily qualifier and reapplying the same history eligibility rule removes both years from each of the three Rio Mora target histories. Across all eight target years, the same 144 original masks remain supported; five targets have unchanged histories. On the Rio Mora 54 masks the plug-in, D2 and stochastic MAEs change from 0.266, 0.180 and 0.174 to 0.242, 0.226 and 0.218 °C, respectively. This post-hoc comparison simultaneously changes input values and historical sample size. A scalar audit of raw qualifiers and annual eligibility passed. A fixed-mask hidden-value perturbation on one mask in each of the eight target years and an isolated single-thread replay of all 144 qualifier-aware cases matched saved inputs and outputs; neither is an exhaustive leakage proof.

The saved reported-daily H30 and an independently summed version differ by at most 7.11×10^{-15} °C, yet intervals often touch an observed lower bound, so exact floating-point comparisons change up to 27 of 144 coverage classifications across these arithmetic paths. A post-hoc absolute 10^{-10} °C comparison tolerance yields identical classifications for the two paths. For Rio Mora, original-history D2 and stochastic reported-daily coverages are 53/54 and 54/54 with this tolerance, versus 52/54 and 54/54 using the saved exact comparison. With qualifier-aware histories, the tolerant coverages are 48/54 and 53/54. This numerical convention is not a measurement-error model. An independent scalar audit verified all 24 coverage summary cells.

The 144 cases have zero, one or two genuinely complete training histories. All are below the estimator-role minimum of three, although the other 67 inputs lie within their individual source extrema; marginal inclusion does not establish joint support. Replacing the history count by the training minimum is knowingly false and leaves every quantile-model and classifier leaf path unchanged, so it is not a correction. The original numerical-release gate requires four complete history years and therefore cannot release numeric values for these cases. An independent scoring audit checked all 144 saved outputs against reported daily references; eight same-process hidden-value probes and an isolated single-thread replay of all masks and model inputs passed. The open USGS/NASA snapshot independently replays reference arithmetic and clock qualification, but omits the restricted source weights and per-case predictions.

S35.1. Reporting resolution and source-recalibrated support margins

We subsequently bounded the annual 15-minute-sample H30 in all eight previously opened USGS years under both fixed-standard and civil reporting clocks. Each missing, flagged, duplicated or otherwise ambiguous quarter-hour value is assumed to lie in $[-5, 50]\,^{\circ}\mathrm{C}$; an off-grid day is treated as entirely unknown. For each reporting day, lower and upper means assign all unknown slots to the corresponding endpoint; each annual bound is the maximum of its 30-day rolling sequence. This hypothetical range is not a validated instrument or physical-water limit. Under fixed standard time, three years have zero-width H30 bounds, all at Rio Mora: 2012 and 2019 each retain a small number of unknown quarter-hours despite full hourly coverage, while 2017 has every quarter-hour slot. Independent UTC-slot and scalar-window code reproduced all 5,844 day-by-clock bounds.

On the 54 masks in those three years, nominal 90% D2 coverage is 52/54 against published daily H30 but 33/54 against the fixed-standard continuous H30; stochastic coverage is 54/54 and 34/54. In 20 masks the largest observed published-day window exceeds the conditional continuous H30 upper bound; the D2 and stochastic 5th percentiles equal the published bound in all 20. The discrepancy has median 0.0034 and maximum 0.0118 °C. All 21 D2 non-overlap distances are at most 0.05 °C, with maximum 0.0203 °C. This threshold was examined after the scores and is a descriptive resolution sensitivity, not a sensor-error estimate or calibrated coverage. Published daily and equally weighted quarter-hour targets can differ through clock, rounding or corrections; an apparent cross-target contradiction is not itself proof of model leakage or physical failure.

To examine the consequence of a measurement-error assumption, we applied support $L-\epsilon$ instead of the original published-window bound L before and after quantile calibration, for both D2 and stochastic methods. For $\epsilon = 0$, 0.01, 0.05 and 0.20 °C, the source offsets were recomputed using the same positive-weight source interval-calibration roles: 13,007 cases from 13 recorded groups. Zero margin reproduces the previous source offsets and target quantiles. New target outcomes were not used in the recalculation, though prior source-panel contact is unresolved. The source calibration reference remains reported daily H30, so its validity for a latent continuous target is not established.

On the same 54 masks, all three positive margins yield 37/54 D2 and 34/54 stochastic continuous-target coverage; against the published daily target they yield 46/54 and 44/54. Under $\epsilon = 0.05$, ten D2 lower interval endpoints and four medians fall below the original published observation bound. Thus a softened support changes the estimand or requires a justified measurement-error model; it cannot silently replace the exact reported-series

constraint. Of 2,828 fixed-standard days with all quarter-hour samples, 264 have an absolute daily-mean discrepancy greater than 0.05 °C; none exceeds 0.20 °C in this observed subset. This does not bound missing days, validate either clock or establish instrument accuracy. A separate scalar implementation reproduced 868 source offset values and all 32 margin/clock/cohort/method summary cells. The experiment is post-hoc, uses one station for the three-year subset and does not establish reliable calibration or deployment safety.

Table S45. Paired exploratory reported-day boundary sensitivity with identical original masks. Coverage is percent and error, width and interval score are in °C. Local stages are refit; source model, source calibration and forcing are held fixed. Two sites and eight station-years are not independent climatic replications.

Shift (h)	Method	Cases	MAE	90% coverage	Width	Interval score
-1	D2	144	0.310	88.9	1.000	1.555
-1	Stochastic	144	0.313	92.4	1.283	1.704
0	D2	144	0.311	88.2	1.000	1.572
0	Stochastic	144	0.313	93.1	1.281	1.706
1	D2	144	0.315	86.8	0.998	1.572
1	Stochastic	144	0.313	93.1	1.278	1.731

Table S46. Truckee exploratory annual H30 scores on 72 controlled masks from one station and four years. MAE, bias, width and interval score are °C; coverage is percent for the nominal 90% interval. The plug-in method supplies no interval.

Method	Available	MAE	Bias	Coverage	Width	Score
Plug-in	72	0.458	-0.458	–	–	–
D2	72	0.131	-0.047	97.2	1.106	1.133
Calibrated stochastic	72	0.188	-0.100	98.6	1.459	1.464

S36. Exploratory process and numerical diagnostics

These checks were developed after inspecting earlier regional and time-support results. They are exploratory and reuse the opened archive; selection of six stations before a numerical replay is not preregistration of the research questions.

S36.1. Common support and common-weight standardization

Common geometric cells contain gap length, training-peak overlap, year and visible-donor bins. Screening to shared cells preserves region-specific cell frequencies and is not balancing. The primary standardized target assigns identical weights proportional to the smaller regional cell count; equal-cell weighting is a sensitivity. Joint-cell frequency TV becomes zero under either common target, but within-bin continuous quantities and training histories remain imbalanced. In the geometry min-count analysis, mean training complete-year counts are 9.78 in FOEN versus 22.96 in Bavaria (SMD -3.052); after adding reconstructor they are 8.50 versus 20.18 (SMD -2.707). Saved CSV files also report continuous means, weighted KS distances and missing-weight fractions. A coverage contrast remains under these common geometric-cell weights; residual within-cell and history differences remain, and the comparison does not identify a reconstructor effect or exclude river processes. The absence of a Swiss air-ridge stratum prevents corresponding cross-region comparison of the main Bavarian failure layer.

S36.2. Annual water–air associations

Pooled annual associations mix station differences and within-station variation. We therefore center both H30 quantities within station and period, before/from 2010, and pool those centered observations. Both periods of a sampled connected group remain paired in

Table S47. Descriptive regional coverage standardization on common cells. Min-count weights are the primary common target; equal-cell weights are a sensitivity analysis. Coverage is in percent and differences are percentage points. Geometry cells contain gap, training-peak overlap, year and visible-donor bins. Adding the reconstructor changes the supported population; it does not identify a reconstructor effect.

Cells	Weights	FOEN	Bavaria	Difference
Geometry	Original	98.75	82.78	15.97
Geometry	Min-count	98.72	82.68	16.04
Geometry	Equal-cell	99.29	81.68	17.62
+ reconstructor	Original	99.58	94.12	5.46
+ reconstructor	Min-count	99.72	96.04	3.69
+ reconstructor	Equal-cell	99.75	93.41	6.33

Reconstructor common support retains 472/1,119 FOEN and 289/1,983 Bavaria cases. Switzerland has no air-ridge cases, so Bavaria's air-reconstruction failure stratum has no cross-region counterpart.

10,000 bootstrap replicates (seed 20261029); station-paired resampling is an additional sensitivity, not an assumption-free replacement. FOEN has two groups, with only three distinct bootstrap group compositions; its intervals are particularly coarse. Similar annual within-slope point estimates neither establish equivalence nor validate gap-day residuals, heat-peak timing, tails or autocorrelation. Each variable is its own annual maximum and their maximizing dates need not coincide.

Table S48. Remaining balance after regional standardization. TV denotes total variation of discrete distributions. History denotes training complete-year count. SMD denotes the FOEN-minus-Bavaria mean difference divided by the pooled population standard deviation in the original two regions, kept fixed across weighting schemes. Full continuous means, missing-weight fractions and weighted Kolmogorov distances are saved with the analysis. Zero joint-cell TV does not ensure balance within bins.

Cells	Weights	Joint TV	Policy TV	History-bin TV	History SMD
Geometry	Full	0.232	0.854	0.948	-3.006
Geometry	Common raw	0.228	0.853	0.948	-3.000
Geometry	Min-count	0.000	0.919	0.979	-3.052
Geometry	Equal-cell	0.000	0.780	0.916	-3.176
+ reconstructor	Full	0.884	0.854	0.948	-3.006
+ reconstructor	Common raw	0.614	0.000	0.696	-2.292
+ reconstructor	Min-count	0.000	0.000	0.722	-2.707
+ reconstructor	Equal-cell	0.000	0.000	0.760	-2.891

S36.3. Year-set comparisons and retained baselines

The original single random selection and incomplete-year-only spacing rule are retained as labeled legacy comparators in the table below. The exact subset expectation and the two selectors that incorporate fixed complete years are the current exploratory extensions. Paired intervals in this table use the common updated resampling specification; the previous resampling tables remain archived with the earlier revision.

Table S49. Hidden-reference-peak recovery by region and daily reconstructor. The hidden-peak label uses withheld complete-year H30. Coverage is nominal 90%. Above-interval is the share of complete references hotter than the 95th percentile.

Region	Reconstructor	Method	Cases	MAE	Bias	90% coverage	Above interval
Switzerland	donor_ridge	D2	633	0.175	-0.130	98.1%	0.9%
Switzerland	donor_ridge	classical	633	0.180	-0.133	96.7%	2.7%
Bavaria	air_ridge	D2	995	0.574	-0.479	67.9%	30.2%
Bavaria	air_ridge	classical	995	0.486	-0.296	66.5%	27.1%
Bavaria	donor_ridge	D2	163	0.249	-0.129	90.8%	8.6%
Bavaria	donor_ridge	classical	163	0.205	-0.079	87.7%	8.6%

S36.4. Common gap metric and archive count audit

The primary comparison uses 2010–2025 to match the forward experiment. The observed eligible archive covers 41 forward-cohort stations; only 29 have incomplete years in this period. There are 109 unique incomplete station-years, without artificial mask repetition. The empirical-calendar and controlled-gap summaries contain 3,806 and 3,777 incomplete station-year–replicate rows, respectively, across the same 41-station forward cohort. Their repeated appearances preserve experimental weighting and do not create new independent observations. All three summaries use each year's longest contiguous missing run, including unions of adjacent or overlapping missing spans.

As an additional historical description, the observed archive over 1981–2025 contains 220 incomplete station-years at 31 stations; the median annual longest run is 3 days and 11.36% have a run of at least 30 days. This historical period is supplementary rather than the primary comparison to forward masks. The calendar masks are closer to common

Table S50. Annual water–air H30 associations in unique complete station-years. Years and slopes are shown before/from 2010; S/G counts stations/connected groups. Within estimates center both variables within station and period. Slopes and correlations pool the centered annual observations, rather than averaging station coefficients. The paired 95% percentile intervals resample the same connected groups in both periods; only two FOEN groups and eight Bavaria groups are available, so these intervals are descriptive and coarse. Station-resampling sensitivity results are saved separately.

Region	Fit	Years	S/G	Slopes	Δb [95% interval]
FOEN	Pooled	98/159	10/2	1.143/1.176	+0.033 [-0.005, +0.047]
FOEN	Within	98/159	10/2	0.715/0.701	-0.014 [-0.015, -0.005]
Bavaria	Pooled	554/295	24/8	1.318/1.409	+0.091 [-0.151, +0.346]
Bavaria	Within	554/295	24/8	0.814/0.809	-0.005 [-0.224, +0.149]

The air and water H30 maxima need not refer to the same dates. Similar annual within slopes do not establish stable daily residuals, hidden-peak timing, tails or autocorrelation, and are not an equivalence test.

observed gap conditions on this metric, whereas 30–90-day controlled masks probe the long-gap tail; differing sampling and repeat weights preclude a same-distribution claim.

The shared-year fraction counts station-years in which at least one other station in the same connected group also has an incomplete year (within the same replicate for trial masks). It does not require overlapping missing days and therefore does not establish synchronous daily outages or unavailable donors on the same dates.

Complete-year counts in the cohort comparison count unique complete station-years in the leave-one-year-out source cases, including the held-out year. The previous 30/7

Table S51. Annual water–air correlations and paired changes. Intervals use connected-group resampling, as in Table S50. Correlations describe annual H30 associations at the stated aggregation scale.

Region	Fit	$r_{<2010}$	$r_{\geq 2010}$	Δr [95% interval]
FOEN	Pooled	0.842	0.817	-0.025 [-0.129, -0.014]
FOEN	Within	0.855	0.725	-0.130 [-0.130, -0.129]
Bavaria	Pooled	0.649	0.708	+0.059 [-0.027, +0.100]
Bavaria	Within	0.826	0.800	-0.025 [-0.066, +0.051]

medians counted training years: every one of the 581 stations differs by exactly one year. Correct archive medians are 31/8. Visible-donor summaries derive from leave-one-year-out cases and describe that design, rather than the forward controlled-gap experiment.

S36.5. Earlier uncalibrated-only trajectory diagnostic

The original three-seed check reports raw, uncalibrated endpoints in the fixed 27-case sample. Its small pooled median does not establish stratified stability. The subsequent experiment section additionally regenerates the original source-calibration roles and re-estimates offsets at each budget; the table below is retained specifically as the uncalibrated reference.

Table S52. Paired resampling sensitivity for within-station annual water–air H30 changes (from 2010 minus before 2010). Each resample carries both periods of a selected group or station. Connected groups are the primary sampling unit; station resampling assumes independence across stations. Neither percentile interval establishes equivalence, and FOEN group intervals have only three distinct possible compositions.

Region	Unit	n	Δb [95% interval]	Δr [95% interval]
FOEN	Group	2	-0.014 [-0.015, -0.005]	-0.130 [-0.130, -0.129]
FOEN	Station	10	-0.014 [-0.043, +0.032]	-0.130 [-0.163, -0.097]
Bavaria	Group	8	-0.005 [-0.224, +0.149]	-0.025 [-0.066, +0.051]
Bavaria	Station	24	-0.005 [-0.123, +0.109]	-0.025 [-0.069, +0.022]

S37. Additional exploratory experiments

S37.1. Outcome-blind year-set selection

All feasible same-count subsets are enumerated while retaining every fixed complete year. There are 5758 subsets across 807 common archives; the 805 actual-mask primary archives contribute 5756 subsets. The two zero-mask settings contribute one subset each. Only 468 actual-mask archives offer a nontrivial choice; the remaining 337 have identical same-count selections by construction. CDF minimizes the retained-versus-full empirical-year-CDF squared distance, including complete years; Sxx maximizes the centered decade-year sum of squares of the entire retained set. Ties use the alternate geometry score and then the earliest incomplete-year tuple. No temperature or error enters either selector. They change 367 and 376 selected sets relative to warm-season screening, respectively. These

Table S53. Exploratory finite-record slope comparisons on 805 actual-mask archives in 17 connected groups. Errors are °C per decade. CDF and Sxx optimize the entire retained set including fixed complete years. Uniform exact averages squared errors over feasible same-count sets within each archive, then averages archives and takes the square root. All-year truth-set error is zero by definition. The last column gives comparator-minus-warm-season estimated RMSE with a paired 95% group-percentile interval, not an equivalence test.

Rule	Years	Estimate	Year-set	Reconst.	Δ RMSE [95% interval]
Warm-season	9.59	0.4459	0.4458	0.0023	Reference
Legacy random	9.59	0.4540	0.4569	0.1030	0.0081 [-0.0103, 0.0177]
Legacy spacing	9.59	0.4566	0.4539	0.0873	0.0108 [-0.0063, 0.0404]
Uniform exact	9.59	0.4492	0.4483	0.0978	0.0033 [-0.0052, 0.0171]
CDF	9.59	0.4284	0.4199	0.1079	-0.0175 [-0.0274, -0.0049]
Sxx	9.59	0.4023	0.3931	0.1185	-0.0436 [-0.0645, -0.0173]
All years	13.32	0.1697	0.0000	0.1697	-0.2762 [-0.3557, -0.1841]

Table S54. Common annual-longest-gap comparison in 2010–2025. Each number describes the longest contiguous missing run within an incomplete year. Observations weight incomplete station-years equally; trial rows weight incomplete station-year–replicates equally, retaining all 20 mask replicates. Repetition and station composition differ, so this is descriptive rather than a same-distribution test.

Source	Rows	Stations	Median (d)	90th percentile (d)	≥ 30 d
Observed archive	109	29	4	33.6	11.93%
Empirical calendar	3,806	41	6	30.0	10.25%
Controlled gaps	3,777	41	60	90.0	100.00%

Table S55. Long-complete evaluation stations versus the remainder of the 581-station source archive. Complete-year counts are unique complete station-years, including the held-out year, rather than training-year counts. USGS counts follow named hydrologic-unit regions.

Panel	Stations	Groups	Median years	Bavaria	Switzerland	USGS
Trend cohort	42	18	31	24	10	7
Source remainder	539	183	8	29	1	312

Table S56. Forward-mask geometry on the 41-station trend cohort with twenty replicates. Year selection after a seven-year burn-in is shared. Segment length and season come from the copied empirical calendars or from the controlled-gap design. Shared-year fraction is the share of masked station-years that are also masked at another station in the same connected group.

Family	Median length (d)	JJA share	Adjacent years	Shared years	Stations
Empirical calendar	4	19.9%	43.9%	60.9%	41
Controlled 30–90 d	60	58.5%	43.5%	60.1%	41

post-review exploratory rules are not deployment recommendations and do not separately identify effects of year count and temporal position.

Uniform exact means $\sqrt{\text{mean}_a\, E_{U|a}(E_a^2)}$, with an equal-probability set U inside each archive a. It is not the mean RMSE over random selections. Over 200 random seeds, the mean RMSE is 0.4505, the root of mean sampled MSE is 0.4506, and the exact conditional root-expected-MSE is 0.4492 °C per decade. The across-seed 2.5th–97.5th percentiles [0.4359, 0.4671] describe selection variation, not a confidence interval; 62/200 seeds improve on the warm-season point result. Paired intervals in Table S53 instead

Table S57. Bavarian whole-archive stations. Regulation labels are project metadata.

Code	Group	Latitude	Longitude	Years	Regulation
10026293	g0032	48.389	9.987	1994–2024	unmatched_gages
10039802	g0032	48.711	10.802	1997–2025	unmatched_gages
10046105	g0032	48.757	11.425	1981–2025	unmatched_gages
10056302	g0032	48.947	12.015	1993–2025	unmatched_gages
10074009	g0032	48.886	12.574	1981–2024	unmatched_gages
10081004	g0032	48.826	12.961	1981–2024	unmatched_gages
10089006	g0032	48.635	13.187	1981–2025	unmatched_gages
10091008	g0032	48.576	13.459	1981–2025	unmatched_gages
10098007	g0032	48.514	13.727	1981–2023	unmatched_gages
11401009	g0034	47.524	10.260	1981–2025	unmatched_gages
11402001	g0034	47.730	10.317	2001–2025	unmatched_gages
11405000	g0034	48.371	9.988	1981–2025	unmatched_gages
16003003	g0036	47.684	11.572	1981–2025	unmatched_gages
16005701	g0036	48.145	11.597	1981–2024	unmatched_gages
16007004	g0036	48.528	12.133	1981–2025	unmatched_gages
17207508	g0039	48.611	13.145	1981–2025	unmatched_gages
18001508	g0035	47.855	12.143	1981–2020	unmatched_gages
18008008	g0035	48.562	13.444	1981–2024	unmatched_gages
18403002	g0031	47.934	12.481	1981–2024	unmatched_gages
24010004	g0037	49.951	10.874	1981–2025	unmatched_gages
24022003	g0037	50.031	10.221	1981–2025	unmatched_gages
24042000	g0037	49.796	9.926	1981–2025	unmatched_gages
24064003	g0037	49.716	9.225	1981–2025	unmatched_gages
24405002	g0033	50.197	10.075	1981–2025	unmatched_gages

Table S58. Non-Bavarian whole-archive stations, including Switzerland (FOEN) and USGS HUC17. Regulation labels are project metadata.

Code	Group	Latitude	Longitude	Years	Regulation
2016	g0029	47.482	8.194	2000–2025	unmatched_gages
2019	g0029	46.746	8.092	2000–2025	unmatched_gages
2029	g0029	47.122	7.283	2000–2025	unmatched_gages
2030	g0029	46.765	7.612	2000–2025	unmatched_gages
2085	g0029	47.056	7.184	2000–2025	unmatched_gages
2113	g0029	47.595	8.225	2000–2025	unmatched_gages
2126	g0030	47.496	8.953	2002–2025	unmatched_gages
2135	g0029	46.933	7.448	2000–2025	unmatched_gages
2205	g0029	47.516	8.235	2000–2025	unmatched_gages
2457	g0029	46.694	7.880	2000–2025	unmatched_gages
1670	g0087	62.598	25.739	1935–1994	unmatched_gages
02172020	g0181	33.094	-79.949	1988–2025	unknown
04137030	g0166	44.473	-83.571	1999–2024	unknown
06036905	g0053	44.620	-110.864	2003–2025	regulated
06054500	g0143	46.147	-111.420	2001–2025	regulated
13342500	g0054	46.448	-116.828	2001–2024	regulated
14166000	g0057	44.270	-123.174	2003–2025	regulated
14211500	g0059	45.477	-122.508	2000–2021	regulated

resample the recorded connected groups 10,000 times, preserving paired archives and their conditional subset-average MSEs before taking roots.

For each fixed set $E = S + R$, hence $\mathrm{MSE}(E) = \mathrm{MSE}(S) + \mathrm{MSE}(R) + 2\,\mathrm{mean}(SR)$. Averaging over uniform subsets preserves this identity; negative cross terms can cancel squared errors. Neither RMSE subtraction nor normalized squared terms are independent causal contributions.

S37.2. Larger-budget source-role calibration

The source calibration union contains 2370 cases at 19 stations and 102 unique station-years. The Swiss/Bavarian target roles contain 4/4 positive-weight interval-calibration groups under the original saved role assignment. All source calibration years precede 2010; held target groups remain excluded and original integer bootstrap multiplicities are unchanged. For each seed and budget the order is seven raw source quantiles → observed-bound projection → weighted source residual order statistics; target raw quantiles → bound → add source offset → sort → bound. A cell requires at least 50 finite positive-weight cases before multiplicity; otherwise the finite positive calibration pool is used. No target truth is used for offset estimation.

Native20 calls the original bridge with 20 draws, whereas nested20 is the prefix of a 500-draw call; they need not be identical because the random arrays are generated by day. Native20 source and target references are checked separately against original case/repair and archive files. Seeds 0, 1 and 2 share fixed daily models, residual histories and masks. The uncalibrated, fixed-old-offset and updated-source-offset layers score the same target raw draws. D2 retains its original seven annual quantiles, with no refit or recalibration. Every CRPS uses the common bounded normal-spacing 256-point annual approximation;

these are annual scores, not newly computed full-archive trends. The 27 target years span only two connected groups; 500 paths do not prove convergence or broad ranking robustness.

CRPS integration precision.: Keeping every saved annual quantile and the tail assumption fixed, we score 783 method–case distributions on 256, 1,024 and 4,096 probability points. An independent order-statistic formula reproduces the original 256-point scores to floating-point precision. The maximum casewise 256-to-4,096 change is 0.000063°C. The fixed-seed-0 updated-source nested20/nested500 comparison crosses the fixed-D2 CRPS at all three grids. This checks integration of the stipulated seven-quantile approximation; it does not validate its tail assumption or establish infinite-trajectory convergence.

S37.3. Regional and annual water–air influence diagnostics

Removing a connected group and recomputing common cells changes both support and the standardized target population; it is an influence check, not a fixed-population causal contrast. Geometry min-count coverage gaps stay positive, whereas the smaller geometry-plus-reconstructor subset can reverse sign. Training-history imbalance and the missing Swiss air-ridge counterpart remain. Annual water–air sensitivity also depends on whether years or stations receive equal total weight. A group deletion can leave only one Swiss station, and the Bavarian aggregate change masks group influence. None of these checks establishes daily reconstruction stability or equivalence.

S37.4. Same-day missingness with an evaluable-peer denominator

Peer-evaluable rows require another station in the same connected group with an eligible recorded missingness pattern in that year. Observed peers use the eligible-year roster,

which includes incomplete years; trial peers use the complete-reference-year roster on which masks are imposed. Group-years with no such peer are excluded and counted. Same-day overlap means at least one missing day coincides with a peer missing day. Shared station-days count each focal station's missing days that coincide with any peer, divided by its eligible missing station-days. Observed rows are incomplete station-years; trial rows are station-year–replicates. Same-day co-missingness does not identify an outage cause, the donor actually used, or a naturally unobserved failure process.

S38. External reference qualification overview

Strict independent-reference accuracy remains unresolved. Table S66 keeps complete or bounded reference years, stations meeting each history rule, quality and clock, independence, and scored cases in separate fields. Nine Greater Wellington sites stayed at the metadata screen because temperature series were not requested, so their complete hot-year count is unknown (Greater Wellington Regional Council, 2016). The saved screens found zero strict complete years at 24 NEON sites and zero complete July–June thermal years at five Réunion sites; eight new USGS candidates contain 8 complete station-years, but 0 stations reach four years (Marzolf, 2025; Office français de la biodiversité & Bureau de recherches géologiques et minières, 2026; U.S. Geological Survey, n.d.). That USGS count is not an hourly physical reference and does not prove independence. ECCC Red River shares a river with a source station and retains grade and time-basis questions (Environment and Climate Change Canada, 2024; Redoloza et al., 2025). An Australian candidate archive had prior evaluation exposure: 79 of 80 candidate station-years were previously evaluable, including all 10 designated test years (Australian Bureau of Meteorology, 2026). These exclusions do not establish that the observations are erroneous.

Two exploratory comparisons bound what external evidence can show. British Columbia has 4 stations and 16 station-years. Of 372 masks, 288 are controlled gaps and 84 are calendar gaps. Plug-in, D2 and source-role-calibrated stochastic H30 mean absolute errors on the controlled gaps are 0.516, 0.306 and 0.302 °C (Moore & Callahan, 2024; Callahan & Moore, 2025). Three stations share the Fraser basin; daily means may use only 20 hourly observations, their day convention is unresolved, and source quality processing used water–air comparisons (Water Survey of Canada, n.d.). Nova Scotia covers 2 stations, 8 station-years and 144 masks selected after inspecting availability; D2 and stochastic mean absolute errors are 0.311 and 0.313 °C (Government of Nova Scotia, Environment and Climate Change, 2026). Neither comparison is strict independent-climate validation.

The continuous USGS support at Rio Mora is the clearest warning about target definition. Three years have at least one approved unqualified continuous observation in every fixed-standard-time hour. Only 2017 has every quarter-hour slot. The continuous targets for 2012 and 2019 assign unknown quarter-hour values to the conditional range $[-5, 50]$ °C; they are not fully verified physical years. The 54 hourly-supported masks give D2 MAE 0.180°C against the published daily target. Using the original histories, D2 interval coverage is 53/54 against published daily H30 and 33/54 against that conditional continuous target; qualifier-aware histories yield 48/54 and 28/54. The difference can combine clock, rounding, corrections, daily preprocessing and the bound assumption. It does not identify a single source error or justify a universal physical calibration. Truckee scores 72 controlled-gap masks on one station and four reported years. The reservoir comparison scores 306 exploratory bounded-reference cases at 4 stations and 17 target station-years. Both remain descriptive checks of bounds and clock assumptions, not independent accuracy tests.

Table S59. Fixed-input trajectory sensitivity, 20 to 500 nested draws, by seed and region–reconstructor stratum. Med./max. are absolute 95th-percentile changes in °C. Tail count is the number with either endpoint moving by more than 0.05°C. Intervals are uncalibrated; each stratum is one connected group.

Seed	Stratum	n	Med./max.	Tail count	Width 20/500	Covered 20/500
0	Switzerland / donor	18	0.0011/0.1770	4	0.138/0.157	15/16
0	Bavaria / air	9	0.1140/0.4346	6	0.677/0.792	8/9
1	Switzerland / donor	18	0.0033/0.0887	2	0.151/0.153	16/16
1	Bavaria / air	9	0.1057/0.3824	6	0.700/0.821	8/9
2	Switzerland / donor	18	0.0043/0.1369	5	0.153/0.153	16/16
2	Bavaria / air	9	0.0194/0.2570	5	0.790/0.831	9/9

Table S60. Estimated slope RMSE sensitivity to weighting and selection opportunity. Archive, station and group columns weight each respective unit equally within the 805-archive cohort; the last column uses only the 468 nontrivial-choice archives and equal archive weights. These are exploratory summaries, not equivalence tests.

Rule	Archive wt.	Station wt.	Group wt.	Choice-only
Warm-season	0.4459	0.4436	0.3776	0.3944
Uniform exact	0.4492	0.4471	0.3806	0.4008
CDF	0.4284	0.4262	0.3572	0.3594
Sxx	0.4023	0.4003	0.3413	0.3031
All years	0.1697	0.1689	0.1524	0.1711

Table S61. Same-target annual comparison by region–reconstructor stratum. D2 and historical native20 are fixed original references. Updated rows regenerate the source-role offsets at the stated nested budget; entries are minima–maxima across seeds 0, 1 and 2, not confidence intervals. MAE and mean 90% width are in °C. CRPS uses the same seven-quantile normal-spacing 256-point approximation for every method. Each stratum contains one connected group; these counts do not estimate population coverage or establish a general method ranking.

Method / paths	Covered	MAE	Width	CRPS256
Switzerland / donor ridge; 18 target station-years				
Fixed D2	18/18	0.0569	0.7347	0.0483
Historical native20	18/18	0.0785	0.5623	0.0554
Updated nested20	18 of 18	0.0632–0.0666	0.5802–0.6716	0.0504–0.0509
Updated nested100	18 of 18	0.0642–0.0724	0.5677–0.5715	0.0490–0.0521
Updated nested500	18 of 18	0.0693–0.0730	0.5334–0.5533	0.0500–0.0516
Bavaria / air ridge; 9 target station-years				
Fixed D2	9/9	0.2311	1.0531	0.1634
Historical native20	9/9	0.2435	0.7612	0.1530
Updated nested20	8–9 of 9	0.1914–0.2321	0.7357–0.8804	0.1281–0.1621
Updated nested100	8–9 of 9	0.1824–0.2129	0.7526–0.8547	0.1262–0.1467
Updated nested500	9 of 9	0.1915–0.2015	0.7607–0.8006	0.1287–0.1334

Table S62. Updated source-offset annual results for every nested seed and budget. Panels retain the original region–reconstructor strata. Counts are descriptive on one group per panel; all scores use the same bounded 256-point annual approximation.

Seed / paths	n	Covered	MAE	Width	CRPS256
Switzerland / donor ridge					
0 / 20	18	18	0.0666	0.5802	0.0504
0 / 100	18	18	0.0700	0.5677	0.0521
0 / 500	18	18	0.0730	0.5334	0.0513
1 / 20	18	18	0.0649	0.6716	0.0507
1 / 100	18	18	0.0724	0.5715	0.0513
1 / 500	18	18	0.0700	0.5533	0.0516
2 / 20	18	18	0.0632	0.6092	0.0509
2 / 100	18	18	0.0642	0.5711	0.0490
2 / 500	18	18	0.0693	0.5503	0.0500
Bavaria / air ridge					
0 / 20	9	8	0.2321	0.7949	0.1621
0 / 100	9	8	0.2129	0.7526	0.1467
0 / 500	9	9	0.2015	0.7607	0.1334
1 / 20	9	9	0.1914	0.7357	0.1281
1 / 100	9	9	0.1845	0.7643	0.1262
1 / 500	9	9	0.1925	0.7740	0.1323
2 / 20	9	9	0.1960	0.8804	0.1308
2 / 100	9	9	0.1824	0.8547	0.1317
2 / 500	9	9	0.1915	0.8006	0.1287

Table S63. Both endpoint sensitivities after source-role offset re-estimation, from nested-prefix20 to 500 paths. Entries are the median/maximum absolute change in °C across the stated target stratum; the last column counts cases with either endpoint changing by more than 0.05°C. Each region–reconstructor stratum is one connected group, and these descriptive changes do not establish convergence.

Stratum	Seed	n	$\lvert\Delta q_{.05}\rvert$ med./max.	$\lvert\Delta q_{.95}\rvert$ med./max.	Either > 0.05
Swiss donor	0	18	0.0000/0.1353	0.0691/0.1159	11/18
Swiss donor	1	18	0.0000/0.0561	0.1169/0.2619	13/18
Swiss donor	2	18	0.0000/0.0388	0.1024/0.2667	12/18
Bavarian air	0	9	0.0099/0.2223	0.0609/0.3688	7/9
Bavarian air	1	9	0.0433/0.2918	0.1011/0.3140	6/9
Bavarian air	2	9	0.0034/0.4496	0.0670/0.2367	7/9

Table S64. Influence checks. Panel A shows FOEN-minus-Bavaria coverage gaps in percentage points using min-count weights; group deletion recomputes common support. Panel B shows after-minus-before-2010 within-station annual slopes; equal-station weighting changes the target relative to equal annual observations. Deletion ranges are not confidence intervals.

Panel A: cells	Main gap	Group-deletion gap	Sign reversals
Geometry	16.037	12.392–17.772	0
+ reconstructor	3.686	-1.120–5.292	2
Panel B: region	**Annual-weight Δ**	**Station-weight Δ**	**Group-deletion Δ**
Switzerland	-0.0143	-0.0137	-0.0152 to -0.0047
Bavaria	-0.0054	-0.0343	-0.0882–0.1089

Table S65. Same-day overlap denominators. Any overlap is the percent of peer-evaluable incomplete rows with at least one shared missing day. Day share uses eligible missing station-days. Repeated trial masks retain their original weights; the historical observed row supplements the primary 2010–2025 comparison.

Source / period	All	Peer	Excl.	Any (%)	Shared / eligible days	Days (%)
Observed / 2010–25	109	74	35	18.92	40/2,086	1.92
Calendar / 2010–25	3,806	2,810	996	36.01	4,946/41,507	11.92
Controlled / 2010–25	3,777	2,779	998	58.58	64,164/165,870	38.68
Observed / 1981–25	220	147	73	32.65	158/3,946	4.00

Table S66. External reference qualification. Complete or bounded reference years, stations meeting the stated history rule, quality and clock, independence, and scored cases are separate fields. Unknown is written as unknown. A zero appears only when the saved screen counted that quantity.

Source	Reference years	History-rule stations	Quality and clock	Independence	Scored cases
Greater Wellington	Temperature series not acquired	9 potential sites; quality-code screen not completed	Quality-code values were not requested	Unknown	Not scored
NEON	0 strict complete years	0 of 24 sites meet the four-year strict rule	UTC day and final quality flag required	Climate independence not established	Not scored
New USGS candidates	8 complete station-years	0 of 8 sites reach four complete years	Daily screen only; hourly reference not verified here	Not established	Not scored. The observation-only witness check is separate.
Réunion (Hub'Eau)	0 complete July–June thermal years	0 of 5 sites meet the four-year rule	Reported-day time zone unresolved	Unknown	Not scored
ECCC Red River	0 complete local wall-clock years	Strict physical reference not met	Grade and time basis unresolved	Shares a river with a source station	Not scored
Australia candidate	80 candidate station-years; strict 24-hour sampling not established	79 of 80 were previously evaluable, including 10 of 10 designated test years	The saved screen allowed 20 observed hours	Not independent: prior evaluation exposure	Not scored as a new test
British Columbia	16 published station-years; 0 strict 24-hour stations	4 stations scored; 0 meet the strict 24-hour rule	Days may contain 20 hours; clock unresolved	Not established; three stations share the Fraser basin	372 masks total: 288 controlled gaps and 84 calendar gaps. The reported MAEs use the 288.
Nova Scotia	8 reported-clock station-years	2 stations; strict reference not met	Source QC and time zone unresolved	Not established	144 controlled-gap masks scored
Truckee, California	4 reported years	1 station; strict reference not met	Clock and estimation flags unresolved	Not established; one managed station with prior basin contact	72 controlled-gap masks scored
USGS Rio Mora	Complete quarter-hour year: 2017. Bounded years: 2012 and 2019 (2012: 3 unknown slots; 2017: 0 unknown slots; 2019: 2 unknown slots)	1 station	Unknown quarter-hour slots are assigned to the conditional range $[-5, 50]\,^{\circ}\mathrm{C}$	Not an independent climate	54 hourly-supported masks scored against the reported daily target and the conditional continuous target
Reservoir network	0 complete finite years among 69 station-years	0 of 10 files have four finite years	Strict quarter-hour calendars were not met. The scored bounds assume missing values lie in $[-5, 50]\,^{\circ}\mathrm{C}$	Not established	306 exploratory bounded-reference cases scored at 4 stations and 17 target station-years

S39. Post-hoc retained-truth and point-summary checks

These checks reuse the saved forward and offline predictions. They do not refit models, open new stations, or supply an independent validation. Replacing a retained annual value by the experimental truth cannot be done when that year is actually missing.

On each common station–repetition, the warm-season year set is held fixed and every retained D2 annual value is replaced by the known complete-record H30 before the slope is recomputed. The resulting root-mean-square error is the error of the year set alone. Table S67 shows that this truth-in-set error is nearly the same as the error of the saved D2 values, especially for forward controlled gaps (0.445242 versus 0.445331 °C per decade). For a finite complete record, the warm-season screen changes the slope mainly by changing which years and ordinary-least-squares weights remain. Making the retained years exact does not restore the years that were removed.

The two root-mean-square errors are not subtracted, and neither is divided by the other to name a percentage caused by retention. The squared error of the warm-season slope relative to the complete-record slope also contains the cross term between the year-set change and the reconstruction error on the retained years. The four-cell symmetric decomposition in the main text keeps its original definition. The target remains the slope of a particular complete archive, not a latent climate-process trend.

The main analysis publishes the annual median and then scores the trend of that published series. A conditional mean is the relevant point for squared error of a slope when the conditional distribution is correct; the median does not automatically have that property. Table S68 replaces only the annual point, using the mean of the same approximate distribution on the 738 archives where both methods have a forward controlled-

gap result. The mean still depends on interpolation between the seven quantiles and on the normal-spacing tails. Both root-mean-square errors change slightly, and the original ordering does not reverse. This is not a claim that the mean is a verified optimum, and it does not replace the published-median analysis.

Table S67. D2 slope error when retained years keep their saved annual values or are replaced by the known complete-record H30. Units are °C per decade. The truth replacement is an evaluation device.

Gaps	Timing	Common settings	All years	Warm season	Warm season, true H30
Calendar	Offline	840	0.008732	0.024041	0.023985
Controlled	Offline	840	0.027772	0.100603	0.100603
Calendar	Forward	820	0.067277	0.113458	0.111852
Controlled	Forward	807	0.169474	0.445331	0.445242

Table S68. Trend root-mean-square error, in °C per decade, when the annual point is the published median or the mean of the same approximate distribution. Both use the 738 common forward controlled-gap archives.

Method	Published median	Approximate-distribution mean
D2	0.163814	0.157811
Calibrated stochastic	0.146577	0.143719

S40. Independent observed-record arithmetic re-audits

Using the same local canonical float32 daily archive and an independent convolution implementation, we recomputed 2,139 complete reference station-years and 545 incomplete station-years at 96 eligible stations. All 35 witness identities, saved 30-day values and complete-year reference maxima agreed within the stated numerical tolerance. This is an arithmetic re-audit of the same processed archive, not an independent check of provider-native records or sensor error.

Comparing 971,673 finite values from 34 selected project input panels with the canonical archive yielded no missingness or float32-cast discrepancies. The largest absolute rounding difference is 0.00000153 degrees C; 0 witness identities and 0 classifications at the 0.5-degree excess threshold changed. The input panels are processed project files, not provider-native instrument records.

Across 2,700 eligible station-years, independently recalculated annual coverage, available 30-day maxima, July–August completeness and longest missing runs reproduced 210 witness–rule decisions. For each of 35 witness years, increasing its observed values by 5 degrees C did not change the leave-target-year-out warm-season climatology. The old fixed-season fraction remains sourced from saved metadata; these probes do not cover every year or other pipelines.

For the ten additional USGS sites, independent parsing of the saved daily API responses reproduced 8 approved-complete reference station-years and the 9 certified observed windows. The original quality-screened window selection also replays from the saved payloads, but the separate arithmetic audit does not independently implement that selection. Approved estimated values remain in complete-year references, but the certified

windows contain no estimated days. These records do not establish subdaily completeness, physical measurement accuracy or independent recovery performance.

A separate candidate review subset contains processed daily values for 95 of 96 previously eligible source stations, including all 35 witnesses. One LUBW station is omitted pending permission. The subset verifies the included reported-series arithmetic and rule decisions, but cannot independently confirm the omitted station or the original full eligible-station denominator.

References

Australian Bureau of Meteorology. (2026). *Water data online: water temperature time series.* http://www.bom.gov.au/waterdata/.

Callahan, L., & Moore, R. D. (2025). Evaluation of the hybrid air2stream model for simulating daily stream temperature during extreme summer heat wave and autumn drought conditions. *Hydrological Processes*, *39*(1), e70033. doi: 10.1002/hyp.70033

Environment and Climate Change Canada. (2024). *Automated fresh water quality monitoring and surveillance data.* Retrieved from https://open.canada.ca/data/en/dataset/f258b0c8-7871-4572-b567-1ba2bd55f1b6 (Open Government Licence; ECCC Red River files and metadata accessed October 2026)

Government of Nova Scotia, Environment and Climate Change. (2026). *Surface water quality monitoring network continuous water quality data.* Retrieved from https://data.novascotia.ca/datasets/bkfi-mjgw (Open Government Licence – Nova Scotia; accessed October 2026)

Greater Wellington Regional Council. (2016). *Environmental monitoring: quality codes.* Retrieved from https://cms.graphs.gw.govt.nz/envmon/whats-new (Quality-

code explanation; accessed October 2026)

Marzolf, N. (2025). *Ecosystem metabolism estimates from the national ecological observatory network stream and river sites.* figshare. (Public author archive, version 1) doi: 10.6084/m9.figshare.28074452.v1

Moore, R. D., & Callahan, L. (2024). *Evaluation of the hybrid air2stream model for simulating daily stream temperature during extreme summer heat wave and autumn drought conditions: data sets and scripts.* Zenodo. (Version 1, CC BY 4.0) doi: 10.5281/zenodo.14502248

Office français de la biodiversité, & Bureau de recherches géologiques et minières. (2026). *Hub'Eau: continuous stream temperature data.* Retrieved from `https://hubeau.eaufrance.fr/api/v1/temperature/chronique` (Quality-qualified Réunion records retrieved October 2026)

Redoloza, F. S., Glas, R. L., Nustad, R. A., & Ryberg, K. R. (2025). *Evaluating drought risk of the red river of the north basin using historical and stochastic streamflow upstream from emerson, manitoba* (Tech. Rep.). U.S. Geological Survey. Retrieved from `https://pubs.usgs.gov/publication/sir20255002/full` (Identifies the Emerson ECCC station and USGS streamgage alias) doi: 10.3133/sir20255002

U.S. Geological Survey. (n.d.). *Statistics documentation: supporting information for daily data statistical values.* Retrieved from `https://waterdata.usgs.gov/statistics-documentation` (Approved daily sample counts, accessed October 2026)

Water Survey of Canada. (n.d.). *Frequently asked questions: water level and flow.* Retrieved from `https://wateroffice.ec.gc.ca/contactus/faq_e.html` (Temperature is an auxiliary output without standardized quality assurance; accessed October 2026)