Abstract Thermal archives are usually screened by completeness thresholds that measure outage size, not the support of the target statistic. We develop a framework that conditions on how much of that support a gap hides, links it to reconstruction error and archive admission, and calibrates risk. On 5114 complete river station-years from 146 networks (51524 designed outages), a frozen hottest-30 screen on a retrospective 36-network holdout cuts consequential errors from 9.7% to 2.5% at 70% retention (about [−8.97, −5.36] points); the empirical arm is unresolved. A sealed never-used USGS panel (39 HUC4 networks; 8973 cases) scored once recovers about −7.9 points ([−10.2, −5.8]), with a smaller non-zero empirical-arm gain; multi-agency confirmation remains open. Continuum, crossover and dilation results from the opened reserved panel stay exploratory outside that seal. On 2422 real incomplete years, a peak-window rule disagrees with length guidance on 15.1% of admissions and flags wholly masked *climatological* peak windows (a proxy). Difficulty-based screening can thin high-elevation catchments unless stratified.

Plain Language Summary Rivers are getting warmer, and managers ask how hot a river got last summer. The answer needs a year of daily temperatures, but records have holes. Screening today mostly counts missing days. We argue the better question is whether the missing days are ones the number you want depends on, and we test a screening rule built that way on many river networks. On held-out networks with designed gaps the rule reduces large errors for the summer peak at equal retention; under the gaps these networks actually experience the gain is not yet clear. Results about exact crossover days, autumn maintenance benefits, and "peak never observed" counts are exploratory or use proxies and need confirmation before they guide operations.

This paper has not been peer-reviewed

1. Introduction

The quantity this paper is about is the error of a derived thermal statistic, not the error of a daily temperature series. Cold-water habitat assessment, thermal-regime classification and criterion-exceedance accounting all reduce a year of daily water temperature to a single number: a hottest 30-day mean, an annual mean, a seasonal amplitude (Caissie, 2006; Isaak et al., 2017). That number is what enters a management judgement, and a station-year is admitted to an archive or excluded from it on the strength of it. An error of one degree in a summer peak converts a thermally stressful year into an unremarkable one; on our panel that error is not hypothetical, because computing the peak from recorded days alone carries a mean absolute error of 1.305 °C, larger than one year-to-year standard deviation of the quantity itself.

The error of a derived statistic is not the same object as the error of the series it is computed from, and the difference is what makes the admission decision hard. A reconstruction with excellent daily skill can be badly wrong about a 30-day maximum if its errors are correlated inside the window that achieves the maximum, and a reconstruction with mediocre daily skill can recover that maximum exactly if the window survived. Daily skill is what the modelling literature reports; the derived statistic is what a manager reads. Between the two sits a decision nobody writes down — whether this station-year may be used at all — which is made for every record in every thermal archive, is made by a rule, and is almost never made with the derived statistic in mind.

Two families of rule decide admission today, and they differ in what they assume rather than in when they were written (Table 1). Agency collection standards specify what must be present: Mauger, Shaftel, Trammell, Geist, and Bogan (2015), writing minimum standards for a state whose records feed regional analyses, require a minimum sampling duration of one calendar month and recommend a deployment period of 1 June to 31 August "to capture the thermal maxima". Quantitative guidance instead constrains what may be absent: Johnson et al. (2021) report that "datasets with less than seven weeks of consecutively missing data enabled estimation of all annual signal parameters with reasonable accuracy" and that "imputation of missing data expanded this range to approximately 20 weeks". The gap-filling literature optimises the accuracy of the filled series rather than the admissibility of the completed year (Mohseni et al., 1998; Li et al., 2017), and the closest prior work fills incomplete summer records so that a thermal peak can be computed, but applies the filling uniformly and declines no station-year (Beaufort et al., 2022). Irregular sampling and sampling artifacts already bias mean and trend estimates in stream-temperature series (Grey et al., 2024),

and long-term reconstruction frameworks have been developed for regulated and data-sparse basins including Chinese catchments (Huang et al., 2023). Those studies answer prediction and reconstruction questions; they do not yet close the archive-admission question for a named annual estimand. Across the completeness thresholds in common use the conditioning variable is a property of the record. None is a property of the statistic being computed from it, and the transfer does happen in practice: Mauger, Shaftel, Leppi, and Rinella (2017), for example, retain a site-year for a mean July temperature only "when at least 90% of the days were captured", and then report the maximum weekly maximum temperature from the same records.

That is the assumption we examine, and its consequence is not confined to a prediction. An annual mean is a sum over days: removing k days removes k terms and it matters little which. A hottest 30-day mean is an extreme supported on a short window: if the outage misses that window the statistic is recoverable almost exactly however long the outage is, and if the outage covers it the statistic must be inferred however short the outage is. A single completeness threshold therefore cannot be correct for both, and the direction of its error is opposite in the two cases. Applied to a summer extreme, a threshold on gap length can admit station-years whose climatological peak window is wholly masked and discard station-years whose realised peak is intact; which of the two dominates is an empirical question about the archive, not a matter of taste. The consequence is an archive whose composition, and not only whose accuracy, depends on a choice nobody records.




Figure 1: Problem frame (schematic). The same station-year and outage mask feed an annual mean and a hottest 30-day mean; different supports imply different governing axes. Boxes that quote numerical gradients are mechanism diagnostics from the scored panel, not the sealed never-used confirmation.

Three things make the question answerable now that were not available to the studies that raised it. Water-temperature archives from nine national agencies can be assembled into a panel large enough to hold out whole river networks, so a screening rule can be scored on held-out river networks, and — after external recruitment — on networks never used anywhere in the programme. Because a fully observed calendar year supplies a truth, a gap can be placed on it deliberately, which converts the comparison between conditioning variables from an observational contrast into a designed one: the two candidate properties of a gap can be varied independently, which they never are in a record that failed on its own. And the share of a statistic's support that a gap hides is computable at decision time from the station's own earlier years, so the alternative

This paper has not been peer-reviewed

Table 1: What each body of work optimises, decides on, and conditions on. The rows are ordered by what they assume about the conditioning variable, not by date.

body of work	optimises	decision unit	conditioning axis
completeness guidance (Johnson et al., 2021)	tolerable amount of missing data	station-year	outage size
collection standards (Mauger et al., 2015)	comparability across a programme	deployment	season and duration
gap filling (Mohseni et al., 1998; Li et al., 2017)	accuracy of the filled series	day	air temperature, donors
peak reconstruction (Beaufort et al., 2022)	spatial coverage of the thermal peak	reach	air regression, applied uniformly
large-sample models (Isaak et al., 2017; Corona & Hogue, 2025; Kratzert et al., 2023)	prediction skill	day or reach	meteorology, catchment attributes
selective prediction (Geifman & El-Yaniv, 2017; Zaoui et al., 2020)	error at a fixed rejection rate	case	model confidence
this work	**error of the derived statistic among retained years**	**station-year**	**share of the statistic's support hidden**

conditioning variable is not merely definable but deployable by an agency with no data it does not already hold.

We therefore ask three linked questions, and the last matters most, because a screening rule is not evaluated by its discrimination but by the archive it leaves behind. First, which property of a data gap governs the error of a derived thermal statistic, and does the answer depend on which statistic is being computed? Second, if it does, is that dependence a dichotomy between two regimes or a continuous transition, and where does it turn over? Third, what does conditioning on the answer change — for the error among retained station-years, for which stations survive in the archive, and for the 2422 records in this same archive that really do have gaps and have never been decided?

2. Study system, data, and the quantity to be explained

2.1. The statistic, its support, and the share a gap hides

Let a station-year be a sequence $y = (y_1, \dots, y_T)$ of daily mean water temperatures, $T \in \{365, 366\}$, and let $q(\cdot)$ be the derived statistic of interest. The family we sweep is the hottest consecutive k-day mean

$$q_k(y) = \max_t \frac{1}{k} \sum_{i=t}^{t+k-1} y_i, \tag{1}$$

which is the annual maximum at $k = 1$ and the annual mean at $k = T$; the coldest 30-day mean, the annual amplitude, and the sine-fit mean, phase and amplitude of Johnson et al. (2021) are scored alongside.

A gap is a binary mask $m \in \{0, 1\}^T$; the analyst observes the days where $m_t = 0$. A reconstruction $\hat{y} = R(y_{\bar{m}}, Z)$ completes the year using side information Z — synchronous donor stations, air temperature, the station's own history — and reports $q(\hat{y})$. The quantity that matters is the error of the reported statistic,

$$L = |q(\hat{y}) - q(y)|, \tag{2}$$

and a station-year is called *consequential* when $L > \tau$ for a tolerance τ in the units of q.

The property of the gap we propose as the governing one follows from how much each day moves the statistic. Writing $w_t^q = |\partial q(y)/\partial y_t|$ for the discrete influence function of q at y (Hampel, 1974), the *hidden mass* of a gap is the share of that influence it removes,

$$h_q(m) = \frac{\sum_{t:\, m_t = 1} w_t^q}{\sum_t w_t^q}. \tag{3}$$

For the annual mean every w_t is equal and h_q is the missing fraction; for the hottest k-day mean the influence is concentrated on the k days achieving the maximum and h_q is the share of that window the gap hides; for

This paper has not been peer-reviewed

the sine-fit parameters the weights are smooth functions of day of year — a constant for the fitted mean, $|\sin(\theta_t - \phi)|$ for the amplitude, which is largest at both extremes, and $|\cos(\theta_t - \phi)|$ for the phase, which is largest a quarter-year from the peak. Equation (3) therefore contains the incumbent axis as a special case rather than opposing it, which is why the two regimes reported below are ends of one scale.

Equation (3) cannot be evaluated at decision time, because the weights depend on the days the gap hides. What an analyst has is the *climatological* support: the window the station's other complete years say the maximum should fall in. That estimate has an error, and Section 4.5 shows the error is large enough to change what a screening rule should condition on.

2.2. Records, and what each dataset is for

Daily water-temperature series were assembled from nine national providers: ARSO (Slovenia), CHMI (Czechia), FOEN (Switzerland), GKD Bayern and LUBW (Germany), NVE (Norway), Rijkswaterstaat (Netherlands), SYKE (Finland) and the USGS (United States). Acquisition contracts differ. USGS series are requested as NWIS daily means (`parameterCd=00010`, `statCd=00003`); NVE HydAPI series are measured daily river temperature; Rijkswaterstaat values are aggregated to daily means in the project parser. Several other providers publish daily tables whose native statistic (24 h mean versus a timed reference reading) is not yet proven station by station. In particular, SYKE's public day values can be morning nearshore references rather than 24 h means; the three SYKE stations in the panel are therefore flagged in `results/wrr_submission_` and should be read as a sensitivity stratum, not as proven daily means. A *network* is one provider's group of stations on a common river system. It is the unit at which donor stations are available to one another, and it is the clustering unit for every interval reported here, because station-years within a network share weather, regulation and instrumentation.

Waterbody class was assigned per network and non-river objects — seas, lakes, canals, estuaries, deltas — are excluded rather than averaged into a river result. Reference series are never imputed: a calendar year with any missing day supplies no truth. Stations with fewer than four fully observed calendar years are excluded because the estimator protocol requires them. This leaves **146 networks, 457 stations and 5114 fully observed station-years**.

The same 457 stations contribute **2422 further station-years that are genuinely incomplete**. These carry no truth and are therefore not part of the scored population; they are the population a screening rule exists for. Section 4.12 reports overlap-rule decisions on that panel; the frozen certificate can also be applied through a truth-free `predict` interface (repository amendment `results/wrr_submission_ready/e2_cer`

Station coordinates were assembled for all 457 stations, from the providers' own catalogues, from the USGS NWIS site service for 195 United States gauges, and from ARSO's public station feed, and daily 2 m air temperature was joined from NASA POWER for the whole panel, 1981 to 2026 (NASA Langley Research Center, 2026). For the United States subset, basin characteristics were joined from GAGES-II (Falcone, 2011) on the eight-digit gauge identifier, matching **140 of 204** eligible stations across 57 networks. Basin attributes are used to ask which rivers are hard (Section 4.7) and which rivers a screen keeps (Section 4.11); they are deliberately not used inside the screening rule, because they are unavailable for the other eight agencies.

Table 2 lists every quantity the analysis uses, what it means physically, where its value comes from, and whether it was calibrated. Networks — never station-years — were split into development and reserved, stratified on provider and on donor regime, with the seed recorded before any scoring. This gives **110 development and 36 reserved networks**. That reserved panel is a retrospective network holdout for the frozen v17 screen: 28 of 36 networks have prior development, descriptive or scoring use in earlier programme stages (`results/wrr_submission_ready/prior_access_ledger_reserved_networks.csv`). It is not an unused confirmation set for analyses developed after opening. A sealed confirmation was therefore recruited externally: 39 never-used USGS HUC4 networks (91 stations; 18 HUC2 contexts; 8973 designed outage cases) were ingested, hashed, and scored once with the frozen certificate (`results/wrr_submissio`
At 70% retention and 0.5 °C the pooled contrast is about −7.9 percentage points (clustered interval about [−10.2, −5.8]); the empirical arm is smaller but excludes zero (about −1.7 points, [−3.0, −0.7]). That panel is USGS-only and uses provisional HUC4 groupings rather than agency river systems. A never-used NVE

This paper has not been peer-reviewed

probe reached only two basins with qualifying complete years (62 cases) and does not support transfer (*p0_07_nve_probe/verdict.json*); multi-agency confirmation therefore remains open by an explicit failure closure rather than by silence.




Figure 2: Study design and data flow from nine providers through quality control, the network stratified split, and the freeze. The v17 reserved panel is a retrospective holdout with mixed prior access; sealed never-used confirmation uses a separate USGS ingest (and a thin NVE probe).

2.3. Gap design: two arms, and why both are needed

A case is one station-year under one gap realization. Two arms are used and reported separately, because they answer different questions and are not one population. The *empirical* arm samples real gap runs — length and start month — from the same network's own records and transplants them onto fully observed years, so that operational numbers reflect the geometry agencies experience. The *stratified* arm crosses gap length $\{14, 30, 60, 90, 120, 180, 270\}$ days with the overlap between the gap and the station's climatological peak window, so that the response surface is estimable rather than accidental. The design yields **51524 cases**, 40704 development and 10820 reserved.

Both arms transplant a mask onto a year whose values were observed, which assumes the mask is uninformative about the values it hides. Section 4.13 tests that assumption against the archive rather than asserting it. Gaps are placed inside one calendar year; failures that straddle New Year damage two annual statistics while hiding neither warm season, and the design cannot represent that case.

Each endpoint is scored against a tolerance equal to the same fraction, 0.47, of *its own* interannual standard deviation at the same stations. For the annual peak that scale is 1.06 °C, giving a primary tolerance of 0.50 °C; the scale falls monotonically with support width, from 1.18 °C at $k = 1$ to 0.55 °C at $k = 365$. The fraction is swept at 0.25, 0.47 and 1.0, and no conclusion about which axis wins is claimed unless it is invariant across

This paper has not been peer-reviewed

Table 2: Quantities used in the analysis.

quantity	physical meaning	source	calibrated
h_q, hidden mass	share of the statistic's influence removed by the gap	mask and the station's own complete years	no
peak-window overlap	h_q for a 30-day extreme, from the climatological window	mask and training years	no
warm-season coverage	share of May–September days observed	mask	no
longest run, missing fraction	size of the gap	mask	no
observed-peak margin	how far the observed days already constrain the annual maximum	observed days and training climatology	no
donor count, coverage, correlation	how much synchronous anomaly information neighbours carry	training-period deseasonalised series	no
K, ridge penalty of the donor and air estimators	strength of shrinkage toward the seasonal mean	leave-one-year-out within station	yes, within station
conformal scale	multiplicative factor turning a predicted bound into a valid one	calibration networks, disjoint from fitting and test	yes, on networks
τ, tolerance	consequential error in the units of the statistic	0.47 of the endpoint's own interannual SD; swept 0.25–1.0	no, declared
basin attributes	elevation, slope, baseflow index, drainage area	GAGES-II, United States subset	no, not an input

the sweep. The reference series carries instrument error of roughly 0.1 to 0.2 °C, so the primary tolerance is about three times sensor accuracy; Section 4.13 perturbs the reference by 0.15 °C and repeats the analysis.

3. Methods

3.1. Reconstruction, and what each estimator is allowed to see

Every case is completed by each supported estimator: the statistic computed from recorded days alone (current practice), a day-of-year median climatology, a smoothed climatology, linear interpolation where both flanks exist, ridge regression on the visible subset of synchronous donors, and an air-driven ridge with lagged air temperature and harmonics. Estimators are fit leave-one-year-out within the station, which is what offline archive repair permits; Section 4.13 repeats the study under the stricter protocol in which only prior years are available. Support is declared per method: donor regression does not read a flanking observation, so requiring one would discard usable cases, and absent donor columns are never imputed but the model is refit on the visible subset.

One weakness in that library was worth removing before the comparison was run. Air temperature was available for 2.3% of development cases in an earlier version of this analysis, so the reconstruction this field actually uses was effectively absent, and 36 of the 110 development networks have no synchronous donor at all. With coordinates joined for all 457 stations and NASA POWER pulled for the whole panel, air forcing now covers 98.6% of development cases and 100% of reserved cases. The retrospective reserved holdout is not re-scored against the new forcing; Section 4.9 reports it as a development-panel sensitivity analysis.

The reconstruction layer is not the contribution of this paper and is kept deliberately simple. Its role is to make the comparison between conditioning variables fair, which requires only that the estimator paired with the incumbent rule is the strongest deployable one; Section 4.9 tests that requirement directly rather than assuming it.

3.2. The screening rule, and what it may not read

For each case the screen predicts an upper bound on the absolute error each supported estimator would produce, selects the estimator with the smallest bound, and emits a conformal interval on the statistic together with an admission decision at an analyst-chosen tolerance. Its inputs are the first six rows of Table 2. No input may read a day the gap hides, and that restriction is enforced in code rather than by convention.

Calibration is split conformal (Lei et al., 2018) with two properties that matter. It is *selection-conditional*,

This paper has not been peer-reviewed

because taking the smallest of several bounds is a winner's curse that per-estimator calibration cannot see, and *multiplicative*, because the error scale spans an order of magnitude between a gap that misses the peak window and one that covers it. Fitting, calibration and test networks are always disjoint.

3.3. Test protocol

Before any rule was fit we measured, on development cases, how much a rule with perfect foresight could win inside each candidate action space, and compared it with the margin the reserved panel can resolve. Two natural framings — adopt or reject a fixed reconstruction, and choose among reconstruction methods — have ceilings of +0.038 and +0.080 °C against a threshold of 0.136 °C, and were not pursued. Retain-or-exclude clears its gate by a factor of five and is the action space studied here. A decision experiment whose ceiling is below its own resolvable margin cannot produce evidence in either direction, and running one consumes a reserved panel for nothing.

Every interval resamples whole networks, because station-years within a network are not independent and reusing a station-year under several gap realizations does not create independent observations. The pre-registered primary criterion, frozen before the reserved panel was opened, is that at the retention of the matched length rule the screen must reduce the consequential-error rate by at least five percentage points with a network-clustered interval excluding zero. Precision was computed in advance: the reserved panel resolves 1.44 points, so the declared effect is 3.5 times the resolution.

Where several endpoints are compared, the direction of each contrast was predicted in advance from the form of the statistic, which is the defence against multiplicity. Three of those predictions were not met, and Sections 4.5 and 4.6 report them as failures rather than reshaping them into successes.

4. Results

4.1. Computing the peak on recorded days alone costs more than a year-to-year standard deviation

Current practice is expensive before any rule is proposed. On development cases, computing the annual peak from recorded days alone — the default once a station-year passes a completeness screen — gives a mean absolute error of **1.305 °C, or 1.23 interannual standard deviations** of the quantity being measured, with 24.9% of station-years wrong by more than 1.0 °C. Substituting a day-of-year climatology over the gap gives 0.402 °C, the best supported reconstruction 0.225 °C with air forcing available for the whole panel (0.242 °C without it), and an oracle over the estimator set 0.130 °C. The error is therefore dominated not by which station-years are kept but by whether the record is repaired at all, which fixes the size of the question the rest of the paper addresses: given that a year has been repaired, which repaired years can be trusted.

4.2. The calibrated bound is honest enough to interpret

Everything that follows uses a predicted error bound to rank station-years, so the bound is checked before it is used. On the reserved networks the nominal 90% interval covers 90.9% of cases marginally, inside the pre-declared band of 0.87 to 0.93, and 86.3% to 94.9% within every pre-declared stratum (Figure 3a). Median widths adapt with the quantity they are meant to track, from 0.06 °C when the peak window is untouched to 1.87 °C when most of it is hidden.

Those strata pool across networks, and conformal validity assumes an exchangeability that fails between river networks. Coverage inside individual reserved networks has mean 0.904 and median 0.913 but a minimum of 0.778, with five of 36 networks below 0.85. The pooled interval is an honest statement about a population of networks and is *not* a guarantee at a single new site; calibrating on the per-network conformal quantiles at the 0.9 level puts no network below 0.85 for a 1.5-fold widening. The screening results below are identical under every calibration to four decimal places, because the rank order the screen uses is a monotone transform of the bound, so a reader who distrusts the intervals need not distrust the frontier.

This paper has not been peer-reviewed




Figure 3: Conformal coverage by stratum (a), interval width against peak-window overlap (b), and the price of site-level validity (c).

This paper has not been peer-reviewed

Table 3: Discrimination of a consequential error by axis (AUC), development cases, each endpoint at 0.47 of its own interannual standard deviation.

endpoint	peak-window overlap	warm-season coverage	outage length	AUC difference (95% CI)
hottest 7-day mean	0.823	0.824	0.752	+0.072 (0.063, 0.096)
hottest 30-day mean	0.819	0.820	0.750	+0.070 (0.061, 0.098)
amplitude	0.758	0.792	0.768	+0.024 (0.015, 0.037)
annual mean	0.672	0.818	0.845	−0.027 (−0.036, −0.018)
coldest 30-day mean	0.447	0.619	0.803	−0.184 (−0.228, −0.156)

Table 4: The governing axis as a function of the support of the estimand, development cases.

support width k (days)	interannual SD (°C)	AUC, peak window	AUC, missing fraction	difference (95% CI)
1	1.18	0.825	0.751	+0.074 (0.053, 0.097)
3	1.17	0.824	0.746	+0.078 (0.055, 0.104)
7	1.16	0.823	0.746	+0.077 (0.053, 0.104)
15	1.12	0.821	0.743	+0.078 (0.053, 0.106)
30	1.08	0.819	0.744	+0.075 (0.049, 0.104)
60	1.02	0.815	0.765	+0.050 (0.028, 0.078)
90	0.94	0.792	0.790	+0.002 (−0.018, 0.027)
182	0.74	0.728	0.823	−0.095 (−0.111, −0.077)
365	0.55	0.672	0.845	−0.173 (−0.192, −0.157)

4.3. The share of the support a gap hides orders every endpoint, and the ordering reverses

Table 3 reports, for five endpoints, how well each candidate property of the gap discriminates a consequential error. The ordering reverses between extremes and averages, every clustered interval excludes zero, and the winning axis is unchanged at tolerances of 0.25, 0.47 and 1.0 interannual standard deviations. The seven-day peak has the largest timing advantage of any endpoint, which the support account predicted before it was measured because a shorter extreme is more localised in time; peak-window overlap has an AUC of 0.447 for the cold extreme, because hiding the summer peak carries no information about the winter minimum.

4.4. The transition is continuous, and it turns over at a three-month support

Table 3 evaluates the question at two points and reads the gap between them as a dichotomy. Sweeping the hottest-k-day mean of Equation (1) from $k = 1$ to $k = 365$ evaluates it everywhere (Table 4, Figure 4a). The advantage of a fixed peak window is flat at about +0.075 AUC for every estimand whose support is a month or less, decays through +0.050 at a two-month support, and crosses zero near a three-month support on the development width grid (interpolated crossover about 91 days; opened-reserved estimates differ). Beyond that the whole-year completeness axis wins, by 0.095 AUC at a half-year support and 0.173 at the annual mean. These continuum numbers are exploratory analyses developed after the reserved panel was opened; they motivate the screening axis but are not the sealed USGS confirmation reported with the data description. The two regimes of Table 3 are useful ends of that exploratory scale, not a proven universal dichotomy.

The mechanism behind that scale is measurable rather than only plausible. Binning development cases on both properties simultaneously and measuring the growth of error along each with the other averaged out, the hottest 30-day mean grows 3.5× along the missing-days axis and 9.6× along the peak-overlap axis, while the annual mean grows 10.2× and 1.5× respectively: the two ratios swap by roughly an order of magnitude in each direction. For the peak, once the support is in the model, gap length adds nothing measurable — its permutation importance is −0.004 against 0.194 for peak overlap (Figure 5c) — and nested feature sets grouped by network rise from AUC 0.742 with length alone to 0.845 adding the support, 0.858 adding the observed-peak margin and 0.884 adding donor quality (Figure 5b).

Two qualifications belong with that. Length is not irrelevant: holding peak overlap at zero, the consequential-error rate still rises from 0.6% at 1 to 30 days to 8.7% at 121 to 200 days. And in a linear score that already contains warm-season coverage the length term takes a small negative weight, which is collinearity — the

This paper has not been peer-reviewed




Figure 4: Exploratory continuum on the opened panel (not the sealed USGS confirmation). (a) Advantage of a fixed 30-day peak window over whole-year completeness versus support width, with network-clustered intervals; the development-grid crossing is near 91 days and should be read with interpolation uncertainty. (b) AUC for every estimand against every conditioning window. (c) Sine-fit parameters of Johnson et al. (2021), by axis.

This paper has not been peer-reviewed

two correlate at −0.86 — and not evidence that longer gaps are safer. That correlation is also why a timing-versus-length labelling is wrong: warm-season coverage is a count of missing days restricted to a window, which is a length measure with a domain restriction, and it is the domain restriction and not the timing that does the work.

Donor information enters the residual through quality rather than quantity. Halving the donor set by training-period correlation while holding the count fixed costs +0.082 °C (95% interval 0.061 to 0.112), and replacing donor anomalies with phase-shuffled surrogates while preserving each donor's seasonal cycle costs +0.234 °C (0.164 to 0.288), so donors supply synchronous anomaly information and not merely a seasonal shape.




Figure 5: Mechanism: the length-by-overlap reversal (a), nested feature sets (b), permutation importance (c), and the endpoint dependence (d).

4.5. A support must be located before it can be conditioned on, so the useful window is wider than the statistic

Two predictions declared for this round were not met, and both failures have one cause. We predicted that the axis restricted to the estimand's own support would be at least as good as any hand-chosen axis at every k, and that its advantage would decay monotonically. Neither holds: the advantage is a hump peaking at a 60-day support, and for supports of 15 days or less the fixed 30-day peak window beats the estimand's own support, by 0.064 AUC at $k = 1$.

Table 5 gives the reason. The gain from conditioning on a window wider than the statistic tracks the year-to-year wander of the realised window's location almost exactly: it is +0.102 AUC at $k = 1$, where the climatological window is the worst estimate of where this year's window will fall and the wander is 16.7

This paper has not been peer-reviewed

Table 5: Which climatological window a screen should condition on, development cases.

estimand k (days)	best window (days)	AUC there	AUC on own support	gain	support wander (days)
1	90	0.862	0.760	+0.102	16.7
3	60	0.858	0.766	+0.092	16.3
7	90	0.855	0.778	+0.077	16.5
15	60	0.853	0.802	+0.051	15.3
30	90	0.852	0.827	+0.025	13.7
60	90	0.860	0.855	+0.005	9.4
90	90	0.856	0.856	0	7.4
182	182	0.851	0.851	0	6.9
365	365	0.844	0.844	0	—

Table 6: Axis comparison on the sine-fit parameters (AUC), development cases, each at 0.47 of its own interannual standard deviation.

estimand	peak window	missing fraction	outage length	influence-weighted	best axis
fitted mean	0.672	0.845	0.845	0.845	outage length
amplitude	0.724	0.812	0.815	0.831	influence-weighted
phase	0.583	0.835	0.836	0.847	influence-weighted

days, and vanishes at $k \geq 90$, where the wander has fallen to 7.4 days and the support is wide enough to contain its own location error. A screening rule therefore does not condition on the support; it conditions on the support *dilated by the uncertainty in the support's location*, and a dilation of roughly two location standard deviations on each side is what the data choose.

One consequence is practical. AUC at the best conditioning window is between 0.85 and 0.86 for every estimand from the annual maximum to the half-year mean, so an agency needs one climatological window of 60 to 90 days rather than one per metric. A second is uncomfortable: the screen deployed below conditions on a 30-day window and is therefore leaving 0.025 AUC on the table for its own endpoint.

4.6. On the estimands the standing guidance was derived for, the guidance is right

Johnson et al. (2021) fit a sine wave by linear regression and take a mean, a phase and an amplitude; they do not compute a hottest 30-day mean. Reproducing that estimator on our panel puts the two accounts on the same footing for the first time (Table 6, Figure 4c). A completeness threshold is the correct axis for the fitted mean, at 0.845 against 0.672 for peak-window overlap, and is within 0.02 AUC of the best axis for the phase. The ordering is the same under the authors' own protocol, in which the sine is fitted to whatever days are present with no repair at all: peak overlap 0.667, 0.743 and 0.590 against whole-year missingness 0.902, 0.842 and 0.898 for the mean, amplitude and phase, at base rates of 32% to 34% consequential against 10% to 12% after repair.

The third declared prediction failed here. We expected that for the phase, whose influence is largest a quarter-year from the peak, two 61-day windows around the spring and autumn limbs would beat both peak overlap and gap length. They beat peak overlap decisively (+0.249 AUC, 95% CI 0.231 to 0.270) and they do not beat gap length (0.831 against 0.836). Weighting every day by the influence function of Equation (3) instead recovers the advantage: 0.847, which beats the whole-year missing fraction by +0.013 (0.009, 0.017) and peak overlap by +0.265 (0.249, 0.285).

4.7. Topography, not geology, makes a river hard

The account so far is a property of estimators and estimands and would hold for any time series. For the 140 United States stations matched to GAGES-II across 57 networks, we regressed donor synchrony and completed-year error on basin attributes with the expected direction of each effect declared before the regressions were run (Figure 6). Site elevation and basin slope both reduce donor synchrony (Spearman $\rho = -0.29$,

This paper has not been peer-reviewed

95% CI −0.46 to −0.08, and $\rho = -0.31$, −0.51 to −0.07), and site elevation raises both completed-year error ($\rho = +0.24$, 0.06 to 0.43) and the consequential-error rate ($\rho = +0.25$, 0.08 to 0.43).

The groundwater prediction taken from the literature is not reproduced, and the reason is instructive. Johnson et al. (2021) report that filling can harm groundwater-influenced sites, which predicts larger completed-year error at a high baseflow index. Baseflow index does reduce donor synchrony in the predicted direction, from 0.70 in the lowest tertile to 0.59 in the highest, although the rank correlation interval includes zero ($\rho = -0.19$, −0.42 to 0.10). But the error moves the other way: high-baseflow stations are −0.043 °C easier than low-baseflow stations (95% CI −0.081 to −0.004). Across baseflow tertiles the mean annual peak falls from 22.7 °C to 18.1 °C ($\rho = -0.47$, −0.60 to −0.26) and its interannual standard deviation from 0.98 to 0.76 °C (Figure 6c), so two effects act in opposite directions and the second is larger.

Regulation does not make station-years harder for this endpoint: reference and non-reference basins differ by +0.004 °C (95% CI −0.051 to +0.093). Catchment attributes explain a modest share of the variation, a standardised linear model reaching $R^2 = 0.22$ for donor synchrony and 0.13 for completed-year error, and no attribute predicts how much a screen gains: every correlation between a basin attribute and the per-station screening benefit has an interval containing zero. Whether an attribute predicts what a screen *keeps* is a different question, and Section 4.11 answers it differently.




Figure 6: Catchment attribution on the United States subset: donor synchrony and completed-year error against basin attributes, and the decomposition of the baseflow effect (c).

This paper has not been peer-reviewed

Table 7: Reserved panel at 70% retention, across tolerances.

screen	> 0.25 °C	> 0.5 °C	> 1.0 °C	under by > 1 °C
outage-length rule	17.2%	9.7%	3.4%	2.0%
peak-overlap rule	9.2%	4.2%	1.2%	0.7%
calibrated screen	7.1%	2.5%	0.4%	0.3%
random	23.9%	14.4%	5.6%	3.7%

Table 8: Component ablation, development cases at 70% retention, each variant re-run through the full out-of-fold pipeline.

screen	error rate	points vs the length rule
outage-length rule (incumbent)	9.24%	—
three-variable logistic regression	5.53%	−3.71
peak-overlap alone, no fitting at all	5.62%	−3.62
three-feature gradient boosting	4.89%	−4.35
geometry and timing only (8 features)	4.86%	−4.38
drop donor features	3.78%	−5.46
drop the observed-peak margin	3.13%	−6.11
full screen	2.93%	−6.31
full screen, screening only (no method choice)	3.11%	−6.13

4.8. Conditioning on the support cuts the error rate at equal retention, and three variables carry most of it

At 70% retention on development cases, the share of retained station-years with a peak error above 0.5 °C is 2.9% for the calibrated screen, 5.6% for a rule on peak overlap alone, 9.2% for the gap-length rule and 15.0% for random retention (Figure 7a). On the 36 reserved networks scored as a retrospective network holdout (mixed prior access; see the prior-access ledger), the screen gives 2.5% against 9.7% for the length rule (Figure 7b): a contrast of −7.13 points, 95% network-clustered interval −8.97 to −5.36, against the five-point effect declared for that freeze. Both declared conditions are met on that holdout, in every donor regime (−6.6, −6.7, −7.2, all excluding zero). An independent sealed USGS confirmation recovers a similar pooled contrast (Section 2.2). The holdout result is not carried by the largest agencies alone: case-equal −7.01, provider-equal −6.29, network-equal −5.82, station-equal −5.54 points, with a median across individual reserved networks of −5.0, 83% of networks favouring the screen, and a worst network of +2.8.

Placed at the operating points the published guidance implies rather than at an arbitrary retention: at the seven-week unfilled limit (49 days, 49.2% retention on the reserved panel) the length rule leaves 5.2% and the screen 0.3%; at the twenty-week imputed limit (140 days, 85.1% retention) the length rule leaves 11.3% and the screen 6.9%. The second of those is more permissive than an earlier version of this analysis assumed — it admits 85% of cases rather than 74% — and the screen's advantage there is 4.4 points rather than 6.2.

Most of the benefit does not require the calibrated machinery (Table 8). Fifty-nine per cent is available from a logistic regression on three variables and 57% from peak-window overlap with no fitting at all; method selection contributes 0.2 points. Fitted on development networks and applied unchanged to the reserved panel, the deployable rule is

$$\text{risk} \;=\; \text{logistic}\big(-1.302 + 2.070\,x_1 - 2.412\,x_2 - 0.00187\,x_3\big), \tag{4}$$

where x_1 is the share of the station's climatological hottest-30 window the gap hides, x_2 the share of May-to-September days that remain observed, and x_3 the length in days of the longest consecutive gap. It gives 4.2% against the length rule's 9.6% at 70% retention, a contrast of −5.44 points (−6.85, −3.96), and cuts severe underestimation of the peak from 1.9% to 0.6%.

4.9. The incumbent is not handicapped, including by the absence of the field's own reconstruction

A screening comparison can be rigged by pairing the incumbent with a weak reconstruction, so the incumbent is paired here with every reconstruction in the study, including one nobody can deploy (Table 9).

The screen beats the incumbent armed with a reconstruction nobody can run, and the reconstruction paired

This paper has not been peer-reviewed




Figure 7: Retention-error frontier on development cases (a) and on the reserved panel (b).

This paper has not been peer-reviewed

Table 9: Development cases at 70% retention, with air forcing available throughout.

screen	reconstruction	error rate above 0.5 °C
length rule	information-conditioned policy (used throughout)	9.00%
length rule	always air regression, else climatology	15.16%
length rule	always donor ridge, else climatology	9.57%
length rule	always smooth climatology	19.79%
length rule	oracle over methods, not deployable	3.62%
calibrated screen	deployable estimators it selects itself	2.87%

Table 10: Decomposition by outage arm.

panel	arm	cases	base rate	length rule	screen	contrast (95% CI)
development	empirical	14995	7.4%	2.52%	0.38%	−2.13 (−3.45, −1.26)
development	stratified	25709	19.3%	14.58%	6.17%	−8.39 (−9.25, −7.47)
reserved	empirical	4041	4.2%	0.74%	0.11%	−0.71 (−2.36, +0.05)
reserved	stratified	6779	20.8%	16.59%	7.78%	−8.68 (−10.46, −6.51)

with the incumbent throughout is the strongest of the deployable options rather than the weakest. Pairing the incumbent with an air regression in the manner of Beaufort et al. (2022), applied to every record alike, makes it worse rather than better. The axis conclusion does not depend on the estimator library: the timing-minus-length AUC contrast is +0.070 with the old forcing and +0.081 with the new. The screen uses the new information, selecting the air-driven estimator for 20.0% of cases against 0.07% before, and donor regression for 53.1%.

One result here was not expected: the *fixed* information-conditioned policy gets slightly worse at donor-less stations when air becomes available, from 0.340 to 0.364 °C.

4.10. The gain disappears under the gap calendar these networks keep, because long gaps begin in autumn

The frozen holdout test was run on the pooled case set, which mixes two arms whose base rates differ by an order of magnitude, and they must be reported apart (Table 10). The declared five-point effect is met on the pooled and stratified populations and *not* on the reserved empirical arm, where the interval includes zero. The holdout headline −7.13 points is therefore an effect on a mixture that includes a designed arm. The sealed USGS confirmation likewise finds a much smaller empirical-arm gain than the pooled contrast, so neither number should be quoted as next year's agency experience under natural outages alone.

Why the empirical arm is null is itself a result. Across the 6472 real interior gaps in this archive, gaps shorter than 30 days and gaps of 30 to 89 days are close to uniform in start month ($\chi^2 = 4.0$ on 11 df, $p = 0.97$ for the 30-to-89-day class). Gaps of 90 days or more are not: **72.3% of them begin between September and January and 17.9% between March and July** ($\chi^2 = 86.1$, $p = 9.5 \times 10^{-14}$; Figure 8a). The gaps long enough to destroy a summer peak are seasonal shutdowns, and they start after the warm season has been recorded, so little is at risk and little can be screened out: 4.2% of reserved empirical cases exceed the tolerance with no screening at all.

A counterfactual that holds gap lengths fixed and moves only start month is reported in Figure 8b as an exploratory amendment after correcting year-end truncation in the placement code (`results/wrr_submission_r`). Those percentage-point swings are not sealed confirmation and are not used as agency recommendations.

4.11. Conditioning on difficulty removes mountain catchments unless the threshold is applied within strata

A rule that lowers the average error by discarding hard station-years has bought accuracy with representativeness, and in this application the hard sites are not a random subset: Section 4.7 showed they are high-elevation, steep, donor-poor catchments, which are the cold-water headwaters the summer endpoint exists to describe.

This paper has not been peer-reviewed




Figure 8: (a) Start month of 6472 real interior gaps by length class; long outages begin in autumn and winter (descriptive). (b) Exploratory maintenance-calendar counterfactual after multi-year placement correction; not a verified operational benefit.

This paper has not been peer-reviewed

Table 11: Error against representativeness, reserved panel at 70% retention.

rule	error rate	ρ with elevation	donor spread	retention SD
outage-length rule	8.81%	−0.10	4.5 pts	0.075
peak-overlap rule	3.75%	+0.16	2.4 pts	0.104
calibrated screen, global threshold	2.53%	−0.27	4.3 pts	0.149
screen, balanced within donor regime	2.52%	−0.27	0.1 pts	0.148
screen, balanced within elevation tertile	2.85%	+0.08	4.6 pts	0.148
screen, balanced within both	2.84%	+0.07	0.1 pts	0.144

At 70% retention the calibrated screen's per-station retention correlates with site elevation at Spearman $\rho = -0.26$ ($p = 0.011$) on development and -0.27 ($p = 0.077$) on the reserved panel, while the incumbent length rule is elevation-neutral at $\rho = -0.05$ ($p = 0.60$) and -0.10 ($p = 0.52$). On the reserved panel the screen retains 77.7% of station-years in the lowest elevation tertile and 66.9% in the highest, a spread of 10.8 points against 1.7 for the length rule (Figure 9b). By donor regime the same pattern appears on development cases: 57.1% at donor-less stations against 72.5% at donor-rich ones, a 15.4-point spread, while the length rule tilts the other way.

The distortion is cheap to remove (Table 11). Applying the screen's threshold within stratum rather than globally costs 0.31 points of consequential-error rate on development cases and 0.30 on the reserved panel and flattens both distortions. Stratifying on donor regime alone does not fix the elevation tilt and stratifying on elevation alone does not fix the donor tilt, so an agency that cares about both must declare both.




Figure 9: (a) Consequential-error rate against the correlation between station retention and site elevation, reserved panel. (b) Retention by elevation tertile under three rules.

This paper has not been peer-reviewed

Table 12: Archive composition under each admission rule, on the 2422 real incomplete station-years.

rule	admitted	any day hidden	half hidden	wholly hidden
repair everything, no screening	2422	696	261	186
length ≤ 140 d, with imputation	2221	507	82	17
length ≤ 49 d, unfilled	2016	437	45	2
peak window ≤ 10% hidden	1967	241	0	0

4.12. On records that really have gaps, the two rules disagree on one year in six

Every number above is computed on a complete station-year with a synthetic hole punched in it, which is the only way to have a truth. The same archive contains 2422 genuinely incomplete station-years, a third of the total, which carry no truth and have never been decided. Three things can be asked of them without inventing one (Figure 10).

They are not a marginal population. The median incomplete year is missing 9 days and the 90th percentile 128; 28.7% of them have at least one day of the station's climatological peak window missing, and 261 have more than half of it missing.

The two rules disagree on a sixth of them. At the seven-week unfilled limit the length rule admits 83.2% and a peak-window rule at 10% hidden admits 81.2% — near-equal retention, which is the comparison this paper insists on — and they **disagree on 15.1%** of station-years: 207 are admitted by the incumbent and rejected on the support, and 158 are rejected by the incumbent and admitted on it. At the twenty-week imputed limit the incumbent admits 91.7% and the disagreement is 11.6%, of which 267 are in the dangerous direction and 13 in the wasteful one. The disagreement is present in every provider with more than 20 incomplete station-years, from 11.1% to 25.7%.

What the disagreement costs can be counted without a truth (Table 12). The published guideline agencies are most likely to apply, the twenty-week limit that exists because imputation is assumed, admits 17 station-years whose annual peak was never observed at all and 82 whose peak window is mostly gone. A risk model fitted on development cases from decision-time features only, and checked for calibration on the reserved panel first (largest decile miscalibration 6.2 points, and under 1.6 points in nine of ten deciles), predicts a consequential-error rate of 0.98% among the years the unfilled limit admits, 1.85% among those the imputed limit admits, and 0.26% among those the peak-window rule admits at essentially the same retention. That last figure is an extrapolation from scored cases to unscored records and is labelled as one; Section 4.13 tests the exchangeability it assumes.

At the level of a regional aggregate the choice of rule barely matters: admitting incomplete years at all raises the archive's mean annual peak by about half a degree, and the four rules differ by less than 0.1 °C (+0.52, +0.51, +0.53 and +0.45) and by 0.4 points in the share above 20 °C (+4.55 to +4.17).

The same is true of a multi-year slope. Theil–Sen slopes of the annual peak have RMSE 0.143 °C yr^{-1} on recorded days, 0.044 with reconstruction and no screening, 0.048 under length screening and 0.041 under the screen: the dominant step is again whether the record is repaired. Eleven complete years per station is in any case too short to resolve a decadal trend, so we report the slope error as a diagnostic of the reconstruction and make no trend claim.

Per station-year the choice does matter, and Section 5.5 follows it into the judgement an agency actually makes.

4.13. The conclusion survives instrument error, a stricter fitting protocol, and transfer across agencies

Missingness is weakly informative, and the effect is bounded well below the tolerance. Every case transplants a mask onto a year whose values were observed, which is valid only if the mask is ignorable in the sense of Rubin (1976); loggers are lost to floods, buried by ice and fouled in low flow, and each of those coincides with anomalous conditions. Across the 6472 real gaps, the mean anomaly of the seven days preceding a gap, relative to the station's own day-of-year climatology, is +0.018 °C for gaps of a week or less

This paper has not been peer-reviewed




Figure 10: The rules on 2422 real incomplete station-years. (a) Agreement and disagreement at each published limit. (b) Transported risk of each admitted set. (c) Station-years admitted with the peak window more than half, or wholly, unobserved.

This paper has not been peer-reviewed

($p = 0.44$), +0.038 for 8 to 29 days ($p = 0.56$), −0.002 for 30 to 89 days ($p = 0.98$) and −0.215 for 90 days and longer ($p = 0.081$). Donors recording through a gap run +0.063 °C warm during short gaps, detectable at this sample size ($p = 0.017$) and 3% of the within-station anomaly scale. The design is therefore defended on the magnitude of the effect rather than on its absence, and the class that warrants a limitation is gaps of three months and longer.

Three further checks move the numbers and not the conclusion. Perturbing the reference series by 0.15 °C of normal noise moves the development contrast from 6.32 to 6.21 points, the reserved contrast from 7.06 to 6.67 points, and the mean absolute error from 0.2416 to 0.2418 °C. Restricting estimators to years before the gap raises reconstruction error from 0.242 to 0.270 °C on a panel that falls to 33633 development cases, and moves the contrast from −6.35 (−7.12, −5.52) to −5.92 points (−6.72, −4.99) with coverage 0.906. Under leave-one-provider-out the contrast is negative with an interval excluding zero for all nine agencies, from −11.6 points (Finland) to −1.2 (Switzerland), with conformal coverage between 0.82 and 0.96.

5. Discussion

5.1. What process explains the transition

The transition measured in Table 4 is not a property of rivers, of the estimators, or of the screening rule. It is a property of the functional being computed, and it follows from where that functional puts its weight. A statistic whose influence is concentrated on a few days is destroyed by losing those days and is almost unaffected by losing any others; a statistic whose influence is spread evenly over the year tends to degrade with how many days go missing and is less sensitive to which. Between those two limits the influence is neither concentrated nor uniform, and the discriminating power of a narrow conditioning window falls as the weight it captures falls. On this panel's exploratory width sweep, a fixed 30-day window stops beating a whole-year count near a three-month support; that numerical crossing is panel- and climate-dependent, not a universal constant.

Two measurements distinguish that account from a re-description of the data. First, the gradient ratios swap by roughly an order of magnitude in each direction between the peak and the annual mean, which is a direct measurement of how error grows along each axis rather than an inference from discrimination. Second, and more decisively, the account makes a prediction about a case it was not built on. The sine-fit phase has an influence function with two broad lobes a quarter-year from the peak, so the account predicts that peak-window overlap should be a poor axis for it — and it is, at 0.583 against 0.836 for gap length. A description fitted to the extreme-versus-average contrast would not have predicted that.

The same account explains where it failed. We predicted that conditioning on the estimand's own support would always be best, and it is not, because the support has to be estimated and the estimate wanders. Once the wander is measured the failure becomes a quantitative statement rather than an exception: the gain from widening the conditioning window tracks the wander from 16.7 days at $k = 1$ to 7.4 days at $k = 90$ and disappears when the support becomes wide enough to contain it. This is the part of the result we would expect to transfer to other variables and other climates, because nothing in it is specific to water temperature: any estimator conditioned on an estimated support inherits the uncertainty of that support.

5.2. Why this agrees with the standing guidance rather than against it

An earlier framing of this work held that the incumbent rule gets an ordering backwards. Section 4.6 shows that reading is wrong on the guidance's own terms. Johnson et al. (2021) derive their limits for sine-fit mean, phase and amplitude, and for those three estimands a completeness threshold is the correct axis or within 0.02 AUC of it. What we correct is not the threshold but the habit of carrying it across to an estimand it was not derived for.

That reading also resolves the anomaly the guidance itself reported. Its authors observed that the timing of missing data affected thermal-signal estimates "inconsistently in space and time". Their three estimands have three different influence functions — a constant, two lobes at the extremes, two lobes at the transitions

This paper has not been peer-reviewed

— and only one of the three has weight concentrated narrowly enough for a timing effect to appear. Observing all three at once, inconsistency is the only thing one could see. We do not contradict the observation; we decompose it, and we can now say in advance which of their parameters should show a timing effect and which should not.

A smaller result points the same way and was not expected. The fixed information-conditioned policy gets slightly *worse* at donor-less stations once air forcing becomes available, because its preference order was written when air was rare and it now prefers an air regression to a climatology at sites where the climatology is better. A fixed ladder over estimators is not robust to a change in what is available; a per-case selection is. That is an argument for the method of this paper rather than against it, and we report it because it was the opposite of what we expected.

The disagreement with the gap-filling literature is of a different kind and is not resolved by our result. Beaufort et al. (2022) fill every record alike, and on our panel pairing the incumbent rule with that treatment makes it worse rather than better (Table 9). That is not evidence against air-temperature regression, which is the best available estimator at many of our stations and which the screen selects for 20.0% of cases. It is evidence that a reconstruction chosen once for a whole archive and a reconstruction chosen per station-year are different objects, and that the quantity a screening comparison must hold fixed is the second.

5.3. What generalises, and what does not

Three parts of the result are portable. The governing quantity of Equation (3) is defined wherever a derived statistic is computed from an incomplete record, and its ingredients — an influence function and a mask — are available in any such setting. The protocol that measures the transition is portable in the same way: compute the statistic over a family of support widths, measure how far the support's location moves between periods, dilate by that amount, and condition on the result. And the finding that a screening rule must be specified together with its strata is a statement about selection, not about temperature.

Three parts are not. The numerical crossover near three months depends on how concentrated a river's thermal year is and on how far its peak wanders between years; both are properties of this climate and this variable. The operational gain is regime-dependent: the sealed USGS confirmation recovers a pooled contrast, but the empirical arm remains much smaller, and multi-agency never-used confirmation is still thin. Catchment associations rest on 140 of 204 stations in one country and should be read as attribute correlations on that subset, not as a geology-versus-terrain law.

One boundary deserves naming separately, because it limits interpretation rather than transfer. The mechanism argument is supported by an influence-function derivation, by measured error gradients, by a continuum with a located crossover and by an out-of-sample prediction about the phase. It is not supported by an independent observation of the internal quantity itself, because the hidden mass of Equation (3) is by construction unobservable on the records it matters for. A different representation that reproduced the same gradients would not be excluded by our evidence.

5.4. What the remaining uncertainty changes, and what it does not

Uncertainty enters this study in four places, and it is worth separating what each one moves. Measurement error in the reference series changes the numbers in the third decimal and changes neither the mechanism nor the recommendation: 0.15 °C of noise moves the reserved contrast from 7.06 to 6.67 points. Fitting protocol changes the numbers by about half a point and changes nothing else: restricting estimators to prior years moves the contrast from −6.35 to −5.92. Neither of these is capable of overturning a conclusion.

The other two are. The choice of population changes the practical conclusion completely: on the arm built from the gap geometry these networks actually experience the effect is −0.71 points with an interval containing zero, and any reader who needs a number for their own archive needs the arm that matches their gap regime rather than the pooled figure. And the absence of an independent observation of the internal quantity limits the mechanism judgement, as the previous section states: we can show that conditioning on

This paper has not been peer-reviewed

an estimate of the support discriminates better, and we cannot show that no other representation would do as well.

Two further limits belong here because they are specific rather than generic. Missingness is weakly informative and the transplant design is defended on the magnitude of that effect rather than on its absence, which makes the defence weakest for gaps of 90 days and longer, where the leading flank runs 0.215 °C cold. And the archive-level consequence is small: the choice of rule moves a regional mean annual peak by less than 0.1 °C. We report that because it deflates a claim we might have made. At the level of a regional aggregate the decision that matters is whether to repair; the rule matters per station-year, and it matters for who is in the archive.

5.5. What changes in the judgement an agency actually makes

The archive-level aggregate is the wrong place to look for the value of a screening rule, because it averages over the station-years where the call is easy. A criterion-exceedance judgement is the right place, because it is made one station-year at a time and it is where a wrong peak becomes a wrong decision. On reserved networks the misclassification rate of a 20 °C exceedance is 6.86% when the statistic is computed on recorded days, 1.97% when every year is reconstructed and not screened, 1.53% under length screening and 0.36% under the screen. Restricted to the years where the call is close the four figures are 16.84%, 9.50%, 7.26% and 1.92% within 1.0 °C of the threshold and 19.10%, 13.12%, 10.24% and 3.38% within 0.5 °C, so the advantage is proportionally larger exactly where it is needed. The bias in the count of years above 20 °C falls from −0.76 years per station under recorded days and −0.32 under length screening to +0.06.

That result is not an artefact of the threshold chosen. Swept from 14 to 24 °C the screen's misclassification stays between 0.18% and 0.37% while the length rule stays between 0.84% and 1.53%. The sweep matters because no single number is the criterion: regulatory practice uses a seven-day average of daily maxima with criteria differing by several degrees between jurisdictions and designated uses, 16 °C for core cold-water salmonid habitat and 13 °C for salmonid spawning (Washington State Department of Ecology, 2024), which is why the seven-day endpoint is reported throughout and why the threshold is swept rather than asserted.

For a judgement of this kind, screening is not the best use of the calibrated object. Discarding a station-year is a blunt response to uncertainty about a single call: screening at the 0.5 °C tolerance judges 67.5% of years at 0.36% misclassification, whereas abstaining only where the interval straddles the threshold judges 92.7% at 0.42%, and doing both judges 65.7% at 0.08%. Keeping 93% of station-years rather than 68% at essentially the same error rate matters to anyone assembling a regional archive. The two actions answer different questions — screening decides what enters an archive, abstention decides what one judgement may say — and the same calibrated object serves both, which is an argument for calibrating rather than for screening.

5.6. What an agency should change, and in what order

The first recommendation is to treat screening as estimand-specific. An agency assembling warm-season extremes from maximum gap length should trial conditioning on how much of a station's climatological warm window the gap hides, and should keep length-based rules for annual means, cold extremes and the sine-fit regime parameters for which standing guidance was derived. On the retrospective reserved holdout, substituting a support-aware screen takes the consequential-error rate from 9.7% to 2.5% at equal retention on pooled designed outages; the empirical-arm gain is not resolved, so this is not yet a claim about the networks' natural failure calendar.

Maintenance-calendar pricing that previously appeared to show autumn servicing gains of several percentage points is withdrawn from recommendations until the cross-year outage placement is recomputed: the earlier counterfactual truncated late-year gaps at day 366 while still labelling them by their nominal length. Until that recompute and a never-used-network confirmation exist, schedule changes remain a hypothesis suggested by autumn-heavy long outages, not a verified error-rate benefit. Redundancy investments simi-

This paper has not been peer-reviewed

larly need cost and failure-mode evidence beyond synchrony correlations on the opened panel.

Thresholds should be applied within strata when difficulty correlates with elevation or donor poverty, or the archive can quietly lose mountain stations. Nothing here applies to ungauged sites, because locating support uses a station's own history. A pooled conformal interval is a statement about a population of networks, not an automatic guarantee at a single new site; cluster-aware calibration must be chosen on calibration data only and validated on never-used networks before it is sold as site-level coverage.

6. Conclusions

Whether a data gap matters depends on how much of the target statistic's support it hides. That quantity is a useful risk descriptor and screening axis for archive decisions; we do not claim an unconditional proportion law relating hidden support to loss for every residual structure and every competing peak window. A frozen screen, scored as a retrospective network holdout on 36 held-out networks, reduces consequential error for the hottest-30 mean from 9.7% to 2.5% at 70% retention on pooled designed outages. Continuum, crossover and dilation results that suggest when timing versus length should govern were developed after that panel was opened and remain exploratory until confirmed on never-used networks.

Three conditions bound what can be said now. On the sine-fit estimands standing guidance was derived for, transferring a warm-season threshold is the mistake to avoid rather than the guidance itself. Operational gain under the agencies' own outage geometry is not established on the empirical arm. Difficulty-based screening can thin high, steep, donor-poor catchments unless thresholds are applied within strata. Maintenance-calendar benefits and "annual peak never observed" counts are not retained as verified effects: the former require a corrected multi-year outage geometry, and the latter currently flag wholly masked *climatological* windows.

What follows for practice is correspondingly limited. Prefer estimand-specific rules; keep length-based completeness for averages and sine-fit regime parameters while trialling support-aware rules for warm-season extremes; report retention by catchment attribute alongside error rates; deploy only prediction interfaces that do not require true errors at decision time; and treat autumn maintenance and regulatory hottest-7 claims as unvalidated until recomputed on the correct endpoints and geometries.

This paper has not been peer-reviewed

Open Research Statement

Daily water-temperature records are from nine public providers (ARSO, CHMI, FOEN, GKD Bayern, LUBW, NVE, Rijkswaterstaat, SYKE and the USGS); all are open archives and none required a data-use agreement. Station coordinates are from those providers' own catalogues, the USGS NWIS site service, and ARSO's public station feed. Daily 2 m air temperature is NASA POWER (MERRA-2) (NASA Langley Research Center, 2026). Basin characteristics are GAGES-II (Falcone, 2011).

All analysis code, derived artifacts, figures, the editable draw.io sources of the schematic figures and the frozen pre-registration are in the accompanying repository. A claim registry (`results/wrr_submission_ready`) records each headline claim's artifact, endpoint, unit, denominator, evidence type and whether it may enter the abstract or conclusions. An automated audit checks a declared set of archived manuscript numbers against artifacts; it is not yet a full semantic binding of every live TEX numeral. Input-file hashes for the 146 study networks are in `results/wrr_submission_ready/current_data_manifest.csv`. A permanent archive of the repository and a DOI will be deposited on acceptance.

Acknowledgments

Acknowledgments and funding statements pending author confirmation; see `paper_archived/author_infor` in the repository. The authors declare no competing interests.

Acknowledgements:
Your acknowledgements will appear here.

References

Beaufort, A., Diamond, J. S., Sauquet, E., & Moatar, F. (2022). Spatial extrapolation of stream thermal peaks using heterogeneous time series at a national scale. *Hydrology and Earth System Sciences*, *26*(13), 3477–3495. doi: 10.5194/hess-26-3477-2022

Caissie, D. (2006). The thermal regime of rivers: a review. *Freshwater Biology*, *51*(8), 1389–1406. doi: 10.1111/j.1365-2427.2006.01597.x

Corona, C. R., & Hogue, T. S. (2025). Machine learning in stream and river water temperature modeling: a review and metrics for evaluation. *Hydrology and Earth System Sciences*, *29*(12), 2521–2549. doi: 10.5194/hess-29-2521-2025

Falcone, J. A. (2011). *GAGES-II: Geospatial attributes of gages for evaluating streamflow* (Tech. Rep.). U.S. Geological Survey. doi: 10.3133/70046617

Geifman, Y., & El-Yaniv, R. (2017). Selective classification for deep neural networks. In *Advances in neural information processing systems* (Vol. 30).

Grey, V., Hatt, B. E., Fletcher, T. D., Smith-Miles, K., & Coleman, R. A. (2024). Analysis of irregularly sampled stream temperature time series: challenges and solutions. *Journal of Hydrology*, *636*, 131244. doi: 10.1016/j.jhydrol.2024.131244

Hampel, F. R. (1974). The influence curve and its role in robust estimation. *Journal of the American Statistical Association*, *69*(346), 383–393. doi: 10.1080/01621459.1974.10482962

Huang, F., Qian, B., & Ochoa, C. G. (2023). Long-term river water temperature reconstruction and investigation: a case study of the Dongting Lake Basin, China. *Journal of Hydrology*, *616*, 128857. doi: 10.1016/j.jhydrol.2022.128857

Isaak, D. J., Wenger, S. J., Peterson, E. E., Ver Hoef, J. M., Nagel, D. E., Luce, C. H., … Payne, S. L. (2017). The NorWeST summer stream temperature model and scenarios for the western U.S.: a crowd-sourced database and new geospatial tools foster a user community and predict broad climate warming of rivers and streams. *Water Resources Research*, *53*(11), 9181–9205. doi: 10.1002/2017WR020969

Johnson, Z. C., Johnson, B. G., Briggs, M. A., Snyder, C. D., Hitt, N. P., & Devine, W. D. (2021). Heed the data gap: guidelines for using incomplete datasets in annual stream temperature analyses. *Ecological Indicators*, *122*, 107229. doi: 10.1016/j.ecolind.2020.107229

Kratzert, F., Nearing, G., Addor, N., Erickson, T., Gauch, M., Gilon, O., … Matias, Y. (2023). Caravan — a global community dataset for large-sample hydrology. *Scientific Data*, *10*, 61. doi: 10.1038/s41597-023-01975-w

Lei, J., G'Sell, M., Rinaldo, A., Tibshirani, R. J., & Wasserman, L. (2018). Distribution-free predictive inference for regression. *Journal of the American Statistical Association*, *113*(523), 1094–1111. doi: 10.1080/01621459.2017.1307116

Li, H., Deng, X., & Smith, E. (2017). Missing data imputation for paired stream and air temperature sensor data. *Environmetrics*, *28*(2), e2426. doi: 10.1002/env.2426

Mauger, S., Shaftel, R., Leppi, J. C., & Rinella, D. J. (2017). Summer temperature regimes in southcentral Alaska streams: watershed drivers of variation and potential implications for Pacific salmon. *Canadian Journal of Fisheries and Aquatic Sciences*, *74*(5), 702–715. doi: 10.1139/cjfas-2016-0076

Mauger, S., Shaftel, R., Trammell, E. J., Geist, M., & Bogan, D. (2015). Stream temperature data collection standards for Alaska: minimum standards to generate data useful for regional-scale analyses. *Journal of Hydrology: Regional Studies*, *4*, 431–438. doi: 10.1016/j.ejrh.2015.07.008

Mohseni, O., Stefan, H. G., & Erickson, T. R. (1998). A nonlinear regression model for weekly stream temperatures. *Water Resources Research*, *34*(10), 2685–2692. doi: 10.1029/98WR01877

NASA Langley Research Center. (2026). *POWER daily data, T2M (MERRA-2).* Prediction Of Worldwide Energy Resources project. (`https://power.larc.nasa.gov`)

Rubin, D. B. (1976). Inference and missing data. *Biometrika*, *63*(3), 581–592. doi: 10.1093/biomet/63.3.581

Washington State Department of Ecology. (2024). *Water quality standards for surface waters of the state of Washington, WAC 173-201A-020 and WAC 173-201A-200.* Washington Administrative Code.

Zaoui, A., Denis, C., & Hebiri, M. (2020). Regression with reject option and application to kNN. In *Advances in neural information processing systems* (Vol. 33).

This paper has not been peer-reviewed