Supporting Information for: Discard or reconstruct? A leakage-safe benchmark of gap handling for annual heat extremes and their trends in river-temperature archives

This Supporting Information contains eleven tables that document the station roster and trend signal, the archive-level direction counts, the scope of the uncertainty summaries, the H7 refit robustness check, the open-source and flow diagnostics, the definitions, the software release, and the data-access terms. All values are taken from saved result files; no new model fit or data download was performed for these tables.

S1. Station, group, and reference-year summary

Table S1 summarizes the 41 forward stations by region and connected group. The cohort contains 24 Bavarian stations in eight groups, 10 Swiss stations in two groups, and seven USGS stations in seven groups. The median reference-year set contains 14 complete years (minimum 7, maximum 16), and 24 stations have at least one internal gap year inside their reference span.

S2. Full station-level trend metrics

Table S2 reports the complete-record H30 trend metrics for all 41 forward stations. The OLS slope and its classical standard error are computed on each station's saved reference-year set. The Sen slope is the median of pairwise slopes. The Mann–Kendall p-value includes a tie correction in the variance of S and a continuity correction, but no autocorrelation correction. The H30 interannual SD describes the year-to-year variability of the complete-record annual maximum.

S3. Archive-level direction counts

Table S3 gives the within-archive direction of the all-year comparison for both gap families and for the same-count CDF and S_{xx} calendar controls.

Table S1: Station, connected-group, and reference-year summary by region. Complete-record years are the years in each station's saved reference-year set.

Region	Stations	Groups	Median complete-record years (range)	Median first–last year	Median span (years)	Stations with internal gaps
FOEN (Switzerland)	10	2	16 (15–16)	2010–2025	16	1
GKD Bayern (Bavaria)	24	8	13 (7–16)	2010–2025	15	18
USGS HUC03	1	1	14 (14–14)	2011–2025	15	1
USGS HUC04	1	1	11 (11–11)	2010–2024	15	1
USGS HUC10	2	2	14 (12–16)	2010–2025	16	1
USGS HUC17	3	3	12 (10–14)	2010–2024	14	2
All forward stations	41	17	14 (7–16)	2010–2025	15	24

Notes: A station-year is one station and one calendar year. A connected group links stations by river network or record family and is the cluster unit. The reference-year set contains the complete years available for a station under the design and need not be contiguous. Span is last year minus first year plus one. Internal gaps count stations with at least one missing year inside the reference span. Source: `results/wrr_final_revision_20261031/reference_year_sets.csv` and `results/wrr_final_revision_20261031/retention_choice_counts.csv`.

The all-year rows are the primary comparison. Numerical equality uses an arithmetic 10^{-12} °C per decade tolerance and is not practical equivalence. Archives are station/mask combinations, not independent rivers. The uniform-exact calendar control is not tabulated because its archive-level estimate errors are not saved; its cohort RMSE is 0.449 (controlled) and 0.135 (calendar) °C/10a, with 55 and 8 sign flips.

S4. Scope of the uncertainty summaries

Table S4 states what each of the four uncertainty summaries describes and what it does not establish. The four summaries are annual H30 90% intervals, finite-record slope 95% intervals, paired RMSE-difference 95% group intervals, and three-source-seed or random-path ranges. None is a general coverage guarantee or a climate-process confidence interval.

S5. H7 refit robustness

Table S5 reports the H7 refit. The H7 target is the annual maximum of seven consecutive daily means, not the H30 target used in the main analysis. The refit used 20 seeds and 540 outer-region fits. Panel A gives the danger-scope MAE, CRPS, tail CRPS, and 90% coverage for the four estimators. Panel B gives the direction-stability comparison of D2 relative

Table S2: Full station-level H30 trend metrics for the 41 forward stations. OLS slope, OLS SE, and Sen slope are in °C/10a; H30 interannual SD is in °C. OLS SE is classical and excludes autocorrelation. Mann–Kendall p includes a tie correction in $\mathrm{Var}(S)$ but no autocorrelation correction.

Station	Region	Group	n	OLS slope	OLS SE	OLS p	Sen slope	MK p	H30 SD
foen|2016	FOEN	g0029	16	0.850	0.644	0.208	0.863	0.065	1.217
foen|2019	FOEN	g0029	16	0.496	0.180	0.015	0.557	0.027	0.398
foen|2029	FOEN	g0029	16	0.757	0.645	0.260	0.816	0.079	1.204
foen|2030	FOEN	g0029	16	0.960	0.543	0.099	0.633	0.043	1.070
foen|2085	FOEN	g0029	16	0.349	0.619	0.582	0.331	0.499	1.115
foen|2113	FOEN	g0029	15	0.780	0.790	0.342	0.728	0.138	1.385
foen|2126	FOEN	g0030	16	0.423	0.541	0.447	0.261	0.893	0.984
foen|2135	FOEN	g0029	16	0.841	0.549	0.148	0.723	0.115	1.057
foen|2205	FOEN	g0029	16	0.906	0.648	0.184	0.972	0.053	1.232
foen|2457	FOEN	g0029	16	1.248	0.279	<0.001	1.356	<0.001	0.774
gkd_bayern|10026293	GKD	g0032	13	0.823	0.712	0.273	0.554	0.246	1.148
gkd_bayern|10039802	GKD	g0032	12	0.712	0.808	0.399	0.650	0.304	1.199
gkd_bayern|10046105	GKD	g0032	15	1.235	0.696	0.099	1.261	0.113	1.251
gkd_bayern|10056302	GKD	g0032	16	0.643	0.661	0.348	0.513	0.344	1.217
gkd_bayern|10074009	GKD	g0032	13	0.465	0.908	0.619	1.097	0.360	1.326
gkd_bayern|10081004	GKD	g0032	14	0.745	0.940	0.443	0.585	0.443	1.397
gkd_bayern|10089006	GKD	g0032	15	0.568	0.853	0.517	0.575	0.621	1.398
gkd_bayern|10091008	GKD	g0032	14	1.081	1.037	0.318	1.003	0.274	1.710
gkd_bayern|10098007	GKD	g0032	14	−0.411	1.066	0.706	−0.767	0.511	1.555
gkd_bayern|11401009	GKD	g0034	12	0.539	0.405	0.212	0.718	0.304	0.656
gkd_bayern|11402001	GKD	g0034	14	1.204	0.688	0.106	1.642	0.101	1.239
gkd_bayern|11405000	GKD	g0034	7	0.544	0.508	0.333	0.233	0.368	0.689
gkd_bayern|16003003	GKD	g0036	13	0.867	0.453	0.082	0.813	0.127	0.818
gkd_bayern|16005701	GKD	g0036	9	1.542	0.953	0.150	1.581	0.251	1.433
gkd_bayern|16007004	GKD	g0036	16	0.819	0.450	0.090	0.860	0.137	0.891
gkd_bayern|17207508	GKD	g0039	9	0.930	0.529	0.122	0.549	0.348	0.706
gkd_bayern|18001508	GKD	g0035	8	0.568	0.673	0.431	0.669	0.266	0.509
gkd_bayern|18008008	GKD	g0035	10	1.037	0.775	0.218	1.004	0.107	1.091
gkd_bayern|18403002	GKD	g0031	14	1.078	0.584	0.090	1.167	0.101	1.011
gkd_bayern|24010004	GKD	g0037	12	0.757	0.590	0.229	0.494	0.304	1.032
gkd_bayern|24022003	GKD	g0037	10	0.564	0.603	0.377	0.407	0.721	0.690
gkd_bayern|24042000	GKD	g0037	12	−0.026	0.584	0.965	−0.061	1.000	0.791
gkd_bayern|24064003	GKD	g0037	14	0.754	0.507	0.162	0.743	0.228	0.946
gkd_bayern|24405002	GKD	g0033	9	0.795	0.653	0.263	0.883	0.348	0.952
usgs|02172020	HUC03	g0181	14	0.398	0.270	0.166	0.467	0.125	0.463
usgs|04137030	HUC04	g0166	11	−0.609	0.317	0.086	−0.750	0.087	0.529
usgs|06036905	HUC10	g0053	16	0.701	0.415	0.113	0.438	0.224	0.811
usgs|06054500	HUC10	g0143	12	0.300	0.531	0.584	−0.243	0.945	0.791
usgs|13342500	HUC17	g0054	12	0.273	0.439	0.548	0.000	1.000	0.691
usgs|14166000	HUC17	g0057	14	0.426	0.487	0.399	0.467	0.381	0.727
usgs|14211500	HUC17	g0059	10	1.776	0.468	0.005	1.638	0.004	0.874

Notes: n is the number of reference years. p-values below 0.001 are printed as <0.001. Values are rounded for display; the saved file retains full precision. Source: `results/ems_revision_20261102/trend_signal.csv`. Region labels group the USGS stations into HUC03, HUC04, HUC10, and HUC17. Station keys retain the provider prefix.

Table S3: Archive-level direction of the all-year slope-error comparison. Counts give archives in which each scheme's absolute finite-record slope error is smaller than, larger than, or numerically equal to warm-season screening; the all-year rows are the primary comparison and the CDF and S_{xx} rows use the same-count controls. Median change is the within-archive absolute-error change in °C per decade. Numerical equality uses an arithmetic 10^{-12} °C per decade tolerance, which is not practical equivalence. The controlled 30–90-day gap and empirical-calendar families are separate cohorts, and archives are station/mask combinations rather than independent rivers; a changed year set means that the selected years differ from those retained under warm-season screening.

Family	Scheme	Subset	n	Smaller abs. error	Larger abs. error	Numerically equal	Median change
Controlled 30–90 day	All years	All archives	805	630	169	6	−0.1395
Controlled 30–90 day	All years	Changed year set	800	630	169	1	−0.1400
Controlled 30–90 day	CDF	All archives	805	197	170	438	0.0000
Controlled 30–90 day	CDF	Changed year set	367	197	170	0	−0.0196
Controlled 30–90 day	S_{xx}	All archives	805	218	158	429	0.0000
Controlled 30–90 day	S_{xx}	Changed year set	376	218	158	0	−0.0421
Empirical calendar	All years	All archives	818	231	57	530	0.0000
Empirical calendar	All years	Changed year set	288	231	57	0	−0.0578
Empirical calendar	CDF	All archives	818	144	91	583	0.0000
Empirical calendar	CDF	Changed year set	235	144	91	0	−0.0256
Empirical calendar	S_{xx}	All archives	818	151	83	584	0.0000
Empirical calendar	S_{xx}	Changed year set	234	151	83	0	−0.0490

Notes: "Smaller abs. error" and "Larger abs. error" count archives relative to warm-season screening at the same station and mask. The all-year comparison is the primary row; the CDF and S_{xx} rows are same-count calendar controls. Median change is the median within-archive change in absolute slope error, not a cohort RMSE. The `uniform_exact` scheme is not shown as a row because its archive-level estimate errors are not saved for this cohort; its saved cohort RMSE is 0.449 (controlled) and 0.135 (calendar) °C/10a, with 55 and 8 sign flips. Source: `results/wrr_comprehensive_review_20261102/archive_level_comparisons.csv`; the uniform-exact cohort values are from `results/ems_revision_20261102/trend_signal.csv`.

Table S4: Scope of four uncertainty summaries. Annual H30 90% intervals describe the annual predictive distribution under fixed methods and information. Finite-record slope 95% intervals propagate that annual distribution and the specified joint construction to the slope. Paired RMSE-difference 95% group intervals describe comparison variability conditional on the saved annual predictions, the predefined connected groups and the rules compared. Three-source-seed or random-path ranges describe numerical simulation variation at a fixed budget. None establishes a general coverage guarantee, a climate-process confidence interval, or uncertainty from retraining source models and selecting the rule. At 4,096 samples, the nominal 95% trend intervals remain 87.4–89.9%, below nominal 95%, as a saved diagnostic rather than a new experiment.

Summary	What it describes	What it does not establish
Annual H30 90% intervals	Annual predictive distribution under fixed methods and information; empirical coverage of the annual target	Not evidence that every region or reconstructor attains 90%; endpoint comparisons use an arithmetic tolerance.
Finite-record slope 95% intervals	Propagates the annual distribution and the specified joint construction to the slope on the reference-year set	Not a 95% confidence guarantee for a climate-process true trend, another reference-year set or a complete calendar.
Paired RMSE-difference 95% group intervals	Comparison variability conditional on saved annual predictions, predefined connected groups and the rules compared	Not retraining or rule-selection uncertainty, multiplicity across sensitivities, or new climate samples.
Three-source-seed / random-path ranges	Numerical or simulation variation at a stated budget with fixed data and rules	Not a confidence interval across independent hydrological samples; more paths do not establish convergence.

Notes: The 87.4–89.9% range gives the minimum lower and maximum upper bounds of the 4,096-draw coverage intervals across the saved methods and constructions, not point estimates; the corresponding point estimates span 87.5–89.7%. It is a saved diagnostic for the nominal 95% trend intervals, not a new experiment or a general coverage guarantee. Source: `results/wrr_adversarial_review_20261027/rank_convergence/rank_convergence_summary.csv`.

to calibrated stochastic regression: the median paired MAE reduction is 2.94% in the crossed family and 4.85% in the calendar family, and CRPS and tail CRPS improve in 20 of 20 seeds in both families. Panel C gives trend-target missingness-component coverage: D2 multiple imputation has median coverage 81.2% under MCAR and 80.7% under the strong-heat scenario, but 4.5% under the non-air-trend scenario. The near-81% value therefore describes the MCAR and strong-heat trend scenarios, not the non-air-trend scenario.

S6. Open-source candidate-provider sensitivity

Table S6 reports the candidate-provider sensitivity. Replacing the restricted source providers with seven candidate providers changes the crossed/forward D2 trend RMSE from 0.164 to 0.158 °C/10a and the classical trend RMSE from 0.147 to 0.156 °C/10a over 738 four-way common pairs in 17 groups. D2 retention falls from 93.9% to 93.7% on 807 factor cases, and nominal 95% D2 interval coverage falls from 88.0% to 87.6% on 820 coverage cases. The ranking of the two approaches is unchanged. This is a candidate open reproducibility route, not a legal clearance.

S7. Zero-flow flow-proxy logistic results

Table S7 reports the station-fixed conditional-logit associations between a warm-season-incomplete station-year and within-station standardized summer air temperature, within-station standardized summer log flow, and a linear trend. The flow-adjusted odds ratios are 0.594 [0.409, 0.813] for the recorded-zero cohort and 0.474 [0.265, 0.676] for the at-least-30-zero-day cohort. The air-temperature odds ratios are 0.993 [0.659, 1.305] and 0.855 [0.580, 1.275]. These are associations in a historical proxy cohort, not causal effects or a missing-at-random test.

S8. Flow-comparator summary

Table S8 reports availability and mean absolute error for the opened flow-comparator methods. Historical D2 has full availability and the lowest MAE in both scopes. Donor-dependent methods are available on 81.7% of crossed/all eligible cases and do not lower paired MAE. The paired comparisons use 66,099 common cases in the all scope and 33,609 in the danger scope, covering

Table S5: H7 refit robustness. Panel A reports native H7 source refits over 20 seeds and 540 outer-region fits; seven-day peak supports, masks, history summaries, conditional functionals, MOVE augmentation, annual models, and disjoint calibration were recomputed. Panel B reports the direction-stability check for D2 (distribution regression) relative to calibrated stochastic regression. Panel C reports trend-target coverage of the missingness component for the multiple-imputation pipelines. These are H7 results, not the historical H30 endpoint.

Panel A: H7 danger-scope refit scores (20 seeds).

Family	Estimator	Seeds	MAE (°C)	CRPS (°C)	Tail CRPS (°C)	Coverage 90%
Calendar	Distribution regression (D2)	20	0.494	0.365	0.065	0.874
Calendar	Functional debiasing	20	0.522	–	–	–
Calendar	Daily plug-in	20	0.638	–	–	–
Calendar	Calibrated stochastic	20	0.516	0.380	0.068	0.858
Crossed	Distribution regression (D2)	20	0.348	0.257	0.046	0.892
Crossed	Functional debiasing	20	0.375	–	–	–
Crossed	Daily plug-in	20	0.423	–	–	–
Crossed	Calibrated stochastic	20	0.360	0.265	0.048	0.893

Panel B: D2 relative to calibrated stochastic regression.

Family	Comparison	Median paired MAE reduction (%)	CRPS gain positive seeds	Tail CRPS gain positive seeds
Crossed	D2 vs calibrated stochastic	2.94	20/20	20/20
Calendar	D2 vs calibrated stochastic	4.85	20/20	20/20

Panel C: Trend-target missingness-component coverage (median, seed range).

Scenario	D2 multiple imputation	Calibrated stochastic MI	Stochastic regression MI
MCAR	81.2% (79.1–86.2)	78.4% (75.7–79.7)	75.6% (73.9–76.7)
Strong heat	80.7% (78.9–87.4)	78.5% (75.5–81.3)	76.0% (73.8–78.4)
Non-air trend	4.5% (3.8–5.6)	3.9% (2.9–4.5)	3.7% (2.7–4.2)

Notes: Panel A values are medians over 20 seeds in the dangerous decision-time stratum. Coverage is the empirical coverage of nominal 90% annual H7 intervals. Panel B "median paired MAE reduction" is the median over saved station-level paired relative reductions in MAE for D2 relative to the calibrated stochastic comparator across the 20 seeds; positive CRPS and tail-CRPS gains mean lower D2 scores. Panel C covers the trend target, not the annual H7 target; values are medians with minimum–maximum seed ranges. H7 is a different target and is not used as confirmation of the H30 results. Source: `results/wrr_review_completion_20261024/evidence/H7_refit_scorecard_summary.csv`, `results/wrr_review_completion_20261024/evidence/H7_refit_direction_stability.csv`, `results/wrr_review_completion_20261024/target_windows/h7/full/trends_per_seed.csv`, and `results/wrr_review_completion_20261024/target_windows/h7/full/completion.json`.

Table S6: Open-source candidate-provider sensitivity on the crossed/forward cohort. The candidate route refits the source models on seven candidate providers with recorded reuse paths while retaining the target masks, daily features, and plug-in baselines. Retention and coverage are D2 quantities.

Metric	Original source	Candidate source
D2 trend RMSE (°C/10a)	0.164	0.158
Classical trend RMSE (°C/10a)	0.147	0.156
D2 retention in the symmetric MSE decomposition (%)	93.9	93.7
Nominal 95% D2 interval coverage (%)	88.0	87.6
Common four-way pairs	738	738
Connected groups	17	17
Factor cases for retention	807	807
Coverage cases	820	820

Notes: The candidate route leaves the ranking of D2 and the classical comparator unchanged. It is a candidate open reproducibility route, not a legal clearance: LUBW rights and the provenance of neighbor records still require confirmation. Candidate-source tables are descriptive, repetitions are not independent climates, and the filtered inputs remain processed study data rather than provider-native records. Source: `results/ems_revision_20261102/flow_diagnostic.json`, extracted from `paper/wrr_tr/open_source_values.tex`.

210 sites. The full cohort contains 255 sites, 8,999 complete climatic years, 134,925 cases, and 17 HUC2 region groups.

S9. Definitions and symbols

Table S9 collects the terms and symbols used in the main text and in this Supporting Information. The decomposition $E = S + R$ is an arithmetic decomposition of the finite-record slope error, not a causal variance decomposition. The observed lower bound L_y is a property of the reported series, not an annual estimate.

S10. Software and reproducibility

Table S10 summarizes the `gapbench` 1.2.0 release. The package exposes six public functions, passes 13 tests, includes a synthetic quickstart, and provides a Docker smoke environment that imports LightGBM and runs the quickstart without restricted observations. The version is consistent across the package metadata and citation file. An archival DOI and a v1.2.0 release

Table S7: Zero-flow flow-proxy conditional-logit results. The response is a warm-season-incomplete station-year (the hidden-peak condition). Air and flow covariates are within-station standardized summer means; the model also includes a linear trend per decade. Odds ratios are shown with 95% confidence intervals.

Cohort	n	Events	Stations	Groups	z_{air} OR (95% CI)	z_{flow} OR (95% CI)
Recorded zero flow	282	129	21	17	0.993 (0.659–1.305)	0.594 (0.409–0.813)
At least 30 zero days	215	104	16	14	0.855 (0.580–1.275)	0.474 (0.265–0.676)
Fewer than 30 zero days	4,417	1,814	271	103	1.024 (0.944–1.122)	0.969 (0.897–1.046)
No zero in available flow	4,350	1,789	266	103	1.020 (0.937–1.125)	0.968 (0.895–1.050)

Notes: The underlying USGS missingness sample contains 6,479 station-years at 336 physical stations; 21 stations in 17 groups recorded at least one zero-flow day and 16 stations in 14 groups recorded at least 30. Confidence intervals use 400 bootstrap draws. Lower hidden-peak odds is an association in this proxy cohort, not evidence that low flow protects against underestimation. Zero flow is a historical proxy; no recorded zero does not prove perennial flow, and a zero does not identify an intermittent river, sensor exposure, or a dry bed. Source: `results/wrr_review_completion_20261024/zero_flow_missingness/conditional_logit_summary.csv` and `results/wrr_review_completion_20261024/zero_flow_missingness/cohort_counts.csv`.

tag are pending. The D2 model and the conditional AR(1) bridge remain in the research compendium.

S11. Data and access summary

Table S11 summarizes the data providers and access terms. FOEN, GKD Bayern, USGS NWIS, and NASA POWER are public sources with provider-specific attribution requirements. LUBW observations are restricted; reviewer access remains pending provider agreement and editorial/legal confirmation. The MIT license applies to the `gapbench` code and does not relicense observational data. Processed annual tables are distributed in the public repository; restricted daily observations are not. Archival DOIs for the code release and processed tables are pending.

S12. Wild cluster bootstrap sensitivity

Table S12 adds wild cluster bootstrap intervals for the paired RMSE differences of the main text. The wild procedures reweight the group-level influence function of the RMSE difference with Rademacher or Webb six-point signs and never resample the 17 connected groups, so they probe the same comparisons without the few-cluster resampling approximation. The

median wild-to-group width ratio is 1.02. Directions are unchanged where both procedures agree: the calendar CDF interval includes zero under both procedures, the calendar S_{xx} and all-year intervals exclude zero under both, and the controlled-gap D2−classic interval excludes zero under both. One comparison is method-dependent: the calendar D2−classic interval includes zero under group resampling but excludes zero under both wild procedures, so the common-case classical advantage on the calendar family is not promoted to a stable ranking. These intervals describe comparison variability conditional on the frozen per-archive tables; they are not new climate samples or a coverage guarantee.

Table S8: Opened flow-comparator scorecard for the crossed family. Availability is the fraction of eligible cases with a valid estimate; MAE is mean absolute error in log-flow units computed on each estimator's valid cases. The "all" scope contains 80,931 eligible cases and the "danger" scope 41,132; paired log-MAE comparisons use 66,099 and 33,609 common cases, respectively.

Estimator	Availability, all	MAE, all	Availability, danger	MAE, danger
Historical D2	1.000	0.0500	1.000	0.0743
Historical multidonor Ridge	1.000	0.0564	1.000	0.0829
Single-donor Ridge	0.817	0.0602	0.817	0.0896
Observed days	1.000	0.0646	1.000	0.1004
Stochastic AR(1) median	0.817	0.0716	0.817	0.1038
MOVE.1	0.817	0.0734	0.817	0.1062

Notes: The tabulated MAE is log-MAE (mean absolute error in log-flow units) on each estimator's valid cases: all eligible cases for historical D2, historical multidonor Ridge, and observed days; the donor-available subset for single-donor Ridge, stochastic AR(1), and MOVE.1 (availability 0.817, i.e., 66,099 of 80,931 eligible cases in the all scope and 33,609 of 41,132 in the danger scope). Paired log-MAE comparisons are restricted to these common cases and cover 210 sites; the main-text paired basis yields D2 MAE 0.0495 (all) and 0.0745 (danger), slightly different from the per-estimator values tabulated here. The full cohort contains 255 sites, 8,999 complete climatic years, 134,925 cases, and 17 HUC2 region groups. The 1.000 availability rows are historical D2, historical multidonor Ridge, and observed days; the donor-dependent rows do not lower paired MAE. The experiment is an opened positive-flow perennial subset, not a new blind transfer or a 20-seed stability claim. Paired differences are stored as left and right MAE on common cases; no derived subtraction is saved. Source: `results/wrr_review_completion_20261024/flow_comparators/scorecard.csv` and `results/wrr_review_completion_20261024/flow_comparators/paired_effects.csv`; cohort counts are from `results/wrr_review_completion_20261024/flow_comparators/status.json` and `results/wrr_review_completion_20261024/flow_comparators/design.json`.

Table S9: Definitions and symbols used in the main text and this Supporting Information.

Term or symbol	Definition
Station-year	One station and one calendar year.
Archive	One station × one source repetition × one gap setting or empirical calendar. The controlled-gap cohort contains 805 analyzed archives; the empirical-calendar cohort contains 818.
Connected group	A set of stations linked by the same river network or record family. It is the cluster unit for group-disjoint calibration and resampling.
Reference-year set Y_i	The complete years available for station i under the evaluation design. It need not be contiguous, and years outside it are not imputed as part of the target. The target is the finite-record slope on Y_i.
H30	Annual maximum of the 30-day running mean water temperature in year y. It is a sustained-heat summary, not an ecological standard.
H7	Annual maximum 7-day mean water temperature. It is a different target, used only as a robustness check and not as confirmation of the H30 results.
L_y	Largest fully observed 30-day mean in an incomplete year y. It satisfies $\text{H30}_y \geq L_y$ for the reported series and is an observed lower bound, not an annual estimate.
SE	Classical OLS standard error of the finite-record slope on the reference years. It excludes autocorrelation and is a lower bound on the trend's sampling uncertainty.
S, R, E	With β_F the complete-record slope on Y_i, β_S the complete-record slope on the selected year set, and β_{SE} the estimated slope on that set: $S = \beta_S - \beta_F$, $R = \beta_{SE} - \beta_S$, and $E = \beta_{SE} - \beta_F = S + R$. The squared-error identity includes the cross term 2 mean(SR); the terms are an arithmetic decomposition, not causal variance shares.
Controlled gap	A contiguous 30-, 60-, or 90-day mask applied jointly to every year of an archive, with at least half overlap of the training-period climatological peak. This is a severe stress, not an estimate of outage frequency.
Empirical calendar	A year-specific gap calendar copied from the observed archive. It preserves each year's gap length and seasonal position but not the exact missing days, so it is closer to observed annual longest-gap conditions than the controlled stress.
Warm-season rule	A station-specific 90% coverage rule requiring coverage in the five consecutive warm months. The 95% annual-coverage rule is a secondary comparator.

Notes: Definitions follow the main text and this Supporting Information; the 805 controlled-gap and 818 empirical-calendar archive counts are from `results/ems_revision_20261102/trend_signal.csv`.

Table S10: Software and reproducibility summary for `gapbench` 1.2.0. The released package contains the reusable primitive layer; fitted D2 and AR(1) bridge models remain in the research compendium.

Item	Status
Package	`gapbench` 1.2.0; Python $\geq$ 3.10; NumPy $\geq$ 1.24; pandas $\geq$ 2.0; MIT license for code.
Public API	Six functions exported from the package root: `observed_window_bound`, `finite_record_slope`, `select_years`, `decompose_slope_error`, `group_splits`, and `mask_archive`.
Tests	13 tests. Clean test command: `pip install -e "gapbench[test]"` followed by `python -m pytest -q gapbench/tests`; all 13 pass.
Quickstart	Synthetic 20-year daily series with four gaps. The example measures the best fully observed 30-day window, selects 18 years under both calendar rules while retaining all 16 complete years, and decomposes the slope error. It runs in under one second.
Docker	Python 3.11 slim image; installs the review and development extras, imports LightGBM, and runs the quickstart. The container contains no restricted observations.
Version consistency	`pyproject.toml`, `CITATION.cff`, and the preferred citation report version 1.2.0; the consistency check reports PASS with a warning that the historical v1.0.0 git tag does not describe version 1.2.0.
Archival DOI	Pending. No Zenodo deposit or v1.2.0 release tag has been created.
Reviewer access	The public repository requires no credentials. Restricted LUBW observations are not redistributed; their access route remains pending provider agreement and editorial/legal confirmation.
Program size and replay memory	24 files, 1,012 lines of Python, 264 KiB uncompressed (23 KiB zipped). Measured on one CPU core: quickstart 0.13 s wall clock and 35 MiB peak resident set; full packaged replay of all 1,623 archive-level masks (8,115 strategy rows) 9.7 s and 289 MiB peak resident set.
Research compendium	The D2 quantile model and the conditional AR(1) bridge are not part of `gapbench`. This separation keeps the published primitives stable while the fitted models depend on restricted data.

Notes: The MIT license applies only to original software and project-authored documentation. It does not relicense observational data or upstream provider products. Source: `gapbench/pyproject.toml`, `CITATION.cff`, `Dockerfile`, and the test suite under `gapbench/tests`.

Table S11: Data and access summary. Provider terms govern access, use, and redistribution. The MIT license applies to code, not to observational data.

Provider	Data	Access and terms	Distribution in this study
FOEN (Switzerland)	Daily water temperature	Public water-observation service; data are published under the opendata.swiss "Open use. Must provide the source." license. Attribution to FOEN/BAFU and a link to the dataset are required.	Processed annual tables only; daily observations are not redistributed.
GKD Bayern (Bavaria)	Daily water temperature	Public official downloads; CC BY 4.0. Attribution: "Datenquelle: Bayerisches Landesamt für Umwelt".	Processed annual tables only; daily observations are not redistributed.
USGS NWIS (United States)	Daily water temperature	Public Water Data APIs. U.S. Government work, generally publishable after acquisition; cite USGS Water Data.	Processed annual tables only; daily observations are not redistributed.
NASA POWER	Daily air temperature	Public point API. U.S. Government work, generally publishable after acquisition and citation.	Processed annual tables only.
LUBW (Baden-Württemberg)	Daily water temperature	Restricted. Dataset-specific UIS terms are identified, but their scope and vintage remain pending; redistribution is not established.	Not redistributed. Reviewer access remains pending provider agreement and editorial/legal confirmation.
`gapbench` code	Software	MIT license for original code and project-authored documentation.	Public repository; no credentials required for the code.
Processed annual tables	Derived annual quantities	No restricted daily observations; derived from provider records under the terms above.	Included in the public repository; archival DOI pending.

Notes: Code availability does not imply redistribution rights for provider data. The restricted daily observations are not part of the public repository or the Docker image. Source: provider terms and the manuscript's software and data-availability statement; no saved numerical file underlies this table.

Table S12: Wild cluster bootstrap intervals for the paired RMSE differences, alongside the saved group-resampling intervals. Clusters are the 17 connected groups and are never resampled by the wild procedure: group contributions of the influence function of the RMSE difference are reweighted with Rademacher (and Webb six-point) signs, 10,000 draws, seed 20261102. Positive estimates mean the first rule or estimator has the larger RMSE. The group-resampling columns reproduce the frozen intervals. Units are °C per decade.

Family	Contrast	n	Estimate	Group bootstrap [95%]	Wild Rademacher [95%]	Wild Webb [95%]
Calendar	Uniform exact − warm-season	818	0.0219	−0.0036, 0.0369	0.0032, 0.0406	0.0031, 0.0405
Calendar	CDF − warm-season	818	−0.0152	−0.0451, 0.0004	−0.0366, 0.0065	−0.0372, 0.0070
Calendar	S_{xx} − warm-season	818	−0.0277	−0.0497, −0.0169	−0.0472, −0.0081	−0.0473, −0.0080
Calendar	All years − warm-season	818	−0.0462	−0.0706, −0.0295	−0.0720, −0.0208	−0.0721, −0.0205
Controlled	Uniform exact − warm-season	805	0.0033	−0.0051, 0.0172	−0.0063, 0.0126	−0.0063, 0.0126
Controlled	CDF − warm-season	805	−0.0175	−0.0274, −0.0052	−0.0292, −0.0060	−0.0294, −0.0059
Controlled	S_{xx} − warm-season	805	−0.0436	−0.0649, −0.0172	−0.0697, −0.0178	−0.0699, −0.0176
Controlled	All years − warm-season	805	−0.2762	−0.3574, −0.1839	−0.3841, −0.1691	−0.3852, −0.1697
Calendar	D2 − classic	701	0.0104	−0.0028, 0.0178	0.0011, 0.0197	0.0011, 0.0197
Controlled	D2 − classic	738	0.0172	0.0036, 0.0270	0.0056, 0.0289	0.0055, 0.0289

Notes: wild intervals are on the influence function of the RMSE difference and need no cluster resampling; the median wild/group width ratio is 1.02 (all-year contrasts 1.24, estimator contrasts 0.90–0.99). The calendar CDF interval includes zero under both procedures; the calendar D2−classic interval includes zero under group resampling but excludes it under the wild procedures, so the common-case classical advantage on the calendar family remains method-dependent and is not promoted to a stable ranking. A wild interval for the paired mean difference of squared errors is also saved. Source: `results/ems_revision_20261102/wild_cluster_bootstrap.csv`, `results/ems_revision_20261102/wild_cluster_bootstrap.json`; group-resampling columns from `results/wrr_final_revision_20261031/retention_paired_intervals.csv` and `results/wrr_final_revision_20261031/estimator_paired_comparisons.csv`.