Incomplete river-temperature years: separating recovery, screening and trend inference

[Author names][1]

[1][Affiliations]

Key points

- Observed windows can disprove a complete-year-only heat record, but most witnesses pass a 90 percent warm-season rule
- Distribution regression and calibrated stochastic reconstruction have similar annual accuracy in difficult gaps
- Retaining years improves the evaluated finite-record trends; forward interval coverage and useful release remain limited

Abstract

Incomplete river-temperature years can contain information about sustained heat, but the evidence needed to recover an annual value, release it, or infer a trend differs. At 20 of 96 long-record stations, fully observed 30-day windows in 35 incomplete years exceed every complete-year heat maximum. Of these witnesses, 32 pass a station-specific 90% warm-season rule, so the observation-only result mainly challenges strict annual completeness. Across twenty source repetitions, annual distribution regression and equally calibrated stochastic reconstruction have similar errors on difficult gaps. We jointly degrade multiyear archives at 42 stations before local refitting and compare retention and estimator effects on the same finite-record slopes. Bounded annual distributions use a stated tail extrapolation and identical marginals across three joint-rank constructions. For controlled gaps, nominal 95% D2 trend intervals cover 94.0% offline and 88.0% in retrospective forward evaluation. Task-level calibration is available for 4.8% of forward controlled-gap cases under empirical support checks, with no population coverage guarantee. A three-output rule retains bounds or withholds values, but its individual classifier score does not certify group risk. Exploratory Canadian comparisons use published daily means or reported-hour records with unresolved quality or time basis; independent strict-reference validation remains open. These findings support preserving observed heat information while restricting claims about numerical recovery, transfer and trend uncertainty.

Plain Language Summary

Gaps in a river's water-temperature record do not necessarily erase information about a hot year. If thirty consecutive observed days are already warmer than the record among fully observed years, the complete-year-only record is incomplete. Most such examples in our archive would also be retained by a less restrictive warm-season rule. We compared several ways to estimate missing values and then tested what they imply for a trend across many years. A calibrated classical reconstruction was about as accurate as a more elaborate annual model. Trend intervals were less reliable when applied to later years, and screening rules did not provide a general safety guarantee. The analysis distinguishes observed facts, estimated values and cases where the evidence is insufficient.

1 Introduction

Stream-temperature records support studies of thermal habitat, climate change and water management (Caissie, 2006; Michel et al., 2020; van Hamel & Brunner, 2024). The annual maximum of a 30-day mean summarizes sustained heat and is related to the thermal peak used in river-network studies (Beaufort et al., 2022). A monitoring gap can affect this statistic very differently depending on its timing, the observations remaining in the year and the information available from other years or stations.

Prior work has already shown that partial stream-temperature records can support some annual analyses. Johnson et al. (2021) evaluated missing intervals and imputation for annual mean, phase and amplitude, finding limitations that varied among rivers. Beaufort et al. (2022) used summer observations and climate correction for a multiyear thermal-peak metric. Neither should be interpreted as requiring every calendar year to be complete. Our question is narrower: what do incomplete daily records support about an annual sustained-heat maximum, numerical publication, and a finite-record trend? We compare strict completeness with a station-specific warm-season rule and distinguish a mathematical observation bound from model-based estimation.

Regression record augmentation and maintenance-of-variance methods have a long hydrological history (Matalas & Jacobs, 1964; Hirsch, 1982; Vogel & Stedinger, 1985). A point prediction, a predictive distribution and a decision to release a value have different targets. Distributional imputation should be assessed for the downstream quantity rather than only for daily squared error (Näf et al., 2023; Gneiting & Raftery, 2007). Here we compare a daily plug-in, simple corrections, calibrated stochastic reconstruction and annual distribution regression. We then jointly hide portions of a station's multiyear record before local refitting and evaluate whether retaining years changes the slope of that finite record. Finally, we examine whether the annual distributions support trend uncertainty and whether visible information can identify cases that should not receive a numerical estimate.

The hypotheses are deliberately conditional: target-oriented recovery can improve on a daily plug-in without necessarily beating a calibrated stochastic comparator; retaining years can improve a finite-record slope under a specified degra-

Table 1. Evidence map. Each claim is paired with the comparison and metric used here and with the boundary that limits it. The table indexes the results; it is not an additional test.

Claim	Primary comparison	Metric	Boundary
Observed windows can overturn a complete-only record	Complete-year H30 vs. fully observed 30-day window L	Count of L above the complete-year maximum	Reported measurements only; not the unique warmest year
Target recovery beats a daily plug-in	D2 vs. plug-in, functional debiasing and calibrated stochastic regression	H30 MAE, CRPS and 90% coverage	No unique gain over calibrated stochastic regression
Retaining years improves finite-record trends	Warm-season rule vs. all supported years at the same estimator	Trend RMSE; symmetric retention and estimator shares	Shares are setting-specific, not a universal percentage
Annual and trend uncertainties need distinct evaluation	Bounded normal-tail vs. constant-tail marginals; three rank couplings	Coverage, width, interval score and availability	Tail, annual and dependence errors are not uniquely separated
Numerical release has a risk–retention trade-off	Fixed score cap vs. three-output rule	Selected risk, retention and group excess	A raw individual score is not a group-risk guarantee
Visible diagnostics do not restore extrapolation safety	Clean vs. hot-year, future-year and air-exposure stress	Selected risk, coverage and retention	Synthetic stress severity; not an archive prevalence

dation mechanism; and annual and trend uncertainty require separate evaluation. The evidence map in Table 1 identifies the comparisons and limits.

2 Data and study design

The source archive contains previously assembled European and US daily observations: 581 physical stations, 185 connected-record groups, 27 regions and 6,170 unique complete station-years. NASA POWER daily air temperature is the common meteorological input (NASA Langley Research Center, 2026). Artificial controlled gaps of 30, 60 or 90 days are crossed with overlap with a training-period climatological peak window; empirical incomplete calendars provide a separate family. Each of twenty source repetitions redraws masks and group-role composition. In the leave-one-year-out source comparison, fitting, interval calibration, score learning and rule calibration use disjoint connected groups outside each held region. These repetitions describe variability within the available archive, not twenty independent climates.

We use complete daily station-years as reference outcomes for artificial masking. Water observations hidden by a mask never enter the local fit for that case.

Features available when a value is evaluated include the visible gap geometry, a training-period climatological peak, local water history, air conditions and available neighboring stations. The true peak location and hidden water values are evaluation quantities. Observation quality is inherited from the assembled source records; a reported value is not assumed to be error-free physical truth. An observed lower bound is valid for the reported series if its thirty measurements are comparable and valid.

The annual target is

$$H30_y = \max_j \frac{1}{30} \sum_{d=j}^{j+29} Y_{yd}. \tag{1}$$

For an incomplete year, the largest fully observed 30-day mean L_y satisfies $H30_y \geq L_y$ for the same reported series. For 1981–2025 we select stations with at least fifteen complete reference years and compare every incomplete-year bound with the largest complete-year H30. We also evaluate whether those same witnesses pass a 95% annual-coverage rule and a 90% station-specific warm-season rule. The five consecutive warm months are obtained from the station's climatology after excluding the evaluated year. These are explicit study comparators; the seven-week missing-period result of Johnson et al. (2021) concerns annual signal parameters, not a validated H30 record rule.

3 Methods

3.1 Annual recovery and decision to release

The daily plug-in computes H30 after a selected daily reconstruction. Functional debiasing corrects its annual value, whereas D2 learns annual-error quantiles using visible features and simulated trajectories. A stochastic regression comparator generates conditional residual bridges and calculates H30 for each realization. D2 and the stochastic comparator have the same source interval-calibration roles and bootstrap multiplicities. Point estimates are medians; MAE, CRPS, 90% interval coverage and interval score are reported on available cases, with paired comparisons on common cases.

In the twenty historical source fits, we apply the same observed-window support projection to both methods after their original calibration. The seven stored quantiles do not determine their outer tails; for comparable CRPS integration we




Figure 1. Evidence architecture of the study. Complete target observations are used only for scoring; every model is fitted on the visible archive defined by a repetition. The lower row names the evaluation objects, and the bottom band lists rules that are held fixed across comparisons. The editable draw.io source is archived with the figure.

extend the tails using adjacent normal-quantile spacings and enforce L_y when available. The original model weights and calibration offsets remain fixed in this source re-analysis. In the whole-archive study described below, offsets for both methods are instead re-estimated on their original independent calibration roles after projection. Neither approach makes the assumed tail shape a verified full distribution.

The conditional-mean mechanism is useful but limited. For observed information X and 30-day averaging vectors w_j, convexity gives

$$\max_j w_j^\top E(Y \mid X) \le E\{\max_j w_j^\top Y \mid X\}.$$

It concerns the true conditional mean and conditional expectation; an approximate learner need not underestimate every individual year. Calibration, local adaptation, model bias and the chosen point summary remain empirical questions.

We evaluate a rule that reports an estimated value, an observed lower bound or an unknown state. Numerical release requires an available D2, at least four

complete historical years, visible features within source-derived limits and a raw individual classifier score no greater than 0.10. The score was trained for the original unprojected D2 and is not a calibrated group-risk certificate for the projected value. A seven-day close match between visible water and air temperature is tested for the imposed exact-air-substitution scenario; that detector is not evaluated for other stresses. No observed bound is released when this quality flag is raised. The primary factorial trend comparison is separate. A second exploratory analysis passes released values and bounds into a Gaussian working likelihood, treating released numerical estimates as exact and each bound L as a right-censored response with contribution $P(H30 \quad \geq \quad L)$. Unknown cases remain in availability denominators. The resulting regression interval includes working-model variability and is not a task-calibrated missingness-only interval.

3.2 Whole-archive degradation and task-level uncertainty

The new trend benchmark hides every selected year of a target station before fitting its daily models, climatology or residual histories. Neighboring physical stations remain observed covariates in this experiment. The benchmark contains 42 stations with at least twenty complete reference years in seven regions. One Finnish record ends in 1994, leaving 41 stations for forward evaluation. Twenty paired mask repetitions use both empirical calendars and controlled 30–90-day gaps; after seven observed burn-in years, each reference year is degraded with probability 0.35. These 3,320 archive experiments reuse no jointly hidden target observation for fitting another year. Original complete records enter scoring only. The archived single-year plug-in is retained as an optimistic-information paired comparator on identical masks.

Offline daily fits use the remaining visible history. Forward annual models and interval offsets use source observations before 2010 and exclude the evaluated test stations and their connected groups. The original partitioned role assignment can retain other training-only groups in the same named region: this occurs in the HUC17 estimator and in NVE interval calibration. Accordingly this forward experiment is held-test-group, not uniformly held-region, and is applied to 2010 onward. Forward local daily fitting and dependence estimation also exclude water observations from 2010 onward. These are current-vintage retrospective forward experiments, not a

historical real-time field trial. Annual regression weights remain source-only and use source-fit seed zero throughout. Their independent source observations do not change across archive masks, so those fits are reused; every local stage that uses the degraded target archive is refitted. Twenty archive masks are not twenty new global model fits.

On each degraded archive, the retention set (90% warm-season rule or all supported years) is crossed with the annual estimator (daily plug-in, equally calibrated stochastic regression or D2). Every slope is scored against the same complete-record slope. For the four MSE cells A (rule/plug-in), B (all/plug-in), C (rule/alternative) and D (all/alternative), the symmetric retention contribution is $\{(A-B)+(C-D)\}/2$, and the estimator contribution is $\{(A-C)+(B-D)\}/2$. They sum to $A-D$; the interaction is $(A-B)-(C-D)$. Complete paired cells, rather than different available subsets, determine each decomposition.

We refit both methods' annual quantile offsets on their original disjoint source calibration groups after imposing the observed-window lower bound. Both distributions and their samples respect the bound when one is available. Seven stored quantiles identify only the central 90% of the distribution. We extrapolate both outer tails using the adjacent normal-quantile spacing and project samples onto the bound; the endpoint-constant alternative is retained on paired cases. The normal extrapolation is an assumption, not a validated tail law. We use 256 stratified samples per year and compare independent ranks, auxiliary shared year-bootstrap regression states, and three-year residual blocks on exactly the same annual marginals. Shared states come from a small harmonic/air Ridge model, not the D2 tree posterior or the selected donor model. These couplings are dependence sensitivities, not posterior samples or coverage theorems.

The task-level comparator calibrates absolute slope error divided by an independent-draw interval half-width. Every source connected group receives equal total weight in the empirical 95th percentile. Calibration uses only unused rule-calibration groups, re-predicted with the held-region model; forward errors end before 2010. We withhold the calibration when fewer than three nondegenerate source groups remain, the calibration factor is effectively zero, or the target half-width lies outside the source 1st–99th percentile range. These thresholds are




Figure 2. Fully observed thirty-day windows in incomplete years exceed complete-only station records. Crosses on the left are additional approved-record witnesses with sparse one-to-three-year complete reference support; they are not included in the original long-record denominator or the source-station bars on the right. The evidence concerns reported measurements and requires no reconstruction.

empirical support checks rather than a finite-sample guarantee. Source support, availability and matched-case width, coverage and interval score are reported. All uncertainty scores target the finite complete-record slope, not climate-trend uncertainty.

Two small stress studies select one long-record station per region and five mask repetitions. One applies the target's exact masks simultaneously to all observed donor stations; the other changes annual missingness probabilities to 0.90 or 0.18 according to whether the year's air H30 exceeds the first-seven-year 80th percentile. The latter changes the realized masks and missing-year count as well as the heat association. A station with insufficient air data is unavailable; neither study constitutes a new climatic replication.

4 Results

4.1 What the observations establish

At 20 of 96 eligible stations, 35 incomplete years contain an observed window above the maximum complete-year H30. Six stations have a witness exceeding the complete-year record by more than 0.5°C. Another incomplete year could still be

warmer, and measurement error or a changed sensor location can affect physical interpretation.

The practical consequence depends on the rule. Of the 35 witnesses, 32 pass the station-specific 90% warm-season criterion and 31 pass 95% annual coverage. Among those rejected by the warm-season criterion, 2 exceed the largest *observed window* among rule-accepted years. A rule-accepted incomplete year may itself have an unobserved higher H30, so that comparison concerns reported windows rather than a proven full-record ordering. The evidence therefore establishes a cost of strict complete-year selection; it does not attribute all 35 omissions to the more permissive rule.

4.2 Additional observation-only corroboration

We obtained approved daily-mean USGS temperature records at ten additional metadata-selected stream stations. None was a source-model target. Seven candidates lie in HUC2 regions 13 or 20, absent from the source station regions; three are additional station checks within already represented HUC2 regions. The candidates and all qualification failures are retained. This unregistered intake was not a random sample or a blinded confirmation, and the reference histories are shorter and sparser than the original long-record panel.

The unchanged four-complete-year reconstruction criterion yields zero eligible numerical-recovery references. The observation-only claim needs a different comparison: an incomplete year can contradict the maximum of the complete-year record whenever at least one approved-complete reference year exists. We retain every approved finite reference value, including reference years that an extra heuristic-QC filter might remove. Candidate windows must contain thirty consecutive approved, nonestimated daily values after source QC, and the incomplete year must have an actual absent, null, contradictory or unapproved provider day. Heuristic censoring alone cannot create a witness.

There are four stations with a nonempty approved-complete reference, each with only one to three reference years. At 3 of these stations, 9 incomplete station-years contain verified windows exceeding every approved-complete H30 in the analyzed record (Figure 2; supporting information). For example, the Rio Mora

2020 window exceeds the complete-only record by approximately 0.883°C, and a Pecos River 2020 window by 0.643°C. These facts corroborate record omission without fitting any imputer. They do not estimate an omission incidence, verify a unique warmest year, establish external reconstruction accuracy, or validate trend or risk guarantees. Every witness includes the thirty original approved values; agency reporting and measurement validity remain assumptions.

4.3 Annual point and distribution comparisons

On controlled gaps overlapping at least half the training climatological peak, the median source-repetition H30 MAE is approximately 0.428°C for the daily plug-in, 0.355 for functional debiasing, 0.337 for D2 and 0.336 for calibrated stochastic regression (Figure 3). Median 90% interval coverage is 89.3% and 89.8%, respectively, with CRPS 0.246 and 0.245. On common cases D2 improves CRPS in 7 of twenty repetitions and upper-tail CRPS in 0; equality of the methods is not established. Both distributions use the same support projection and tail approximation in this comparison. The practical evidence is a gain over the simple plug-in, not a demonstrated universal need for D2.

4.4 Whole-archive refitting, retention and uncertainty

The whole-archive benchmark comprises 3,320 station/repetition/setting evaluations at 42 long-record stations in 18 connected groups. Figure 4 and Table 2 use common station/repetition rows. Retaining years improves the finite-record slope comparison in the evaluated settings. The ranking of D2 and calibrated stochastic regression varies with time setting and region; results are conditional on the donor availability and mask design. Twenty masks at one station are not twenty independent rivers or climates.

The factorial comparison identifies a retention contribution rather than assigning one step of a sequential pipeline all of its improvement. For offline controlled gaps, the symmetric retention contribution is 93.0% of the D2-versus-rule/plug-in MSE gain. This is a result for this experiment, not a universal percentage. Estimator effects and interactions are retained in the full paired table.

Annual marginals and trend uncertainty give different answers (Figure 5). Under controlled gaps the D2 missing-data trend interval covers 94.0% offline and 88.0% forward; residual blocks give 89.0% forward. The tail and dependence sensitivities are separated: the same bounded annual marginals are used for each rank coupling, whereas endpoint and normal-tail assumptions are compared on paired cases. Annual miscalibration, extrapolated tails and dependence can all contribute to trend errors; the experiment does not uniquely attribute the residual undercoverage to any one of them. Task calibration is available for 39 of 820 forward controlled-gap D2 cases (4.8%), representing two stations; both task calibration and the independent-draw comparator cover all of these same cases. This selected-subset result does not demonstrate general improvement. Withholding and regional support remain explicit.

4.5 Observed facts, numerical estimates and unknown outputs

The release logic is stated in Figure 6. It is a design schematic; the empirical support for each gate is reported below and does not constitute an operational certification. The three-output comparison uses a point estimate only within the source feature range and a raw individual classifier-score cap, an observed-window bound when a numerical estimate is withheld, and unknown when no valid bound is available or visible readings raise a quality flag. The classifier was trained for the original unprojected D2 and is not a calibrated group-risk bound for the projected estimate. Visible anomaly limits come from the corresponding source training split, including pre-2010 data for forward evaluation. A seven-day water–air agreement flag addresses the specified exact-air-substitution scenario only; it is not evaluated in the other stress scenarios and is not a general sensor diagnostic. Where a valid seven-day water–air window is unavailable, this diagnostic is unassessed and does not itself block release. The primary factorial trend comparison is separate; an additional exploratory analysis passes released values and bounds to a Gaussian censored working likelihood.

Figure 7 reports the cost of withholding numbers. Source-range diagnostics do not make hot-year extrapolation or temporal shift safe. Exact-air substitution can be detected while sacrificing almost all numerical output in that condition; zero publication has undefined selected risk and is not successful numerical recovery.

Observed bounds retain heat information only when the reported measurements are valid. Event tables distinguish detected hot-year cases, missed or withheld cases, and false hot declarations; mask realizations are not counted as independent physical events. Mixture curves vary the imposed contamination fraction; they do not estimate its prevalence in real archives.

In the exploratory release-then-trend analysis, the forward controlled-gap RMSE is 0.202°C per decade when selected values and censored bounds enter the Gaussian working likelihood, compared with 0.173 for retaining all D2 estimates on the same cases. The nominal 95% working interval covers 99.8%, but its average width is 3.26°C per decade. That interval includes fitted regression variation, treats selected numerical estimates as exact and is not a calibrated missingness-only interval. Selection, informative censoring and the working distribution can all affect the result. In offline controlled gaps, 185 bound cases have no evaluable air-agreement diagnostic. Withholding those bounds leaves numerical publication unchanged and gives a working-likelihood RMSE of 0.040; this sensitivity does not establish sensor safety.

A seed-zero audit of 98 source model bundles found no held test station or held test connected group in any source role. However, the forward HUC17 estimator used 140 cases from 1 other training-only station in 1 connected group within the same named region, and the forward NVE interval-calibration role used 431 cases from 2 other same-region groups. The former affects interpretation of 120 forward whole-archive station/repetition settings at 3 HUC17 stations; the two affected regions account for 6,198 rows in the saved future-year release output. These are not held-target observations, but the forward source models are not uniformly trained outside the named target region.

A post-hoc strict-region sensitivity removes every source group appearing in each affected region, retains the other original role assignments and refits the two affected models. For forward controlled gaps, the common whole-archive comparison includes 738 station/repetition cases at 41 stations; original and strict D2 trend RMSE are 0.1638 and 0.1645, and stochastic RMSE are 0.1466 and 0.1466 °C per decade. Across 27,846 forward controlled-gap annual cases, the three-output publication rate changes from 63.1% to 64.5%, while failure among published values

changes from 10.9% to 11.3%. These selected sets differ. On 820 common forward controlled-gap archives, release followed by the working likelihood yields RMSE 0.2024 and 0.1986°C per decade, with average working interval widths 3.26 and 3.29. Among 2,146 annual controlled-gap cases where the complete reference exceeds the visible bound, D2 nominal 90% coverage changes from 83.2% to 82.9%. These are outcome-defined, repeated-mask comparisons. This smaller-source-sample sensitivity does not replace a strict-region rerun of every reported analysis or establish independent-climate transfer.

A further post-hoc source-rights sensitivity refits all 13 source models using seven candidate providers with recorded reuse paths, reassigns the source roles, and excludes all groups occurring in the target region. It reuses the original target masks and daily features. Among 738 common forward controlled-gap station–repetition cases in 17 connected groups, D2 and calibrated stochastic trend RMSE are 0.16381 and 0.14658 °C per decade under the original source models, versus 0.15806 and 0.15600 under the candidate models. On a different four-cell factorial cohort of 807 cases, the symmetric retention share for D2 is 93.9% originally and 93.7% for the candidate; these are ratios of mean contributions, not universal causal effects. On 820 common interval cases, nominal 95% D2 coverage is 88.0% and 87.6%, respectively. Source providers, role assignments and strict region exclusion change together, so these shifts cannot be attributed to one factor. The analysis is retrospective and does not establish independent-climate transfer or permission to release the derived models and case data.

In the small synchronized-donor study, D2's forward controlled-gap RMSE changes from 0.079 to 0.116°C per decade on 30 matched station/repetition cases with identical target masks. In the heat-associated study, the offline controlled-gap comparison changes from 0.035 to 0.058 on 30 available paired station/repetition cases; five cases from one station are unavailable because its air history is insufficient. Different year masks and missing rates prevent attributing the latter difference solely to the heat association. Both studies have few stations and are diagnostic boundaries rather than population estimates. Under strict-region source sensitivity, the same controlled-gap donor comparison changes from 0.0803 to 0.1159°C per decade on 30 common cases; the corresponding heat-associated stress RMSE is 0.0902 on that common cohort.

4.6 External qualification and stress boundaries

A subsequent Australian evaluation of a different model pipeline used the same public source archive (Australian Bureau of Meteorology, 2026). Of 80 station-years previously identified as candidate references, 79 entered that evaluation, including all 10 years previously designated for a separate F2 test. The historical F2 intake recorded its own formal test as closed, and this audit did not run it; the records are nevertheless not unopened targets for a new independent evaluation of the present pipeline. The earlier sample-support criterion allowed 20 observed hours per day, so it does not establish a strict complete 24-hour reference.

Nine Greater Wellington sites yielded no complete reference hot years under the original quality-code-600 criterion. The provider describes QC200 as unverified (Greater Wellington Regional Council, 2016), while the meanings of other returned legacy codes remain unresolved. A separate public NEON PRT archive (Marzolf, 2025) supplied half-hourly records at 24 river or stream sites. Using one fixed location code per site, zero sites had four complete years under either all 48 quality-passing half-hours per UTC day or a 46-of-48 sensitivity. Location codes do not establish identical hardware over time. The latter daily sensitivity is an observed-interval average and is not an exact complete day. An additional USGS intake of eight metadata-selected sites also produced no eligible four-year reference after prior-contact screening and daily quality checks. Five metadata-selected Réunion sites supplied Hub'Eau records (Office français de la biodiversité & Bureau de recherches géologiques et minières, 2026) with quality code 1, but none supplied four complete July–June years after requiring 24 distinct valid hours per reported day; the provider-day time-zone convention remains unresolved. These qualification outcomes are not evidence that the temperatures are wrong, and they yield no external reconstruction-accuracy estimate.

We acquired 6 ECCC Red River at Emerson files (Environment and Climate Change Canada, 2024); the site has a USGS alias (Redoloza et al., 2025) and shares a river with an upstream source station, so it is not an independent basin. The provider dictionary describes validation status V and quality grades 1–4, whereas 16,209 rows use the word VALIDATED without a grade and 13,932 rows use a station identifier different from the catalog. Under the fixed literal V and numeric-

grade-1/2 rule, 0 complete local-clock years qualified; the daylight-saving convention remains unresolved. These exclusions do not establish that the other observations are erroneous, and no recovery accuracy was scored.

We also examined all 23 published British Columbia station records (Moore & Callahan, 2024; Callahan & Moore, 2025); 4 stations had at least four calendar years with a finite published value every day. The Water Survey of Canada describes water temperature as an auxiliary output without standardized quality assurance (Water Survey of Canada, n.d.); the source study subsequently screened hourly measurements using spot checks and water–air comparisons and accepted daily means based on at least 20 hourly observations. The released derivative does not report the count and quality of each individual hour, so a complete year of published daily values is not a strict 24-hour-per-day reference. On 16 station-years with leave-one-year-out local fitting and 288 controlled-gap masks, the plug-in, D2 and calibrated stochastic H30 MAEs are 0.516, 0.306 and 0.302°C, respectively; D2 and stochastic nominal 90% coverage are 95.1% and 91.3%. These case-weighted summaries are exploratory: masks repeat within years, three sites lie in the Fraser River basin (Nechako, Thompson and Fraser), the source investigators selected their 23 sites for data availability, the published daily time-zone convention remains unresolved, and NASA air forcing uses coordinates rounded to 0.01 degree. The reference QC used water–air comparisons, so it does not independently test against that failure mechanism. This is a new station-record comparison against a weaker published-daily reference, not strict independent-climate or natural-outage validation.

A separate Nova Scotia exploratory amendment selected two stations and eight station-years after examining reported-clock completeness and agreement between two provincial releases (Government of Nova Scotia, Environment and Climate Change, 2026), but before calculating model prediction errors. Each reference day averages 24 finite, uniquely labelled hourly values; individual quality and estimated-value flags, the water time basis, and its alignment with local-solar-time NASA forcing remain unresolved. On 144 paired controlled-gap masks, D2 and calibrated stochastic H30 MAE are 0.311 and 0.313°C; nominal 90% coverage is 88.2% and 93.1%, with widths 1.000 and 1.281°C. These are repeated masks at 2 stations and 8 station-years, with single-year leave-out and three other complete years available for local fitting. A post-hoc sensitivity refits the local stages under reported-day

boundaries shifted by −1 or +1 hour while holding masks, source model, calibration and forcing fixed. The maximum change in the eight reported reference H30 values is 0.0023°C, while the largest D2 point-prediction change over the 144 paired cases is 0.094°C; this does not establish the actual time basis or the cause of model sensitivity. This after-the-fact availability and source-agreement selection is not strict reference qualification, independent-climate confirmation, whole-archive degradation or a natural-outage test.

A further metadata-fixed USGS preflight excluded the known source, prior panels, downloaded caches and an API pilot station whose annual means were inspected and which had also been scored in an earlier E10 analysis. Of five remaining candidates with prior network screening, none had four calendar years of complete approved daily counts in 2010–2024 according to the USGS annual statistics service (U.S. Geological Survey, n.d.). The count screen may include estimated values and does not establish subdaily quality; no daily reconstruction was scored for these five stations. This availability-selected exploratory search does not describe the proportion of qualifying USGS stations overall.

An initial 15-station USGS intake illustrates why the access inventory matters: all 15 had already been processed in the national panel and ten had been scored in an earlier external comparison. They are opened historical evidence, not new independent validation. The subsequent eight-station roster was selected after excluding those contacts and other known source, evaluation and cached-value records. The prior-access and river-network screens are best effort; neither they nor different HUC identifiers prove independent weather or errors. All these intakes are exploratory and unregistered.

A complementary clean retrospective comparison selects the score-only cutoff in the held-region model's disjoint source rule-calibration groups, approximately matching the three-output numerical rate in those groups; it uses neither the target outcomes nor the target release count. On held-region empirical calendars, the three-output and source-calibrated score-only rules publish 85.5% and 84.9%, with failure rates 0.701% and 0.030%, respectively. On controlled gaps their publication rates are 72.1% and 71.0%, with failure rates 4.90% and 3.87%. The target retention rates

differ, and this one-seed experiment does not establish a population-level tradeoff or group safety.

In the controlled-gap comparison, 24 of 26 regions with numerical outputs from both rules have a higher conditional error rate for three outputs, while 2 have a lower rate; 1 region has undefined error because three outputs publishes none. These regional signs do not compare equal retention. For example, at the Finnish sites the three-output and score-only rates are 3.94% and 8.86%, while numerical retention is 9.9% and 58.3%. On the 172 of 185 connected groups where both rules publish, the equal-group mean calendar-case error rates are 0.829% and 0.015%. Dropping each evaluation region in turn leaves the pooled error difference positive, with all models and source thresholds frozen; this is not a refitted leave-region-out study or a set of independent significance tests.

A retrospective comparison holds the numerical-output count fixed on each evaluated feature distribution. In clean empirical-calendar cases, both rules publish 27,090 values; three outputs have 190 failures (0.701%), while ranking cases only by the uncalibrated individual score yields 4 (0.015%). For forward controlled gaps, the corresponding 17,583 values have failure rates 10.94% and 6.94%. Score ties use case identifiers, not outcomes. This comparator chooses its threshold from the evaluated score distribution and the three-output count, so it is a transductive descriptive benchmark rather than a prospectively calibrated policy. Equal numbers of numerical outputs do not equalize information retained as bounds or unknown states. In the imposed air-exposure calendar, only 88 numerical values remain; the two methods have 19 and 23 failures, respectively. These case-dependent results offer no group-risk guarantee.

The release comparison also exposes limits. In clean empirical-calendar cases, the fixed-score rule publishes about 94.7% of values at 0.7% observed error, while the three-output rule publishes about 85.5% at a similar error rate. Under the imposed air-exposure stress, it publishes almost no numerical values. Hot-year and later-year stresses remain challenging despite source-range checks. These synthetic scenarios have no inferred field prevalence, and withholding almost every result is not a useful recovery guarantee.

5 Discussion

For the observation-only result, 971,673 finite daily values in 34 selected project input panels agree with their float32 archive casts; the maximum rounding difference is 0.00000153°C. No witness identity or 0.5°C margin classification changes when the selected full-precision panels replace the canonical archive. These panels are themselves processed inputs and do not establish sensor accuracy.

The strongest observation result concerns the choice of reference archive. A fully observed window can establish that a strictly complete-year-only record misses a reported heat event without requiring an imputer. However, most witnesses would survive a less restrictive warm-season screen. The comparison with existing annual-signal and thermal-peak studies therefore matters: the result does not show that every partial-year practice is unnecessarily restrictive. It motivates reporting exactly which data and record statistic a rule excludes.

The whole-archive design shows that preserving supported years can improve the slope of the particular complete record used as reference. The estimate is conditional on the long-record sites, mask design, observed donors and current-vintage forcing. The 42 stations represent 18 connected groups, with 24 stations in Bavaria. Group resampling and leave-region-out comparisons are descriptive and do not establish independent climates. When Bavaria is excluded, the forward controlled-gap D2-minus-stochastic MSE difference changes from -0.000143 in the original analysis to +0.000323 under the strict-source sensitivity, in $(°C/\text{decade})^2$, on 331 common station–repetition cases; the sign of this small difference is sensitive to the source model. The finite-record slope is distinct from an underlying climate-process trend; neither point-error reduction nor a wide total regression interval proves that missing-data uncertainty is calibrated.

On the common forward controlled-gap cohort (818 of 820 station–repetition cases, 41 stations in 17 connected groups), the incomplete-year-weighted D2 90% annual coverage is 90.1%; equal weighting of station–repetition cases gives 89.6%. With matched random ranks in the tail experiment, constant versus normal-spacing tails yield 83.0% versus 88.8% coverage, widths 0.380 versus 0.451 and interval scores 0.987 versus 0.860 for the nominal 95% slope interval. In the separate fixed-marginal coupling experiment, independent, auxiliary shared-parameter and residual-

block ranks yield coverage 88.0%, 87.9% and 89.0%, with interval scores 0.868, 0.854 and 0.830, respectively. The two experiments use different random draws; annual calibration has not been experimentally repaired. These contrasts are not additive causal contributions, and their cohort excludes cases where a comparison is unavailable.

The bounded annual representations also contain many degenerate 90% intervals. On the 3,674 common forward empirical-calendar cases with a finite bound, D2 has 1,476 zero-width intervals (40.2%) and calibrated stochastic reconstruction has 2,977 (81.0%); all of these coincide with the complete reported reference within numerical tolerance. In the 3,683 common forward controlled-gap cases, calibrated stochastic reconstruction has 1,091 zero-width intervals, 3 of which miss the reference across 2 stations; the largest miss is 0.143°C. Among the 1,186 D2 cases whose median equals the bound, the reference is higher in 64; among the 1,375 corresponding stochastic cases, it is higher in 61. These are repeated masked cases, not independent rivers. Matching an observed bound can accurately recover the reported complete-record statistic when its maximum window is already visible, but does not establish zero sensor uncertainty. Replaying the fixed models and projection-aware offsets shows that, in the forward empirical-calendar cases, 380 D2 intervals first become zero-width at the final projection and 1,096 are already zero-width immediately before it. For the stochastic comparator, 0 of its zero-width intervals first appear at this last step; 3,090 raw stochastic intervals are already zero-width before calibration and the explicit projections. These stochastic quantiles come from a finite set of imputations that retains the observed daily values; raw therefore does not mean unconstrained by observations. The pre-final values still reflect the first projection and offsets calibrated with that projection; this replay does not measure the effect of refitting under an alternative calibration protocol.

An outcome-defined diagnostic separates cases in which a fully observed window reaches the complete reported H30, within 10^{-8} °C, from cases in which the reference exceeds the observed bound. In the 3,674 common forward empirical-calendar cases, the bound matches the reference in 3,125; on the remaining 549 cases, D2 and stochastic 90% interval coverage are 88.3% and 87.8%, respectively. In the 2,146 of 3,683 common forward controlled-gap cases whose reference exceeds the bound, the corresponding coverages are 83.2% and 81.6%; descriptive connected-

group bootstrap intervals are [72.6, 93.0]% and [72.2, 91.7]%, respectively. This stratification uses the hidden complete reference and is not a decision-time diagnostic. Repeated masks can place the same station-year in both strata, and group resampling does not establish transfer across independent climates.

The candidate-provider strict-region refits were also compared on the same four-way common annual cases. Among 2146 forward controlled-gap cases at 41 stations in 17 connected groups where the complete reported H30 exceeds the visible bound, D2 nominal 90% coverage changes from 83.2% to 81.9%, while its mean width changes from 1.299 to 1.161°C. Calibrated stochastic coverage changes from 81.6% to 78.2%, with mean width 1.121 to 1.078°C. This outcome-defined stratum cannot guide a real-time decision. For the classical method, the candidate-minus-original coverage change is -3.4 percentage points overall, but +0.3 after omitting FOEN and -8.9 after omitting Bavaria from the fixed evaluation cases. These deletions do not refit models; source composition, role assignment and strict region exclusion change together in the post-hoc refit, so no single change can be assigned as the cause. Neither version establishes reliable coverage on independent climates.

We explored an additive annual interval expansion estimated from source rule-calibration groups excluded from model fitting, the original interval calibration and the held region. It uses the equal-group-weighted empirical 90th percentile of nonnegative interval misses; a region is unsupported when fewer than three source groups contribute or the empirical correction is effectively zero. Forward source cases end before 2010. In the controlled-gap forward comparison, both methods receive a supported correction for only 92 of 3,683 common cases (2.5%), all from 1 station. On those same cases, D2 coverage is 100.0% both before and after expansion, while mean width increases from 0.532 to 0.543°C. The stochastic interval likewise covers 100.0% before and 100.0% after, with width increasing from 0.555 to 0.567°C. Neither improvement in general coverage nor a finite-sample group guarantee follows: the empirical correction is exploratory, the source groups are few, and the supported cases are highly selected. A 90% unrandomized rank based on one maximum nonconformity score per group requires at least nine calibration groups for a finite correction; at most 4 groups are available here. Such a rank would also require appropriate group exchangeability. This experiment does not recalibrate a joint trend distribution.

A subsequent post-hoc experiment uses the candidate-source models and single-year masked cases from their rule-calibration groups to estimate an equal-group-weighted empirical 90th percentile of annual interval misses. These groups are disjoint from estimator fitting and original interval calibration, although some contributed to the separate risk-rule component; they are not a newly unopened dataset. Forward source cases precede 2010. Corrections are withheld for fewer than three contributing groups or a zero empirical radius. On 3411 of 3683 common forward controlled-gap annual cases (38 stations, 14 connected groups), both methods have a supported correction. On this same selected subset, D2 90% coverage changes from 88.7% to 89.9%, with width 0.705 to 0.755°C and interval score 1.4599 to 1.4602; the classical method changes from 86.6% to 89.1%, with width 0.608 to 0.696°C and interval score 1.4867 to 1.4601. In the outcome-defined hidden-peak subset of 1982 supported cases, coverage is 81.0% to 82.9% for D2 and 76.9% to 81.3% for the classical method. Single-year source masks do not reproduce jointly degraded target archives. The forward rule role has at most eight groups, so a finite 90% one-maximum-per-group rank is unavailable; the empirical correction has no finite-sample or independent-climate coverage guarantee and does not by itself identify a calibrated joint trend distribution.

A further post-hoc propagation uses the same masks and rank matrices for the source-supported annual corrections. The original forward controlled-gap sample has 820 station–repetition settings; 760 have source support for both methods, and 685 nontrivial settings at 38 stations in 14 connected groups have both annual distributions and all three rank constructions available. Among the supported settings, 73 lack at least one annual distribution and two contain no incomplete year. Under the independent-rank construction, D2 nominal 95% trend coverage is 87.9% before adjustment, 89.3% when only the annual 5th and 95th quantiles are shifted, and 88.9% when interior quantiles are also tapered outward; corresponding widths are 0.382, 0.409 and 0.403°C per decade. The classical method has coverage 83.4%, 86.9% and 85.5%, with widths 0.344, 0.387 and 0.376°C per decade. Both adjustment schemes have identical annual 90% endpoints, preserve the median and use the same ranks, but assume different interior and normal-spacing outer-tail shapes. The source calibration is based on single-year masks and few groups, and these retrospective results do not establish a calibrated joint distribution or independent-climate coverage.

Annual and trend coverage also depend on the assumed marginal tails. Extending the stored quantile function changes coverage even with an unchanged rank ordering, while auxiliary parameter and residual-block couplings test only particular dependence structures. A credible future joint model must account for its actual fitted-model uncertainty and retain support constraints. Task-level calibration can be useful only where independent calibration groups and relevant source difficulty are available; the small supported forward subset here cannot justify a general 95% claim.

The decision results caution against interpreting a low average selected error as protection for every river. The imposed exposure stress is more easily detectable than unknown field faults, and measurement validity is a prerequisite for treating L_y as a physical bound. Changes in groundwater influence, flow, regulation and sensor deployment can alter the water–air relationship. Their effects cannot be inferred from a model residual alone. New high-quality reference measurements are needed to evaluate natural outages and transfer beyond these archives.

6 Conclusions

Incomplete river-temperature records can preserve observed sustained-heat information, and retaining years reduced finite-record trend RMSE relative to the evaluated warm-season screen in the tested degradation settings. The benefit of rejecting or retaining a year depends on the actual completeness rule and the intended scientific quantity. A well-calibrated classical reconstruction remains a strong annual comparator. Forward trend intervals, three-output release rules and external transfer require their own evidence; the present results do not certify universal numerical recovery or group safety.

Open Research

The source observations, processing code, case-level results and figure scripts require a permanent repository and citations to be supplied by the authors. A local review candidate includes processed daily values from 95 of the 96 previously eligible source stations, permitting an independent arithmetic audit of all 35 observation witnesses and their screening-rule decisions. The LUBW station is omitted pending permission, so the full eligible population is not reconstructed. The candidate also

includes ten additional USGS daily API snapshots and their observation-only replay; neither check reconstructs the main models or verifies subdaily sampling. A local workflow, source inventory and unresolved data-rights details are maintained in the accompanying repository. The source includes observations from LUBW; its reuse and redistribution terms require provider-specific resolution rather than an assumed open licence. Although all whole-archive target stations have recorded reuse paths, 13 of 13 source-model estimator roles and 6 interval-calibration roles include positive-weight LUBW cases; the access and sharing conditions of the resulting models and case-level derivatives remain unresolved. The BC published daily derivative and NEON qualification use the CC-BY author archives (Moore & Callahan, 2024) and (Marzolf, 2025); the USGS intake uses the public daily-values collection (U.S. Geological Survey, 2026). The Nova Scotia reported-clock subset is derived from the provincial open-data release (Government of Nova Scotia, Environment and Climate Change, 2026) and uses NASA POWER forcing (NASA Langley Research Center, 2026); the separately accessed legacy workbooks require their own reuse clarification. Derived files record candidate rosters and unavailable cases. Author, affiliation, funding, conflicts of interest, permitted reviewer access and permanent deposit information remain to be supplied before submission.

References

Australian Bureau of Meteorology. (2026). *Water data online: water temperature time series.* `http://www.bom.gov.au/waterdata/`.

Beaufort, A., Diamond, J. S., Sauquet, E., & Moatar, F. (2022). Spatial extrapolation of stream thermal peaks using heterogeneous time series at a national scale. *Hydrology and Earth System Sciences*, *26*, 3477–3495. doi: 10.5194/hess-26-3477-2022


Caissie, D. (2006). The thermal regime of rivers: a review. *Freshwater Biology*, *51*, 1389–1406. doi: 10.1111/j.1365-2427.2006.01597.x

Callahan, L., & Moore, R. D. (2025). Evaluation of the hybrid air2stream model for simulating daily stream temperature during extreme summer heat wave and autumn drought conditions. *Hydrological Processes*, *39*(1), e70033. doi: 10.1002/hyp.70033


Environment and Climate Change Canada. (2024). *Automated fresh water quality*

318 **Table 2.** Whole-archive trend errors on identical available station/repetition rows, in degrees C
319 per decade.


Setting	Reference	Paired n	D2 RMSE	Reference RMSE
calendar / forward	Calibrated stochastic	701	0.062	0.051
calendar / forward	Daily plug-in	820	0.067	0.088
calendar / offline	Calibrated stochastic	835	0.009	0.011
calendar / offline	Daily plug-in	840	0.009	0.014
crossed / forward	Calibrated stochastic	738	0.164	0.147
crossed / forward	Daily plug-in	820	0.173	0.228
crossed / offline	Calibrated stochastic	838	0.028	0.032
crossed / offline	Daily plug-in	840	0.028	0.046



687 **Figure 3.** Annual estimation under difficult controlled gaps. Both historical source-calibrated
688 methods receive the same observed-bound projection and normal-spacing tail model. Bars show
689 empirical repetition ranges within the same archive; these are not independent climates.



690 **Figure 4.** Whole-archive trend errors and the symmetric retention/estimator decomposition.
691 Left: D2 with identical masks and either retention set. Right: D2 and calibrated stochastic
692 alternatives, with complete paired MSE cells. Contribution percentages are specific to each
693 experimental setting.



694 **Figure 5.** Annual 90 percent and finite-record 95 percent trend intervals are evaluated sepa-
695 rately. The rank methods preserve identical bounded annual marginals. Tail extrapolation is an
696 assumption and source-group uncertainty is not shown by the bars.




Figure 6. The evaluated three-output decision and exploratory trend analysis. The cap applies to an individual raw classifier score, not a certified group risk. The seven-day water–air check was assessed only for the clean and imposed exact-air-substitution cases. The selected-trend working likelihood treats released values as exact and observed bounds as censored.




Figure 7. Left: three outputs for empirical-calendar masks. Right: numerical-publication rates and selected errors for controlled gaps. Source diagnostics do not remove extrapolation failures. The air-agreement flag addresses the specified exact-substitution stress only; zero publication is shown as unknown rather than zero-risk success.

monitoring and surveillance data. Retrieved from `https://open.canada.ca/data/en/dataset/f258b0c8-7871-4572-b567-1ba2bd55f1b6` (Open Government Licence; ECCC Red River files and metadata accessed October 2026)

Gneiting, T., & Raftery, A. E. (2007). Strictly proper scoring rules, prediction, and estimation. *Journal of the American Statistical Association*, *102*, 359–378. doi: 10.1198/016214506000001437

Government of Nova Scotia, Environment and Climate Change. (2026). *Surface water quality monitoring network continuous water quality data.* Retrieved from `https://data.novascotia.ca/datasets/bkfi-mjgw` (Open Government Licence – Nova Scotia; accessed October 2026)

Greater Wellington Regional Council. (2016). *Environmental monitoring: quality codes.* Retrieved from `https://cms.graphs.gw.govt.nz/envmon/whats-new` (Quality-code explanation; accessed October 2026)

Hirsch, R. M. (1982). A comparison of four streamflow record extension techniques. *Water Resources Research*, *18*(4), 1081–1088. doi: 10.1029/WR018i004p01081

Johnson, Z. C., Johnson, B. G., Briggs, M. A., Snyder, C. D., Hitt, N. P., & Devine, W. D. (2021). Heed the data gap: guidelines for using incomplete datasets in annual stream temperature analyses. *Ecological Indicators*, *122*, 107229. doi: 10.1016/j.ecolind.2020.107229

Marzolf, N. (2025). *Ecosystem metabolism estimates from the national ecological observatory network stream and river sites.* figshare. (Public author archive, version 1) doi: 10.6084/m9.figshare.28074452.v1

Matalas, N. C., & Jacobs, B. (1964). *A correlation procedure for augmenting hydrologic data* (Tech. Rep. No. 434-E). U.S. Geological Survey. doi: 10.3133/pp434E

Michel, A., Brauchli, T., Lehning, M., Schaefli, B., & Huwald, H. (2020). Stream temperature and discharge evolution in Switzerland over the last 50 years: annual and seasonal behaviour. *Hydrology and Earth System Sciences*, *24*, 115–142. doi: 10.5194/hess-24-115-2020

Moore, R. D., & Callahan, L. (2024). *Evaluation of the hybrid air2stream model for simulating daily stream temperature during extreme summer heat wave and autumn drought conditions: data sets and scripts.* Zenodo. (Version 1, CC BY 4.0) doi: 10.5281/zenodo.14502248

Näf, J., Spohn, M.-L., Michel, L., & Meinshausen, N. (2023). Imputation scores. *The Annals of Applied Statistics*, *17*, 2452–2472. doi: 10.1214/22-AOAS1727

NASA Langley Research Center. (2026). *POWER project data sets: daily 2 m air temperature (MERRA-2).* `https://power.larc.nasa.gov/`.

Office français de la biodiversité, & Bureau de recherches géologiques et minières. (2026). *Hub'Eau: continuous stream temperature data.* Retrieved from `https://hubeau.eaufrance.fr/api/v1/temperature/chronique` (Quality-qualified Réunion records retrieved October 2026)

Redoloza, F. S., Glas, R. L., Nustad, R. A., & Ryberg, K. R. (2025). *Evaluating drought risk of the red river of the north basin using historical and stochastic streamflow upstream from emerson, manitoba* (Tech. Rep.). U.S. Geological Survey. Retrieved from `https://pubs.usgs.gov/publication/sir20255002/full` (Identifies the Emerson ECCC station and USGS streamgage alias) doi: 10.3133/sir20255002

U.S. Geological Survey. (n.d.). *Statistics documentation: supporting information for daily data statistical values.* Retrieved from `https://waterdata.usgs.gov/statistics-documentation` (Approved daily sample counts, accessed October 2026)

U.S. Geological Survey. (2026). *Water data APIs: daily values collection (OGC API–features).* `https://api.waterdata.usgs.gov/ogcapi/v1/collections/daily`.

van Hamel, A., & Brunner, M. I. (2024). Trends and drivers of water temperature extremes in mountain rivers. *Water Resources Research*, *60*, e2024WR037518. doi: 10.1029/2024WR037518

Vogel, R. M., & Stedinger, J. R. (1985). Minimum variance streamflow record augmentation procedures. *Water Resources Research*, *21*(5), 715–723. doi: 10.1029/WR021i005p00715

Water Survey of Canada. (n.d.). *Frequently asked questions: water level and flow.* Retrieved from `https://wateroffice.ec.gc.ca/contactus/faq_e.html` (Temperature is an auxiliary output without standardized quality assurance; accessed October 2026)