

A Cretaceous Time Scale¹

J. E. VAN HINTE²

Houston, Texas 77001

Abstract An emendation of published Cretaceous time scales is proposed on the basis of presently available age data. The standard geochronologic subdivision is combined with a linear numeric scale, biostratigraphic framework, and geomagnetic-reversal time scale. The biostratigraphic framework includes standard Tethyan and boreal pelagic macrofossil zonations and planktonic as well as benthic microfossil zones and biohorizons. Basement ages at some Deep Sea Drilling Project sites are evaluated and related to the geomagnetic-reversal scale.

INTRODUCTION

Integration of marine and terrestrial geophysical, paleontologic, paleomagnetic, and other historical geologic data and application of new quantitative techniques important in stratigraphic exploration necessitates an increasingly refined numeric geologic time scale. This paper is an attempt to combine a Cretaceous numeric time scale with biostratigraphic schemes and the geomagnetic-reversal scale.

In 1964 the Geological Society of London published the Phanerozoic time scale (PTS), which it emended in 1971 (Harland and Francis, 1971). Other scales have been published since 1964 (see Fig. 1), but the original PTS scale still is the most widely used for the Cretaceous. Casey (1964, p. 199), who covered the Cretaceous for the PTS, clearly stated that the scale was "set up purely as a basis for discussion in the absence of a more positive scale of calibration." In other words, the scale was considered to be "no more than a simple working hypothesis" (Internat. Union Geol. Sci., 1967, p. 378). The PTS 1964 scale assumed that all Cretaceous stages represented equal time intervals (6 Ma each; 1 Ma = 1 megayear = 1 million years).

Using the PTS during the past few years as a working model, the writer somewhat changed it (and will do so further) either when new relevant data became available or for pragmatic reasons when inconsistencies in the results of its application suggested such change. An early version of this scale was published in 1972 with a geomagnetic-reversal scale and a planktonic foraminiferal scheme (van Hinte, 1972). A later unpublished version was used successfully by Vail et al (1974), by Berggren et al (1975), and in unpublished reports. A wider distribution of its present emended

form (Fig. 2) seems useful in anticipation of establishment of a more authoritative scale. For further general considerations, see van Hinte (1976).

RADIOMETRIC DATA

Published Cretaceous radiometric age determinations have been compiled by the Geological Society of London (1964; Harland and Francis, 1971) in a numbered-item list and generally are referred to as "PTS items." To obtain an overview of the published information, the writer plotted the radiometric age of selected PTS items, and of data published after 1971, against a relative geologic time scale (Figs. 3, 4). In these figures, the numeric dates are presented without their analytic error (about 2 to 5 percent), but their possible relative-age span is given by a vertical line.

The relative-age span was determined from the published information in accordance with the biostratigraphic scheme of Figure 2. This scheme enables one to plot directly most of the later European dates, and very few adjustments had to be made in the stratigraphic ages of the PTS items. For the North American data, the writer followed Obradovich and Cobban (1975) by drawing the boundary of the Campanian/Maastrichtian³ at the base of the *Baculites reesidei* Zone, the Santonian/Campanian boundary between the *Desmoscaphtes bassleri* Zone and the *Scaphites*

© Copyright 1976. The American Association of Petroleum Geologists. All rights reserved.

¹Manuscript received, August 29, 1975; accepted, October 10, 1975.

²Exxon Production Research Co.

I thank W. A. Berggren (Woods Hole) for continuously stimulating me to improve and expand on earlier published and unpublished versions of "a Cretaceous time scale," and am grateful to Exxon Production Research Company's management for giving the opportunity to complete this work. R. M. Jeffords and D. O. Smith (both of Houston, Texas) critically read the manuscript, and their corrections and suggestions are highly appreciated. This compilation would have been impossible without the support of EPRCo.'s library staff.

³The author's spelling of Maastrichtian is contrary to the spelling used by the *Bulletin* (Maestrichtian) but has been retained at the author's request. His request to use "planktic" in lieu of "planktonic" as the adjectival derivative of "plankton" was not honored.

hippocrepis Zone, the Coniacian/Santonian boundary below the *Inoceramus undulaticus* Zone, the Cenomanian/Turonian boundary below the *Inoceramus labiatus* Zone, and the Albian/Cenomanian boundary above the *Neogastropiles* Zones (see also Owen, 1973). The Turonian/Coniacian boundary, however, is drawn above the *Inoceramus deformis* Zone (cf. Seitz, 1956; Ernst, 1970).

Owens and Sohl's (1973) glauconite dates from the New Jersey Monmouth Group were plotted as Maastrichtian and those from the Matawan Group as Campanian (Olsson, 1964, found *Globotruncana gansseri* and *G. calcarata* in these units, respectively), and the radiometric ages were averaged per formation.

The PTS item 365 Bearpaw Formation data were grouped as in Figure 3 of Folinsbee et al (1965), whose ages also were followed (365 a-h); the Kneehills Tuff sanidine dates were averaged per locality (365 i, j, l), the biotite date being kept separate (365 k). Caldwell (1968) and North and Caldwell (1970) reviewed the relative-age assignment of the Bearpaw Formation.

Lambert (1971a) suggested rounding of pre-Tertiary major boundary ages to 5 Ma. The present scale follows his -65 and -135 Ma for the ages of the end and the beginning of the Cretaceous. With respect to the Cretaceous-Jurassic boundary, Lambert stated somewhat pessimistically: "At the very best the boundary may fall in the range 125 to 145 m.y. and there is no sound reason for taking the mean figure as the most likely." The radiometric age of the end of the Cretaceous is known more accurately and in all likelihood could not be as far from -65 Ma as the round figures -60 or -70 Ma; the correct age, though, very well could be -64 rather than the -65 Ma generally used (Berggren, 1972) and here retained.

More controversial is the third starting point of our scale: -100 Ma for the Lower/Upper Cretaceous (= Albian/Cenomanian) boundary. Age assignments for this boundary in literature range from -93 to -110 Ma (Fig. 1). Bentonite (Obadovich and Cobban, 1975) and glauconite (Juignet et al, 1975) dates obtained recently suggest that the Albian indeed may have ended 5 or 6 Ma later, that is, at -95 or -94 Ma. Other dates (PTS items 202, 226, 319, 336; van Hinte et al, 1975), however, indicate that the early Cenomanian may be considerably older than -95 Ma, and we decided not to change Casey's 1964 estimate of -100 Ma (cf. Rubinshteyn, 1967, p. 113-121).

With the three main points of the scale fixed at conservative figures, somewhat more flexibility seems justified in diverging from the PTS "equal

stage" principle with respect to other stage boundaries. Radiometric dates (Figs. 3, 4) strongly indicate that the Campanian/Maastrichtian boundary should be drawn at -69 or -70 Ma. The Santonian/Campanian and Coniacian/Santonian boundaries rather clearly are at -78 and -82 Ma. The Turonian/Coniacian boundary can be drawn at -86 Ma between the oldest Coniacian glauconite date and the youngest Turonian bentonite date. Acceptance of these boundaries gives the Coniacian and Santonian equal duration, both being shorter than other Late Cretaceous ages, and has the advantage that equal duration can be assigned to Coniacian, Santonian, and Campanian ammonite zones. Most authors place the Cenomanian/Turonian boundary at -90 Ma or slightly younger, but some have it at -94 or -95 Ma and few at -99 or -100 Ma (Fig. 1). In the absence of accurate data, a boundary age of -92 Ma seems reasonable—especially because this gives the same duration to ammonite zones in both the Turonian and the Senonian.

Radiometric dates for the Lower Cretaceous are scarce, and nearly all are based on glauconites which become less reliable with increasing age of the section. Two Albian dates (PTS items 203, 217) led some authors (Bandy, 1967; Suppe, 1969; Dickinson and Rich, 1972; Baldwin et al, 1974) to accept an "old Albian" (see Fig. 1), whereas Casey (1964) and others placed more weight on a younger (-99 Ma) date on item 217 and took into account that the Albian age for item 203 is a youngest stratigraphic age. The Albian and Aptian beginnings are drawn in this work near the oldest glauconite ages at -108 and -115 Ma, respectively.

Having no dates available to help determine the Hauterivian/Barremian boundary, with the Hauterivian likely being younger than 127 Ma (item 75), the Valanginian being older than 124 Ma (Suppe, 1973, 105-124 Ma old ages for metamorphic Hellhole-Williams rocks with Tithonian-Valanginian *Buchia*), and the Berriasian possibly older than 131 Ma (items 177, 215, 328), the writer drew the boundaries at -121, -126, and -131 Ma. Thus, the present model assumes equal duration for the Valanginian and Hauterivian, both being shorter than the Barremian and longer than the Berriasian. However, this assumption is based largely on intuition and on reasons of practical utility.

The feasibility of the proposed model may be judged from Figure 5, which summarizes the available chronometric information and illustrates its position with respect to the preferred scale and an "equal stage" scale.

HISTORY OF CRETACEOUS NUMERICAL TIME-SCALE





FIG. 1—Different numeric time scales proposed since 1959 for the Cretaceous. No attempt has been made to convert numbers of early and Russian scales to 1964 PTS standards. Maastrichtian, Turonian, and Albian ages are shaded for easy comparison. Gill and Cobban (1966), Pergament (1967), Kauffman (1970), and Obradovich and Cobban (in press) presented their scale with a molluscan biostratigraphic scheme; Bandy (1967), Sigal (1967), and van Hinte (1972) scales were accompanied by planktonic foraminiferal zonation.

Alb, Albian; *Apt*, Aptian; *Ba*, Barremian; *Be*, Berriasian; *Ca*, Campanian; *Ce*, Cenomanian; *Co*, Coniacian; *H*, Hauterivian; *J*, Jurassic; *M*, Maastrichtian; *Neoc*, Neocomian; *S*, Santonian; *Te*, Tertiary; *Tu*, Turonian; *V*, Valanginian.



BIOSTRATIGRAPHIC SCHEMES

The French colloquia on the Upper and Lower Cretaceous set the stage for relating macrofossil biostratigraphy to the type sections of stages. The conclusions of the colloquia (Dalbiez, 1959; Anon., 1963) form the basis for the Cretaceous ammonite zonation of Figure 2. Subsequent work has refined this "standard" and established its correlation with the boreal faunal succession (Fig. 2, authors listed at base of column).

The planktonic foraminiferal scheme is after van Hinte (1972) and is, with the alterations (Figs. 6-8) briefly discussed later, firmly related now to the stratotypes of stages through first-order and second-order (macrofossils, benthic foraminifers) correlation. The zones have been given letter-number designations for easy reference.

Recent work on stratotypes and on correlations between benthic and planktonic foraminiferal zones, between foraminiferal and macrofossil zones, and between boreal and Tethyan biostratigraphic subdivisions provided a better basis for relating the planktonic foraminiferal scheme to the Santonian and Coniacian stratotypes and stage boundaries (i.e., Ernst, 1970; Scheibnerova, 1972; Séronie-Vivien, 1972; micropaleontologic literature listed under Figure 6; see also discussion in Barr, 1972, p. 6).

The top of the *Rotalipora cushmani* Zone now is drawn to coincide with the Cenomanian/Turonian boundary for reasons illustrated in Figure 7, a conclusion that had been reached earlier by other authors for the same or for different reasons (cf. Bolli, 1966; Pessagno, 1969, p. 22).

Wider geographic recognition of Moullade's (1966) Albian-Aptian foraminiferal ranges and biostratigraphy and repeated confirmation of the phylogenetic interpretation suggest that a generalization of his scheme is justified. The derivation of the present zonation for this part of the section is illustrated in Figure 8.

Van Hinte (1972) selected planktonic foraminiferal biohorizons that are recognized widely and

most likely have time significance; in Figure 2 some Senonian biohorizons were added after Herm (1965), Premoli Silva and Bolli (1973), and Smith and Pessagno (1973).

The present biostratigraphic scheme includes a neritic and littoral benthic foraminiferal phylozonation for accurate dating of Senonian shallower water sediments in which pelagic markers are lacking (Fig. 6). In Figure 2, other widely distributed benthic markers of proved time significance were added for the same purpose.

Allemann et al (1971) and Thierstein (1973) related their respective calpionellid and Early Cretaceous nannofossil zonations to macrofossil zones and/or to foraminiferal zones and to type sections of stages. The stratigraphic ages of their boundaries and biohorizons, therefore, are reliable. Those of the Late Cretaceous nannofossil scheme and of the radiolarian zonation are less well established and, therefore, had to be drawn as broken lines in Figure 2.

GEOMAGNETIC-REVERSAL SCALE

Since 1968, various geomagnetic-reversal time scales have been published for the Cretaceous (Fig. 9), some based on measurements made on land, others on marine-anomaly data. The scale presented here combines several of the earlier proposals. The subdivision of the Beringov interval is after the Heiztler et al (1968) scale as revised by Sclater et al (1974). The Serra Geral interval is recalibrated after Larson and Pitman (1972) using -145 and -125 Ma as calibration points instead of -155 and -120 Ma (see van Hinte, 1976). Green and Brecher (1974), Green et al (1974), and Jarrard (1974) recorded three brief reversed zones in upper Albian sediments (Proto Decima, 1974) at DSDP Site 263 which are included in the conclusion column as "Site 263 Mixed" Zone. Keating et al (1975) mentioned that the Akoh Formation recently has been dated radiometrically as 92-104 Ma old, which suggests that the Akoh reversals are within the Site 263 Mixed Zone. But because no further reference to

→

FIG. 2—Numeric Cretaceous time scale proposed in this paper. Younger part of geomagnetic-reversal scale is after Sclater et al (1974), older part is after Larson and Pitman (1972, recalibrated see text), and middle zones are after McElhinny and Burek (1971), Green and Brecher (1974), and Jarrard (1974). Geomagnetic interval and zonal names are after McElhinny and Burek (1971) and Pechersky and Khramov (1973). Foraminiferal scheme is newly compiled (see text); other biostratigraphic data are after literature listed at bottom of each column in figure. Foraminiferal literature listed refers to benthic biohorizons. Note that first and last occurrences of *Globotruncana* and *Bolivina* species do not refer to typological species (see text). Most planktonic foraminiferal zones are phylozones, and they as well as selected biohorizons can be expected to have good geochronologic reliability.




FIG. 3—Plot of radiometric age against relative age for Cretaceous dates on minerals other than glauconite. Data after literature listed on figure. PTS items are those selected by Casey (1964) plus items 334, 335, and 336 and minus items 198 and 199.




FIG. 4—Plot of radiometric age against relative age for Cretaceous glauconites. Data after literature listed on figure. PTS items are those selected by Casey (1964) plus items 51, 59, 207, 213, 228, 233, and 239.




FIG. 5—Summary of Figures 3 and 4. Figure illustrates relation of published radiometric data with preferred scale (shaded) and compares line representing equal duration of stages with line representing preferred scale.




FIG. 7—Cenomanian-Turonian macrofossil zonations, some planktonic foraminiferal ranges, and different opinions on position of Cenomanian/Turonian boundary (heavy line). This paper follows resolutions of Colloque sur le Cretace Supérieur Français (Dalbiez, 1959) which are in accordance with Muller and Schenck (1943) in including *Actinocamax plenus* Zone in Cenomanian. No linear vertical scale.

this new radiometric date is given, we provisionally leave the Akoh Reversed Zone at -79 to -77.5 Ma, following Helsley and Steiner (1969) and McElhinny and Burek (1971). The Gatan Reversed Zone is placed near -108 Ma, as it was assigned to the Albian-Aptian boundary by Pechersky and Khramov (1973). Jarrard (1974) suggested that he may have recorded this zone in sediments of DSDP Site 260.

Watkins and Cambray (1971) studied Campanian-Santonian dikes from Jamaica and found that "a normal geomagnetic field dominates (but not completely so)" and that their reversely magnetized dikes might be Maastrichtian or Santonian or "we may have detected the first Campanian reversed polarities." The second alternative would confirm the extrapolation of marine data by Heirtzler et al (1968). Irving and Couillard (1973) updated Helsley and Steiner's (1969) and McElhinny and Burek's (1971) compilation of Cretaceous radiometrically or paleontologically dated geomagnetic field data. They confirmed the long Cretaceous normal interval by consistently finding normal polarity for the PTS interval -109 to -82 Ma (middle Aptian-end of Coniacian), the PTS intervals -82 to -65 Ma and -160 to -109 Ma being of mixed polarity. Later paleomagnetic work by Hanna (1973) on the radiometrically dated Boulder batholith and by Keating et al (1975) on biostratigraphically dated sediments confirmed the mixed-polarity nature for post-Santonian time. A different result was obtained by Shive and Frerichs (1974), who found no reversed polarity in their study of paleomagnetism in 350 samples of the micropaleontologically dated Coniacian-early Maastrichtian Niobrara Formation in Wyoming. This is no surprise for the lower Coniacian/early Santonian part of the formation, *Globotruncana concavata* (= *G. c. primitiva*?), considered early Santonian by Shive and Frerichs (1974) and Frerichs and Adams (1973), but it poses a problem for the upper part unless the late Campanian/early Maastrichtian age assignment is not correct and the entire Niobrara Formation is Coniacian/Santonian as had been suggested by Cobban and Reeside (1962). The latter may be the case, for microfossil evidence for a younger age is very weak. Furthermore, the Niobrara Formation is older than the Elkhorn Mountains volcanic rocks and the Boulder batholith (Klepper et al, 1957), both of which represent times of mixed polarity (Hanna, 1967, 1973, respectively); the younger Boulder batholith has been dated radiometrically as 78-68 Ma old (Hanna, 1973). This would make the base of the Maastrichtian older than 78 Ma, an unlikely high age.

OCEANIC BASEMENT AGES

Holes of the Deep Sea Drilling Project that reach basement of known magnetic-anomaly number offer an opportunity to age-calibrate the marine geomagnetic-reversal scale, and to provide a check on the consistency of the stratigraphic-radiometric time scale being used. In the following discussion, stratigraphic basement ages are estimated and expressed in numbers of the proposed time scale. These estimates are plotted in Figure 9 for ready comparison with the magnetic-reversal time scale. One age (Site 166) was used to calibrate the magnetic-reversal scale (Larson and Pitman, 1972; van Hinte, 1976); the others function as checks on the age assignment for the calibration point and its consequences for the geomagnetic-reversal scale. The figure demonstrates that none of the basement ages found by the drill contradict the age as it would have been predicted from the magnetic-anomaly position of the sites. This suggests that the present scale is a useful work model.

The following DSDP holes bottomed in Cretaceous oceanic basement of known linear magnetic-anomaly position: 10, 137, 138, 166, 239, 303, 304, 355, and 358 (nearly so). The estimates for the age of basement at Sites 166, 303, 304, and 307 (Late Jurassic) were discussed in van Hinte (1976).

DSDP Site 10 was drilled near anomaly 32. Campanian sediment (*Globotruncana stuartiformis* Zone) overlies basaltic basement at -456 m (Peterson et al, 1970; Cita and Gartner, 1971; Smali, 1973). The *Globotruncana calcarata* Zone is well defined in the section above. This zone is 20 m thick (-400 to -420 m); assuming that its duration is 1 Ma (-71 to -70 Ma), the average rate of sediment accumulation is 2 cm/1,000 years. Further, assuming that this rate is constant, it follows that the age at the base of sediment is $71 + 1.8 = 72.8$ Ma old. If the *G. calcarata* Zone represents not 1 but 2 Ma, the average rate of sediment accumulation then would be 1 cm/1,000 years and the lowest sediment would be 75.6 Ma old. An upward extrapolation of the two rates would predict the top of the Cretaceous at -300 and -350 m, respectively. The Cretaceous top was not cored, but recovery of the *Globotruncanella mayaroensis* Zone in Core 10 (-291 to -299 m) indicates that the average rate of sediment accumulation was even slightly higher than 2 cm/1,000 years and supports the assumption of a 1-Ma duration for the *G. calcarata* Zone. Even incorporating the possibilities that the rate of sediment accumulation is not constant and that basement is somewhat older than the oldest sedi-





FIG. 9—Cretaceous geomagnetic-reversal scales proposed by different authors. Linear vertical scale for each column follows original authors. No attempt was made to convert radiometric ages to PTS standards or to reinterpret biostratigraphic ages except for *This Paper* column. *Site 263 Mixed Zone* refers to mixed polarity found by Green and Brecher (1974) and Jarrard (1974) in cores from DSDP Site 263. See also caption of Figure 2. Cretaceous basement ages at DSDP sites are discussed in text.



ment, it seems reasonable to estimate basement to be between 72 and 75 Ma old.

Basement age at Site 137 can be estimated from planktonic foraminiferal biohorizons given by Beckmann (1972): *Planomalina buxtorfi* (–100 Ma) in Core 14 at –341.7 m and *Ticinella* (*Biticinella*) *breggiensis* (–102 Ma) in Core 16 at –381.7 m. An extrapolation of the average rate of sediment accumulation derived from these data (2 cm/1,000 years) dates the base of sediment (–397 m) as 103 Ma old. A fair estimate for the age of basement seems to be –103 to –106 Ma.

Only a broad paleontologic age has been determined for the lowest sediment at Site 138. Radiolaria of Core 6 (–425 to –431 m) compare with those of Core 7 at Site 137, which is at the oldest late Cenomanian and probably closer to 90 than to 95 Ma old. Petrushevskaya and Kozlova (1972) dated the fauna as Cenomanian but mentioned that it could be Albian as well. One estimate for the age of the lowest sediment could be –92 to –105 Ma. But, is this close to basement age? The fact that at Site 138 sediment above the basalt was deposited below calcite-compensation depth might suggest that at time of deposition the location was too deep for the basalt to be “normal” basement and that the hole bottomed in a sill (Berger and von Rad, 1972, p. 878; above the carbonate-clay boundary, or CCB, Berger and Winterer, 1974). Sites 137 and 138 were drilled at comparable levels with regard to basement topography, and we may assume that basement at 138 was, as at 137, originally somewhere near 2,700 m depth (Berger and von Rad, 1972, Fig. 57, Site 137). A bold extrapolation of Berger and von Rad's Site 138 curve (1972, Fig. 57) to 2,700 m suggests that basement is in the order of 110–115 Ma old (Berger and von Rad use a different time scale but have, in Fig. 57, Core 6 at –92 Ma with which we would agree).

DSDP Site 239 was drilled on the older side of anomaly 31 (Schlich, 1974) and bottomed in basaltic basement overlain by less than a meter of Cretaceous sediment that in turn was covered by Paleocene. No Cretaceous planktonic Foraminifera were found, but the benthic fauna and the calcareous nannoflora suggest a Maastrichtian-latest Campanian age (not late Campanian exclusive of Maastrichtian as assumed by Schlich, 1974, and by Larson and Pitman, 1975). Although a late Maastrichtian basement age (–65 to –68 Ma) seems most likely, the writer plotted Site 239 as –65 to –72 Ma in Figure 9.

Sites 355 and 358 were drilled on anomalies 33 and 32, respectively, and sediment above basement was dated as early-middle Campanian and as late Campanian (Anon., 1975). Basement ages

for these sites can be plotted as –78 to –72 Ma and –72 to –70 Ma, respectively.

REFERENCES CITED

- Afanasyev, G. C., 1970, Certain key data for the Phanerozoic time scale: *Eclogae Geol. Helvetiae*, v. 63, p. 1-7.
- et al, 1964, The project of a revised geological time scale in absolute chronology: 22d Internat. Geol. Cong., India, Soviet Contrib. Geol., p. 287-324.
- Afanasyev, S. L., et al, 1972, Duration of the ages of the Late Cretaceous: Moscow Soc. Naturalists Bull., Geol. Ser., v. 42, p. 136-137.
- Allemann, F., R. Catalano, F. Fares, and J. Remane, 1971, Standard calpionellid zonation (upper Tithonian-Valanginian) of the western Mediterranean province: 2d Internat. Conf. Planktonic Microfossils, Rome 1970, Proc., v. 2, p. 1337-1340.
- Anonymous, 1963, Conclusions du colloque de stratigraphie sur le Cretace Inferieur en France, Lyon, Septembre 1963: Soc. Geol. France Compte Rendu, no. 8, p. 292-296.
- 1975, Leg 39 examines facies changes in South Atlantic: *Geotimes*, v. 20, no. 3, p. 26-28.
- Baldwin, B., P. J. Coney, and W. R. Dickinson, 1974, Dilemma of a Cretaceous time scale and rates of seafloor spreading: *Geology*, v. 2, p. 267-270.
- Bandy, O. L., 1967, Cretaceous planktonic foraminiferal zonation: *Micropaleontology*, v. 13, p. 1-31.
- Barr, F. T., 1962, Upper Cretaceous planktonic Foraminifera from the Isle of Wight, England: *Palaeontology*, v. 4, p. 552-580.
- 1966, The foraminiferal genus *Bolivinoidea* from the Upper Cretaceous of the British Isles: *Palaeontology*, v. 9, p. 220-243.
- 1970, The foraminiferal genus *Bolivinoidea* from the Upper Cretaceous of Libya: *Jour. Paleontology*, v. 44, p. 642-654.
- 1972, Cretaceous biostratigraphy and planktonic Foraminifera of Libya: *Micropaleontology*, v. 18, p. 1-46.
- Bartenstein, H., 1974, *Lenticulina* (*Lenticulina*) *nodosa* (Reuss 1863) and its subspecies—worldwide index Foraminifera in the Lower Cretaceous: *Eclogae Geol. Helvetiae*, v. 67, p. 539-562.
- and F. Bettenstaedt, 1962, Marine Unterkreide (Boreal und Tethys), in *Leitfossilien der Mikropalaeontologie*: Berlin, Borntraeger, p. 225-297.
- Beckmann, J. P., 1972, The Foraminifera and some associated microfossils of Sites 135 to 144: Deep Sea Drilling Project Initial Repts., v. 14, p. 389-420.
- Berger, W. H., and U. von Rad, 1972, Cretaceous and Cenozoic sediments from the Atlantic Ocean: Deep Sea Drilling Project Initial Repts., v. 14, p. 787-954.
- and E. L. Winterer, 1974, Plate stratigraphy and the fluctuating carbonate line: Internat. Assoc. Sedi-mentol. Spec. Pub. 1, p. 11-45.
- Berggren, W. A., 1972, A Cenozoic time scale—some implications for regional geology and paleobiogeography: *Lethaia*, v. 5, p. 195-215.
- D. P. McKenzie, J. G. Sclater, and J. E. van Hinte, 1975, Worldwide correlation of Mesozoic magnetic anomalies and its implications, discussion:

- Geol. Soc. America Bull., v. 86, p. 267-269.
- Bodelle, J., C. Lay, and G. S. Odin, 1969, Determination d'âge par la méthode géochronologique "potassium-argon" de glauconies du bassin de Paris: *Acad. Sci. Comptes Rendus*, ser. D, v. 268, p. 1474-1477.
- and A. Parfenoff, 1969, Âge des glauconies crétacées du sud-est de la France (2, Vallée de l'Esteron, Alpes-Maritimes; résultats préliminaires de la méthode potassium-argon: *Acad. Sci. Comptes Rendus*, ser. D, v. 268, p. 1576-1579.
- Bolli, H. M., 1966, Zonation of Cretaceous to Pliocene marine sediments based on planktonic Foraminifera: *Asoc. Venezolana Geologia, Minería y Petróleo Bol. Inf.*, v. 9, p. 3-32.
- Bonhomme, M., G. S. Odin, and C. Pomerol, 1968, Age de formations glauconieuses de l'Albien et de l'Eocène de bassin de Paris: *France, Bur. Recherches Geol. et Minières Mem.* 58, p. 339-346.
- Bukry, D., 1974, Coccolith stratigraphy, offshore Western Australia, Deep-Sea Drilling Project leg 27: *Deep Sea Drilling Project Initial Repts.*, v. 27, p. 623-630.
- Caldwell, W. G. E., 1968, The Late Cretaceous Bearpaw Formation in the South Saskatchewan River Valley: *Saskatchewan Research Council Geology Div. Rept.* 8, 86 p.
- Casey, R., 1961, The stratigraphical palaeontology of the Lower Greensand: *Palaeontology*, v. 3, p. 487-621.
- 1964, The Cretaceous Period, in *The Phanerozoic time-scale*: *Geol. Soc. London Quart. Jour.*, v. 120, supp., p. 193-202.
- Cita, M. B., and S. Gartner, Jr., 1971, Deep sea Upper Cretaceous from the western North Atlantic: 2d Planktonic Conf., Rome 1970, *Proc.*, v. 1, p. 287-320.
- Cobban, W. A., and J. B. Reeside, 1962, Correlation of the Cretaceous formations of the Western Interior of the United States: *Geol. Soc. America Bull.*, v. 63, p. 1011-1044.
- Colin, J.-P., 1974, Précisions sur le Campanien de Dordogne (région de Belvès-Saint-Cyprien, Dordogne, S-O France): *Newsletters Stratigraphy*, v. 3, p. 139-151.
- Cotillon, P., 1971, Le Crétacé Inférieur de l'Arc subalpin de Castellane entre l'Asse et le Var stratigraphie et sédimentologie: *Bur. Recherches Geol. et Minières Mem.* 68, 243 p.
- Creer, K. M., 1971, Mesozoic palaeomagnetic reversal column: *Nature*, v. 233, p. 545-546.
- Dalbiez, F., 1959, Correlations et résolutions: 84th Cong. Soc. Sav. Paris, Dijon, Sec. Sci., *Comptes Rendus Colloque Sur le Crétacé Supérieur Français*, p. 857-867.
- Dickinson, W. R., and E. I. Rich, 1972, Petrologic intervals and petrofacies in the Great Valley sequence, Sacramento Valley, California: *Geol. Soc. America Bull.*, v. 83, p. 3007-3024.
- Donze, P., B. Porthault, G. Thomel, and O. de Villoutreys, 1970, Le Senonien Inférieur de Puget-Théniers (Alpes-Maritimes) et sa microfaune: *Geobios*, no. 3, p. 41-103.
- Ernst, G., 1970, The stratigraphical value of the echinoids in the boreal Upper Cretaceous: *Newsletters Stratigraphy*, v. 1, p. 19-34.
- Evernden, J. F., and R. W. Kistler, 1970, Chronology of emplacement of Mesozoic batholithic complexes in California and western Nevada: *U.S. Geol. Survey Prof. Paper* 623, 42 p.
- Folinsbee, R. E., H. Baadsgaard, G. L. Cumming, J. Nascimbene, and M. Shafiqullah, 1965, Late Cretaceous radiometric dates from the Cypress Hills of western Canada, in *Cypress Hills Plateau, Alberta and Saskatchewan*: *Alberta Soc. Petroleum Geology, 15th Field Conf. Guidebook*, p. 162-174.
- Frerichs, W. E., and P. R. Adams, 1973, Correlation of the Hilliard Formation with the Niobrara Formation: *Wyoming Geol. Assoc. Guidebook, 25th Field Conf.*, p. 187-192.
- Geological Society of London, 1964, Geological Society Phanerozoic time-scale 1964, in *The Phanerozoic time-scale*: *Geol. Soc. London Quart. Jour.*, v. 120, supp., p. 260-262.
- Ghosh, P. K., 1972, Use of bentonites and glauconites in potassium-40/argon-40 dating in Gulf Coast stratigraphy: *PhD thesis, Rice Univ.*, 136 p.
- Gill, J. R., and W. A. Cobban, 1966, The Red Bird section of the Upper Cretaceous Pierre Shale in Wyoming: *U.S. Geol. Survey Prof. Paper* 393-A, 69 p.
- Green, K. E., and A. Brecher, 1974, Preliminary paleomagnetic results for sediments from Site 263, leg 27: *Deep Sea Drilling Project Initial Repts.*, v. 27, p. 405-413.
- and J. Sclater, 1974, Geomagnetic field reversals in the Cretaceous from DSDP sediments (abs.): *EOS (Am. Geophys. Union Trans.)*, v. 55, p. 236.
- Hanna, W. F., 1967, Paleomagnetism of Upper Cretaceous volcanic rocks of southwestern Montana: *Jour. Geophys. Research*, v. 72, p. 595-610.
- 1973, Paleomagnetism of the Late Cretaceous Boulder batholith: *Am. Jour. Sci.*, v. 273, p. 778-802.
- Harland, W. B., and E. H. Francis, eds., 1971, *The Phanerozoic time-scale; a supplement*: *Geol. Soc. London Spec. Pub.* 5, 356 p.
- Heirtzler, J. R., G. O. Dickson, E. M. Herron, W. C. Pitman, III, and X. LePichon, 1968, Marine magnetic anomalies, geomagnetic field reversals, and motions of the ocean floor and continents: *Jour. Geophys. Research*, v. 73, p. 2119-2136.
- Helsley, C. E., 1973, Post-Paleozoic magnetic reversals (abs.): *Geol. Soc. America Abs. with Programs*, v. 5, p. 665 (hand-out figure "Comparison of Mesozoic reversal sequences").
- and M. B. Steiner, 1968, Evidence for long periods of normal magnetic polarity in the Cretaceous Period (abs.): *Geol. Soc. America Abs. with Programs*, p. 133.
- 1969, Evidence for long intervals of normal polarity during the Cretaceous Period: *Earth and Planetary Sci. Letters*, v. 5, p. 325-332.
- Herm, D., 1965, Mikropalaeontologisch-stratigraphische Untersuchungen im Kreideflysch zwischen Deva und Zumaya (Prov. Guipuzcoa, Nordspanien): *Deutsch. Geol. Gesell. Zeitschr.*, v. 115 (1963), p. 277-348.
- Hermes, J. J., 1969, Late Albian Foraminifera from the Subbetic of southern Spain: *Geologie en Mijnbouw*, v. 48, p. 35-66.

- Hiltermann, H., 1963, Zur Entwicklung der Benthos-Foraminifere *Bolivinoidea*, in Koenigswald et al, eds., Evolutionary trends in Foraminifera: New York, Elsevier Pub. Co., p. 198-223.
- and W. Koch, 1962, Oberkreide des nordlichen Mitteleuropa, in Leitfossilien der Mikropalaeontologie: Berlin, Borntraeger, p. 299-338.
- Holmes, A., 1959, A revised geological time-scale: Edinburgh Geol. Soc. Trans., v. 17, p. 183-216.
- Irving, E., and R. W. Couillard, 1973, Cretaceous normal polarity interval: Nature, Phys. Sci., v. 244, p. 10-11.
- International Union of Geological Sciences (IUGS), 1967, A comparative table of recently published geological time scales for the Phanerozoic time—explanatory notice: Norsk Geol. Tidsskr., v. 47, p. 375-380.
- Jarrard, R. D., 1974, Paleomagnetism of some leg 27 sediment cores: Deep Sea Drilling Project Initial Repts., v. 27, p. 415-423.
- Juignet, P., J. C. Hunziker, and G. S. Odin, 1975, Dation numerique du passage Albien-Cenomanien en Normandie; etude preliminaire par la methode a l'argon: Acad. Sci. Comptes Rendus, ser. D., v. 280, p. 379-382.
- Kauffman, E. G., 1970, Population systematics, radio-metrics and zonation—a new biostratigraphy: North Am. Paleont. Conv. Proc., p. 612-666.
- Keating, B., C. E. Helsley, and E. A. Pessagno, 1975, Late Cretaceous reversal sequence: Geology, v. 3, p. 73-76.
- Kemper, E., 1971, Zur Gliederung und Abgrenzung des norddeutschen Aptium mit Ammoniten: Geol. Jahrb., v. 89, p. 359-390.
- 1973a, The Valanginian and Hauterivian Stages in northwest Germany, in The boreal Lower Cretaceous: Liverpool, England, Seel House Press, p. 327-344.
- 1973b, The Aptian and Albian Stages in northwest Germany, in The boreal Lower Cretaceous: Liverpool, England, Seel House Press, p. 345-360.
- P. F. Rawson, Fr. Schmid, and C. Spaeth, 1974, Die Megafauna der Kreide von Helgoland und ihre biostratigraphische Deutung: Newsletters Stratigraphy, v. 3, p. 121-137.
- Kennedy, W. J., 1969, The correlation of the lower Chalk of south-east England: Geologists' Assoc. Proc., v. 80, p. 459-560.
- 1970, A correlation of the uppermost Albian and the Cenomanian of southwest England: Geologists' Assoc. Proc., v. 81, p. 613-677.
- and J. M. Hancock, 1970, Ammonites of the genus *Acanthoceras* from the Cenomanian of Rouen, France: Palaeontology, v. 13, p. 462-490.
- and P. Juignet, 1973, Observations on the lithostratigraphy and ammonite succession across the Cenomanian-Turonian boundary in the environs of Le Mans (Sarthe, N.W. France): Newsletters Stratigraphy, v. 2, p. 189-202.
- Klepper, M. R., R. A. Weeks, and E. T. Ruppel, 1957, Geology of the southern Elkhorn Mountains, Jefferson and Broadwater Counties, Montana: U.S. Geol. Survey Prof. Paper 292, 82 p.
- Kuhri, B., 1970, Some observations on the type material of *Globotruncana elevata* (Brotzen) and *Globotruncana concavata* (Brotzen): Rev. Espanola Micropaleontologia, v. 2, p. 291-304.
- 1971, Lower Cretaceous planktonic Foraminifera from the Miravetes, Argos and Represa Formations (S.E. Spain): Rev. Espanola Micropaleontologia, v. 3, p. 219-237.
- 1972, Stratigraphy and micropaleontology of the Lower Cretaceous in the subbetic south of Cravaca (province of Murcia, S.E. Spain): Koninkl. Nederlandse Akad. Wetensch. Proc., ser. B, v. 75, p. 193-222.
- Kulp, J. L., 1960, The geological time scale: 21st Internat. Geol. Cong., Copenhagen, Rept., pt. 3, p. 18-27.
- 1961, Geologic time scale: Science, v. 133, p. 1105-1114.
- Lambert, R. St. J., 1971a, The pre-Pleistocene Phanerozoic time-scale; a review, in The Phanerozoic time-scale; a supplement: Geol. Soc. London Spec. Pub. 5, p. 9-31.
- 1971b, The pre-Pleistocene Phanerozoic time-scale; further data, in The Phanerozoic time-scale; a supplement: Geol. Soc. London Spec. Pub. 5, p. 33-34.
- Larson, R. L., and W. C. Pitman, III, 1972, World-wide correlation of Mesozoic magnetic anomalies, and its implications: discussion: Geol. Soc. America Bull., v. 83, p. 3645-3661.
- 1975, World-wide correlation of Mesozoic magnetic anomalies and its implications: reply: Geol. Soc. America Bull., v. 86, p. 270-272.
- Magné, J., and J. Polvéche, 1961, Sur le niveau a *Actinocamax plenus* (Blainville) du Boulonnais: Soc. Geol. Nord Annales, v. 81, p. 47-62.
- Maync, W., 1959, Foraminiferal key biozones in the Lower Cretaceous of the western hemisphere and the Tethys province, in El sistema Cretacio, v. 1: 20th Internat. Geol. Cong., Mexico, p. 85-111.
- McElhinny, M. W., 1973, Palaeomagnetism and plate tectonics: Cambridge, England, Cambridge Univ. Press, 358 p.
- and P. J. Burek, 1971, Mesozoic palaeomagnetic stratigraphy: Nature, v. 232, p. 98-102.
- McKenzie, D., and J. G. Sclater, 1971, The evolution of the Indian Ocean since the Late Cretaceous: Royal Astron. Soc. Geophys. Jour., v. 25, p. 437-528.
- Michael, E., 1966, Die Evolution der Gavelinelliden (Foram.) in der NW-deutschen Unterkreide: Senckenbergiana Lethaea, v. 47, p. 411-459.
- Moullade, M., 1966, Etude stratigraphique et micropaleontologique du Cretace Inferieur de la "fosse vocontienne": Lyons Univ. Fac. Sci. Lab. Geol. Doc. 15, 369 p.
- Muller, S. W., and H. G. Schenck, 1943, Standard of Cretaceous system: AAPG Bull., v. 27, p. 262-278.
- North, B. R., and W. G. A. Caldwell, 1970, Foraminifera from the Late Cretaceous Bearpaw Formation in the south Saskatchewan River Valley: Saskatchewan Research Council Geology Div. Rept. 9, 117 p.
- Obradovich, J. D., and W. A. Cobban, 1975, A time scale for the Late Cretaceous of the Western Interior of North America: Geol. Assoc. Canada Spec. Paper No. 13, p. 31-54.

- Olsson, R. K., 1964, Late Cretaceous planktonic Foraminifera from New Jersey and Delaware: *Micropalaeontology*, v. 10, p. 157-188.
- Owen, H. G., 1973, Ammonite faunal provinces in the middle and upper Albian and their palaeogeographical significance, in *The boreal Lower Cretaceous*: Liverpool, England, Seel House Press, p. 145-154.
- Owens, J. P., and N. F. Sohl, 1973, Glauconites from New Jersey-Maryland coastal plain; their K-Ar ages and application in stratigraphic studies: *Geol. Soc. America Bull.*, v. 84, p. 2811-2838.
- Papp, A., 1956, Die morphologisch-genetische Entwicklung von Orbitoiden und ihre stratigraphische Bedeutung im Senon: *Palaont. Zeitschr.*, v. 30, p. 45-49.
- Pechersky, D. M., and A. N. Khramov, 1973, Mesozoic palaeomagnetic scale of the U.S.S.R.: *Nature*, v. 244, p. 499-501.
- Pergament, M. A., 1967, Stages in *Inoceramus* evolution in the light of absolute geochronology: *Paleont. Jour.* no. 1, p. 27-34 (transl.).
- Pessagno, E. A., Jr., 1969, Upper Cretaceous stratigraphy of the western Gulf Coast area of Mexico, Texas and Arkansas: *Geol. Soc. America Mem.* 111, 139 p.
- Peterson, M. N. A., N. T. Edgar, M. B. Cita, S. Gartner, R. Goll, C. Nigrini, and C. von der Borch, 1970, Site 10: Deep Sea Drilling Project Initial Repts., v. 2, 501 p.
- Petrushevskaya, M. G., and G. E. Kozlova, 1972, Radiolaria; Leg 14, deep sea drilling project: Deep Sea Drilling Project Initial Repts., v. 14, p. 495-648.
- Porthault, B., G. Thomel, and O. de Villoutreys, 1966, Etude biostratigraphique du Cenomanien du bassin superieur de l'Esteron (Alpes-Maritimes); le probleme de la limite Cenomanien-Turonien dans le sud-est de la France: *Soc. Geol. France Bull.*, ser. 7, v. 8, p. 423-439.
- Premoli Silva, I., and H. M. Bolli, 1973, Late Cretaceous to Eocene planktonic Foraminifera and stratigraphy of Leg 15 Sites in the Caribbean Sea: Deep Sea Drilling Project Initial Repts., v. 15, p. 499-547.
- Proto Decima, F., 1974, Leg 27 calcareous nannoplankton: Deep Sea Drilling Project Initial Repts., v. 27, p. 589-621.
- Reiss, Z., 1962, Stratigraphy of phosphate deposits in Israel: *Israel Geol. Survey Bull.* 34, p. 1-23.
- Riedel, W. R., and A. Sanfilippo, 1975, Radiolaria from the southern Indian Ocean, DSDP leg 26: Deep Sea Drilling Project Initial Repts., v. 26, p. 771-813.
- Risch, H., 1971, Stratigraphie der höheren Unterkreide der bayerischen Kalkalpen mit Hilfe von Mikrofossilien: *Palaontographica*, v. 138A, p. 1-80.
- Robaszyński, F., 1971, Les foraminifères pelagiques des "Dives" cretacees aux abords du golfe de Mons (Belgique): *Soc. Geol. Nord Annales*, v. 41, p. 31-38.
- Roth, P. H., 1973, Calcareous nannofossils Leg 17 Deep Sea Drilling Project: Deep Sea Drilling Project Initial Repts., v. 17, p. 695-796.
- Rubinshteyn, M. M., 1967, The argon method as applied to some problems of regional geology: *Akad. Nauk Gruzin. SSR, Inst. Geol. Trudy*, n.s., no. 11, 239 p. (in Russian; English summ.).
- Salaj, J., and S. Bajanik, 1972, Contribution a la stratigraphie du Cretace et du Paleogene de la region de l'Oued Zarga: *Tunisia Serv. Geol. Notes*, no. 38, p. 63-71.
- Scheibnerova, V., 1972, Some new views on Cretaceous biostratigraphy based on the concept of foraminiferal biogeoprovinces: *New South Wales Geol. Survey Rec.*, v. 14, p. 85-87.
- Schlich, R., 1974, Seafloor spreading history and deep-sea drilling results in the Madagascar and Mascarene basins, western Indian Ocean: Deep Sea Drilling Project Initial Repts., v. 25, p. 663-678.
- Sclater, J. G., R. D. Jarrard, B. McGowran, and S. Gartner, 1974, Comparison of the magnetic and biostratigraphic time scales since the Late Cretaceous: Deep Sea Drilling Project Initial Repts., v. 22, p. 381-386.
- Seitz, O., 1956, Über Ontogenie, Variabilität und Biostratigraphie einiger Inoceramen: *Palaont. Zeitschr.*, v. 30, Sonderh., p. 3-6.
- Séronie-Vivien, M., 1972, Contribution a l'étude de Senonien en Aquitaine septentrionale, ses stratotypes: Coniacien, Santonien, Campanien: France, Centre Natl. Recherche Sci. Editions, 195 p.
- Shive, P. N., and W. E. Frerichs, 1974, Paleomagnetism of the Niobrara Formation in Wyoming, Colorado, and Kansas: *Jour. Geophys. Research*, v. 79, p. 3001-3007.
- Sigal, J., 1965, Etat des connaissances sur les foraminifères du Cretace Inferieur: *Bur. Recherches Geol. et Minieres Mem.* 34, p. 489-502.
- 1966, Contribution a une monographie des rosalines, pt. 1, le genre *Ticinella* Reichel, souche des rotalipores: *Eclogae Geol. Helvetiae*, v. 59, p. 185-217.
- 1967, Essai sur l'état actuel d'une zonation stratigraphique a l'aide des principales especes de rosalines (foraminifères): *Soc. Geol. France Compte Rendu*, p. 48-50.
- Silver, L. T., C. R. Allen, and F. G. Stehli, 1969, Geological and geochronological observations on a portion of the Peninsular Range batholith of north-western Baja California, Mexico (abs.): *Geol. Soc. America Abs. with Programs*, p. 279-280.
- Smali, M., 1973, Foraminiferi Campaniano-Maastrichtiani della dorsale Medio-Atlantica: *Riv. Italiana Paleontologia e Stratigrafia*, v. 79, p. 35-100.
- Smith, C. C., and E. A. Pessagno, 1973, Planktonic Foraminifera and stratigraphy of the Corsicana Formation (Maestrichtian), north-central Texas: *Cushman Found. Forum. Research Spec. Pub.* 12, 68 p.
- Sornay, J., 1968, Termes stratigraphiques majeurs; Aptien, in *Lexique stratigraphique international*, v. 8: Paris, Union Internat. Sci. Geol., Comm. Stratigr., 109 p.
- Suppe, J., 1969, Times of metamorphism in the Franciscan terrain of the northern coast ranges, California: *Geol. Soc. America Bull.*, v. 80, p. 135-142.
- 1973, Geology of the Leech Lake Mountain; Ball Mountain region, California: *California Univ. Pubs. Geol. Sci.*, v. 107, 82 p.
- Thierstein, H. R., 1973, Lower Cretaceous calcareous nannoplankton biostratigraphy: [Austria] *Geol. Bundesanst. Abh.*, v. 29, 52 p.
- Thieuloy, J.-P., 1973, The occurrence and distribution of boreal ammonites from the Neocomian of south-

- east France (Tethyan province), in *The boreal Lower Cretaceous*: Liverpool, England, Seel House Press, p. 289-302.
- Thomel, G., 1972, Les Acanthoceratidae cenomaniens des chaines subalpines meridionales: *Soc. Geol. France Mem.* 116, 204 p.
- 1973a, A propos de la limite entre les etages Cenomanien et Turonien: *Acad. Sci. Comptes Rendus*, ser. D, v. 277, p. 761-764.
- 1973b, A propos de la zone a *Actinocamax plenus*: principe et application de la methodologie biostratigraphique: *Nice Museum d'Histoire Nat. Annales*, v. 1, supp. h.s., p. 1-28.
- Troeger, K. A., 1968, Zur Bedeutung okologischer Faktoren fur die Leitfossilien des Obercenoman-Unterturon Zeitabschnitts in Mitteleuropa: *Geologie*, v. 17, p. 68-75.
- Vail, P. R., R. M. Mitchum, and S. Thompson, 1974, Eustatic cycles based on sequences with coastal onlap (abs.): *Geol. Soc. America Abs. with Programs*, v. 6, p. 993.
- Van Gorsel, J. T., 1973a, The type Campanian and the Campanian-Maastrichtian boundary in Europe: *Geologie en Mijnbouw*, v. 52, p. 141-146.
- 1973b, *Helicorbitoides* from southern Sweden and the origin of the *Helicorbitoides-Lepidorbitoides* lineage: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 76, p. 273-286.
- 1974, Some complex Upper Cretaceous rotaliid Foraminifera from the northern border of the Aquitaine basin (SW France): *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 77, p. 319-339.
- van Hinte, J.E., 1963, Zur Stratigraphie und Mikropalaontologie der Oberkreide und des Eozans des Krappfeldes (Karnten): *Austria, Geol. Bundesanst. Jahrb., Sonderb.* 8, 147 p.
- 1965, The type Campanian and its planktonic Foraminifera: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 68, p. 8-28.
- 1966a, *Orbitoides* from the Campanian type section: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 69, p. 79-110.
- 1966b, *Orbitoides hottingeri* n. sp. from northern Spain: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 69, p. 387-402.
- 1967, *Bolivinoidea* from the Campanian type section: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 70, p. 254-263.
- 1968, The Late Cretaceous larger foraminifer *Orbitoides douvillei* (Silvestri) at its type locality Belves, SW France: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 71, p. 359-372.
- 1969a, A *Globotruncana* zonation of the Senonian subseries: 1st Internat. Conf. Planktonic Microfossils, Geneva 1967, *Proc.*, v. 2, p. 257-266.
- 1969b, The nature of biostratigraphic zones: 1st Internat. Conf. Planktonic Microfossils, Geneva 1967, *Proc.*, v. 2, p. 267-272.
- 1972, The Cretaceous time scale and planktonic-foraminiferal zones: *Koninkl. Nederlandse Akad. Wetensch. Proc.*, ser. B, v. 75, p. 1-8.
- 1976, A Jurassic time scale: *AAPG Bull.*, v. 60, p. 489-497.
- J. A. S. Adams, and D. Perry, 1975, K/Ar age of Lower-Upper Cretaceous boundary at Orphan Knoll (Labrador Sea): *Canadian Jour. Earth Sci.*, v. 12, p. 1484-1491.
- Vermeulen, J., 1974, Sur une biostratigraphie homophyletique basee sur la famille des Pulchellidae: *Acad. Sci. Comptes Rendus*, ser. D, v. 278, p. 2885-2887.
- Watkins, N. D., and F. W. Cambray, 1971, Paleomagnetism of Cretaceous dikes from Jamaica: *Royal Astron. Soc. Geophys. Jour.*, v. 22, p. 163-179.