ii. *Suspicious morphology* ☐ Milk of calcium ☐ Suture ☐ Amorphous

☐ Coarse heterogenous ☐ Fine pleomorphic ☐ Fine linear or linear branching

iii. Distribution: ☐ Diffuse ☐ Regional ☐ Grouped ☐ Linear ☐ Segmental

g. Architectural distortion: ☐ Right ☐ Left ☐ Both ☐ Not present

h. Asymmetry: ☐ Global ☐ Focal ☐ Developing ☐ Not present

i. Intra-mammary LN: ☐ Right ☐ Left _____

j. Skin lesion ☐ Right ☐ Left _____

k. Solitary dilated duct ☐ Right ☐ Left _____

l. Associated features

  i. Skin retraction ☐ Right ☐ Left _____

  ii. Nipple retraction ☐ Right ☐ Left _____

  iii. Skin thickening ☐ Right ☐ Left _____

  iv. Trabecular thickening ☐ Right ☐ Left _____

  v. Axillary lymphadenopathy ☐ Right ☐ Left _____

  vi. Architectural distortion ☐ Right ☐ Left _____

  vii. Calcification ☐ Right ☐ Left _____

m. 2nd lesion/contralateral breast: _____
_____
_____

n. Distance from skin: ☐ <0.5 cm ☐ >0.5 cm

o. Distance from Pectoralis Major: _____ cm

p. BI-RADS: ☐ 0 ☐ I ☐ II ☐ III ☐ IV ☐ V

87. **3-D Digital Tomosynthesis:** ☐ Done ☐ Not done Date:_____ Number:_____

a. Location of lesion: ☐ Right ☐ Left ☐ Both

☐ UOQ ☐ UIQ ☐ LIQ ☐ LOQ ☐ CQ Position:_____ o clock

b. Size: ☐ <2cm ☐ 2-5 cm ☐ >5cm

c. Distance from skin: ☐ <0.5 cm ☐ >0.5 cm

d. Distance from Pectoralis Major: _____ cm

e. Diagnosis: ☐ Normal ☐ Benign ☐ Suspicious ☐ Diagnostic for cancer

88. **ABVS:** ☐ Done ☐ Not done Date:_____ Number:_____

a. Location of lesion: ☐ Right ☐ Left ☐ Both

☐ UOQ ☐ UIQ ☐ LIQ ☐ LOQ ☐ CQ Position:_____ o clock

b. Size: ☐ <2cm ☐ 2-5 cm ☐ >5cm

c. Distance from skin: ☐ <0.5 cm ☐ >0.5 cm

d. Distance from Pectoralis Major: _____ cm

e. Diagnosis: ☐ Normal ☐ Benign ☐ Suspicious ☐ Diagnostic for cancer