



Grant Medical Foundation
Ruby Hall Clinic
Wanowarie



Operation Details :

- Methylene blue dye injected in periareolar region.
- Under GA, B/L breast prepped & draped in sterile fashion.
- Left breast local infiltration administered in circumareolar fashion.
- Circumareolar skin incision taken and strip of skin around this incision deepithelised.
- Dissection started in upper outer quadrant to reach for left sentinel node based on gamma probe and tracking of blue dye.
- Suspicious node identified & excised.
- Through the same incision, suspicious breast mass also excised with adequate margins.
- Axilla closed in layers.
- Circumareolar skin incision closed in layers.
- Romovac mini drain placed before closure.

Major Post op events :

Uneventful.

Course in the Hospital including complications if any

Course in the Hospital

Patient admitted with the above complaints & history.
All pre-op profile done.
Pre-anaesthetic check-up done.
Patient underwent LEFT BREAST CONSERVATION SURGERY + SENTINEL NODE EXCISION under GA on 14/02/17.
Procedure uneventful.
Post-op shifted to ward, managed with IVF, antibiotic, analgesic and other supportive management.
Vitals monitored.
Discharged with drain.
Patient haemodynamically stable at the time of discharge.

Treatment during Hospitalisation

IV Fluids : RL

Tablets	: T. CEFTUM 500MG	1--0--1
.	: T. VOVERAN SR 75MG	1--0--1
.	: C. PAN-D	1--0--1
.	: -----	

Advice on Discharge

Treatment on Discharge

P.T.O