

- 8 Fig. 8.1 shows the variation with nucleon number of the binding energy per nucleon of a nucleus.





Fig. 8.1

- (a) On Fig. 8.1, mark with the letter S the position of the nucleus with the greatest stability. [1]

- (b) One possible fission reaction is



- (i) On Fig. 8.1, mark possible positions for

1. the Uranium-235 (${}_{92}^{235}\text{U}$) nucleus (label this position U),
2. the Krypton-90 (${}_{36}^{90}\text{Kr}$) nucleus (label this position Kr).

[1]

- (ii) The binding energy per nucleon of each nucleus is as follows.

$$\begin{aligned} {}_{92}^{235}\text{U}: & \quad 1.2191 \times 10^{-12} \text{ J} \\ {}_{56}^{144}\text{Ba}: & \quad 1.3341 \times 10^{-12} \text{ J} \\ {}_{36}^{90}\text{Kr}: & \quad 1.3864 \times 10^{-12} \text{ J} \end{aligned}$$

Use these data to calculate

1. the energy release in this fission reaction (give your answer to three significant figures),

energy = J [3]

2. the mass equivalent of this energy.

mass = kg [2]

- (iii) Suggest why the neutrons were not included in your calculation in (ii).

.....
..... [1]

Copyright Acknowledgements:

Every reasonable effort has been made to trace all copyright holders. The publishers will be pleased to hear from anyone whose rights we have unwittingly infringed.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.