


## Section A

Answer **all** the questions in the spaces provided.

- 1** A spherical planet has mass  $M$  and radius  $R$ .

The planet may be assumed to be isolated in space and to have its mass concentrated at its centre.

The planet spins on its axis with angular speed  $\omega$ , as illustrated in Fig. 1.1.



**Fig. 1.1**

A small object of mass  $m$  rests on the equator of the planet. The surface of the planet exerts a normal reaction force on the mass.

- (a)** State formulae, in terms of  $M$ ,  $m$ ,  $R$  and  $\omega$ , for

- (i) the gravitational force between the planet and the object,

..... [1]

- (ii) the centripetal force required for circular motion of the small mass,

..... [1]

- (iii) the normal reaction exerted by the planet on the mass.

..... [1]

- (b) (i)** Explain why the normal reaction on the mass will have different values at the equator and at the poles.

.....

.....

..... [2]

- (ii) The radius of the planet is  $6.4 \times 10^6$  m. It completes one revolution in  $8.6 \times 10^4$  s.  
Calculate the magnitude of the centripetal acceleration at

1. the equator,

$$\text{acceleration} = \dots \text{ms}^{-2} [2]$$

2. one of the poles.

$$\text{acceleration} = \dots \text{ms}^{-2} [1]$$

- (c) Suggest two factors that could, in the case of a real planet, cause variations in the acceleration of free fall at its surface.

1. ....

.....

2. ....

.....

[2]