


- 2 A ball of mass  $0.030\text{ kg}$  moves along a curved track, as shown in Fig. 2.1.



**Fig. 2.1**

The speed of the ball is  $1.3\text{ ms}^{-1}$  when it is at point A at a height of  $0.31\text{ m}$ . The ball moves down the track and collides with a vertical wall at point B. The ball then rebounds back up the track. It may be assumed that frictional forces are negligible.

- (a) Calculate the change in gravitational potential energy of the ball in moving from point A to point B.

change in gravitational potential energy = ..... J [2]

- (b) Show that the ball hits the wall at B with a speed of  $2.8\text{ ms}^{-1}$ .

[3]

- (c) The change in momentum of the ball due to the collision with the wall is  $0.096 \text{ kg ms}^{-1}$ . The ball is in contact with the wall for a time of 20 ms.

Determine, for the ball colliding with the wall,

- (i) the speed immediately after the collision,

speed = .....  $\text{ms}^{-1}$  [2]

- (ii) the magnitude of the average force on the ball.

force = ..... N [2]

- (d) State and explain whether the collision is elastic or inelastic.

.....  
 ..... [1]

- (e) In practice, frictional effects are significant so that the actual increase in kinetic energy of the ball in moving from A to B is 76 mJ. The length of the track between A and B is 0.60 m.

Use your answer in (a) to determine the average frictional force acting on the ball as it moves from A to B.

frictional force = ..... N [2]

[Total: 12]