



**1**

An object A of mass 9.0 kg and object B of mass 1.0 kg travel towards each other along a smooth horizontal surface in a straight line and collide head-on. The initial speeds of object A and B before the collision are  $u$  and  $3u$  respectively.

In Fig. 1.1, the variation with time  $t$  of momentum  $p$  is shown from  $t = 0$  to 3.0 s for particle A and from  $t = 0$  to 1.0 s for particle B.




Fig. 1.1

(a)

(i)

Momentum is conserved when two objects collide or interact. State the condition under which momentum is conserved.

.....

[1]

(ii)

On Fig. 1.1, draw the variation with  $t$  of  $p$  from 1.0 s to 3.0 s for object B.

[2]

|

**(iii)**

Explain how the principle of conservation of momentum is used to complete the graph in **(a)(ii)**.

.....

.....

.....

[1]

|

**(b)**

Explain, with appropriate working, whether the collision between objects A and B is elastic.

.....

.....

[3]

|

(c)

(i)

Show that the magnitude of the force acting on object A during the collision is  $7.2u$ .

[1]

|

**(ii)**

Explain how the graphs in Fig. 1.1 are consistent with Newton's third law of motion during the collision.

.....

.....

.....

.....

.....

.....



|

[Total: 10]