

- 4 (a) State what is meant by a *longitudinal* wave.

.....
..... [1]

- (b) Fig. 4.1 shows the equilibrium positions of 14 equally spaced air molecules, labelled 1 to 14, along a line AB. The separation between two adjacent equilibrium positions is 0.020 m.

When a point source of sound located on the left of A is switched on, a sinusoidal sound wave travels through the air at a speed of 343 m s^{-1} .

At time $t = 0 \text{ s}$, air molecule 4 is at its maximum displacement towards the right from its equilibrium position, while air molecule 8 is the closest molecule that is at its maximum displacement towards the left.




Fig. 4.1

- (i) Determine, for the sound wave,

1. its wavelength

wavelength = m [1]

2. its frequency

frequency = Hz [1]

- (ii) Label, on Fig. 4.1, with the letter C a point of compression, and the letter R a point of rarefaction.

[1]

- (iii) On the axes in Fig. 4.2, sketch the variation with time t of the displacement y of air molecule 9 for one period of the wave.



Fig. 4.2

[1]

- (iv) The power of the source is P . The intensity of sound recorded by a small detector placed 0.25 m away from the source along AB is I . The detector is then shifted further away along AB until it records an intensity of $\frac{1}{2}I$.

Determine the distance moved by the detector.

distance = m [2]

