

3 (a) The internal energy of an ideal gas is dependent on its state, and is given by the sum of the *random* kinetic energies of all its molecules.

(i) Explain why it is important to include the word *random* in this definition.

.....
.....
..... [1]

(ii) Explain why the potential energy of the molecules is not included in this definition.

.....
.....
..... [1]

(iii) The pressure p exerted by an ideal gas is given by the equation

$$p = \frac{1}{3} \rho \langle c^2 \rangle$$

where ρ is the density of the gas.

Use this equation to derive an expression for the total internal energy U of n moles of an ideal gas at temperature T .

[2]

(iv) State two physical conditions under which a real gas will behave approximately as an ideal gas.

1.
.....

2.

.....

[2]

- (b) A heat engine uses 10 moles of an ideal gas as a working substance. Fig. 3.1 shows the changes in pressure and volume of the gas during one cycle ABCA of operation of the engine.




Fig. 3.1

- (i) Using values from Fig. 3.1, calculate the temperature of the gas at point A.

temperature = K [2]

- (ii) Show that the process $A \rightarrow B$ does not take place at a constant temperature.
 Show your workings in the spaces provided.

.....

..... [1]

(iii) Use Fig. 3.1 to estimate the net work done by the gas during one cycle.

work done = J [2]

(iv) Hence, or otherwise, calculate the amount of heat absorbed by the gas during one cycle.

heat absorbed = J [1]

[Total: 12]