

- 7 In the first half of the last century, numerous experiments were conducted to investigate the absorption and scattering of X-rays by matter.

It was discovered that when a monochromatic beam of X-rays is incident on a light element such as carbon, the scattered X-rays have wavelengths dependent on the angle of scattering.

Compton (1923) assumed that the scattering process could be treated as an elastic collision between an X-ray photon and a 'free' electron, and that energy and momentum would be conserved.

- (a) Explain what is meant by a *photon*.

.....

 [2]

- (b) The elastic collision between a photon and a stationary electron may be represented as in Fig. 7.1.




Fig. 7.1

The incident photon has momentum p and energy E . The photon is scattered through an angle θ and, after scattering has momentum p_s and energy E_s . The electron of mass m , which was originally stationary, moves off with speed v at an angle ϕ to the original direction of the incident photon.

- (i) Write down equations, in terms of p_i , p_s , E_i , E_s , m , v , θ and ϕ , that represent, for this interaction,

1. conservation of energy,

..... [1]

2. conservation of momentum along the direction of the incident photon.

..... [1]

- (ii) Suggest, with a reason, whether the scattered photon will have a wavelength that is greater or less than that of the incident photon.

.....

.....

.....

..... [3]

- (c) In an experiment to provide evidence to justify Compton's theory, measurements were made of the wavelength λ_i of the incident photon, the wavelength λ_s of the scattered photon and the angle θ of scattering. Some data from this experiment are given in Fig. 7.2.

$\lambda_i / 10^{-12} \text{ m}$	$\lambda_s / 10^{-12} \text{ m}$	$\theta / ^\circ$
191.92	193.27	57
153.30	154.65	57
965.04	966.84	75

Fig. 7.2

Use the data in Fig. 7.2 to show that, when a photon is scattered, the change in wavelength produced is independent of the wavelength of the incident photon.

.....

.....

..... [2]

- (d) In this experiment, the uncertainty in the measurement of θ is $\pm 5^\circ$. Determine the value of $\cos \theta$, with its uncertainty, for the angle $\theta = 75^\circ \pm 5^\circ$.

$$\cos \theta = \dots\dots\dots \pm \dots\dots\dots [3]$$

- (e) Compton's theory suggests that the change in wavelength $\Delta\lambda$ is related to the angle θ of the scattering by this expression

$$\Delta\lambda = k(1 - \cos \theta)$$

where k is a constant.

Experimental data for the variation with $\cos \theta$ of $\Delta\lambda$ are shown in Fig. 7.3.




Fig. 7.3

- (i) On Fig. 7.3, draw the best-fit line for the points.

[1]

- (ii) State and explain two different ways by which the constant k may be determined from the graph of Fig. 7.3.

.....

.....

.....

..... [3]

- (iii) Determine the constant k , with its unit.

$k =$ [2]

- (f) For a carbon atom, the binding energy of an electron is of the order of a few electronvolts.

Compton's theory assumes that the electrons are not bound in the atoms but are free.

Suggest whether, for 30 keV photons, this assumption is justified.

.....

.....

.....

..... [2]

[Total: 20]

BLANK PAGE