


1 A  
b  
all  
is  
th  
ro  
w  
n  
fr  
o  
m  
a  
p  
oi  
nt  
P,  
w  
hi  
ch  
is  
at  
gr  
o  
u  
n  
d  
le  
ve  
l,  
as  
ill  
us  
tr  
at  
e  
d  
in  
Fi  
g.  
1.  
1.



**Fig. 1.1**

The initial velocity of the ball is  $12.4 \text{ m s}^{-1}$  at an angle of  $36^\circ$  to the horizontal.

The ball just passes over a wall of height  $h$  at point Q. The ball reaches the wall  $0.17 \text{ s}$  after it has been thrown.

**(a)** Assuming air resistance to be negligible,

**(i)** show that the height of the wall is  $1.1 \text{ m}$ .

[1]

**(ii)** determine the velocity of the ball at point Q.

[3]

magnitude of the velocity = \_\_\_\_\_  $\text{m s}^{-1}$

direction of the velocity = \_\_\_\_\_.

- (b)** A second ball is thrown from point P with the same velocity as the ball in **(a)**. For this ball, air resistance is not negligible. This ball hits the wall elastically at an angle and rebounds.

On Fig. 1.1, sketch a possible path of this ball between point P and the point when it first hits the ground. [2]

[Total : 6]