

- 3 (a) (i) The kinetic theory for an ideal gas of volume V at pressure p leads to the equation

$$pV = \frac{1}{3}Nm\langle c^2 \rangle,$$

where the other symbols refer to their usual meanings.

Use the equation of state for an ideal gas to show that the average translational kinetic energy E_K of a molecule of ideal gas is given by

$$E_K = \frac{3}{2}kT.$$

[1]

- (ii) One helium atom has a mass of 6.68×10^{-27} kg.
Helium may be considered as an ideal gas.

Show that the total kinetic energy of the helium atoms in 1.00 mol of helium gas at 25 °C is 3720 J.

[1]

- (iii) State the value of the internal energy of 1.00 mol of helium gas at 25 °C. Explain your answer.

.....

.....

.....

.....[2]

- (iv) The helium gas is gradually cooled from 25 °C to –150 °C at which the internal energy is 1540 J.

On Fig. 3.1, plot points and draw a line to show the variation with temperature θ of the internal energy U of the helium gas.




Fig. 3.1

[1]

- (v) Explain how your graph leads to the idea of an absolute zero of temperature.

.....

.....

.....[1]

- (b) Gases like hydrogen and helium are found mainly in stars. These gases are at a very high pressure.

Use the assumptions of the kinetic theory of gases to suggest why, in practice, the gas found in stars is unlikely to behave as an ideal gas.

.....

.....

.....

.....[2]

[Total: 8]