

- 3 (a) (i) With reference to the first law of thermodynamics, explain why there is considerable difference in magnitude between the specific latent heats of fusion and vaporisation for the same material.

.....

.....

.....

.....


..... [4]

- (ii) Ethanol has a melting point of  $-120\text{ }^{\circ}\text{C}$  and a boiling point of  $78\text{ }^{\circ}\text{C}$ . The specific latent heat of fusion is  $110\text{ J g}^{-1}$  and specific latent heat of vaporisation is  $840\text{ J g}^{-1}$ . The density and specific heat capacity of liquid ethanol are  $0.79\text{ g cm}^{-3}$  and  $2.4\text{ J g}^{-1}\text{ K}^{-1}$  respectively.

Calculate the minimum thermal energy required to fully vapourise  $2.0\text{ cm}^3$  of ethanol that is initially at  $30\text{ }^{\circ}\text{C}$ .

required thermal energy = ..... J [4]

- (b) Some gas, assumed to behave ideally, is contained within a cylinder which is surrounded by insulation to prevent loss of heat, as shown in Fig. 3.1.



**Fig. 3.1**

Initially, the volume of gas is  $2.9 \times 10^{-5} \text{ m}^3$ , its pressure is  $2.6 \times 10^6 \text{ Pa}$  and its temperature is  $790 \text{ K}$ .

- (i) The gas expands to a volume of  $2.9 \times 10^{-4} \text{ m}^3$  and its temperature decreases to  $314 \text{ K}$ . Calculate the pressure of the gas after this expansion.

pressure of gas after expansion = ..... Pa [2]

- (ii) The work done by the gas during the expansion is  $91 \text{ J}$ . Determine the change in the internal energy of the gas during the expansion.

change in internal energy = ..... J [1]

- (iii) Explain the meaning of internal energy, and use your result in **(b)(ii)** to explain why a decrease in the temperature of the gas takes place during the expansion.

.....

.....

.....

..... [2]

[Total: 13]