

- 9** A monatomic ideal gas in an expandable chamber is cooled from  $50.0\text{ }^{\circ}\text{C}$  to  $10.0\text{ }^{\circ}\text{C}$  at constant pressure  $3.00 \times 10^5\text{ Pa}$ . Total heat given out by the gas is  $1.79 \times 10^4\text{ J}$ .

- (a)** The specific heat capacity at constant pressure of the gas is  $5190\text{ J kg}^{-1}\text{ K}^{-1}$ . One mole of the gas has mass of  $4.0\text{ g}$ .

Determine the number of gas atoms in the chamber.

number of gas atoms = ..... [3]

- (b) (i)** Define an ideal gas. Explain all symbols you may have used in your answer.

.....

.....

..... [2]

- (ii)** Calculate the change in internal energy of the gas when it has been cooled. Explain your answer.

change in internal energy = ..... J [2]

- (iii) Calculate the percentage change in root-mean-square speed of the gas when it has been cooled.

percentage change = .....% [2]

- (c) (i) State the *first law of thermodynamics*.

.....

.....

..... [2]

- (ii) Use the first law of thermodynamics, or otherwise, to determine the work done by the gas it has been cooled.

work done = ..... J [2]

- (iii) Determine the heat given out by the gas if it is cooled from 50.0 °C to 10.0 °C at constant volume instead. Explain your working.

heat given out = ..... J [2]

- (d) (i) Calculate the volume of the gas before and after the change in temperature, considering constant pressure.

volume before = ..... m<sup>3</sup>

volume after = ..... m<sup>3</sup> [1]

- (ii) Hence, sketch in Fig. 9.1 the pressure – volume graph of the gas as it is being cooled from 50.0 °C to 10.0 °C at constant pressure  $3.00 \times 10^5$  Pa.




Fig. 9.1

[2]

- (e) With reference to the heat given out, comment on the difference in your answer in (c)(iii), if any, with the process when the gas is cooled at constant pressure.

.....  
.....  
.....  
.....  
.....

[2]

