

- 9 (a) Define magnetic flux linkage.

.....
.....
..... [2]

- (b) A solenoid of length 30.0 cm is wound evenly with 900 turns of insulated wire. The cross-sectional area of the solenoid is $3.2 \times 10^{-4} \text{ m}^2$.

A flat coil having 250 turns of wire is wound tightly around the centre of the solenoid. A resistor of 35Ω is connected to the coil as illustrated in Fig. 9.1.




Fig. 9.1

- (i) At a particular instant in time, the current through the solenoid is 2.0 A. Show that the magnetic flux through the coil is $2.41 \times 10^{-6} \text{ Wb}$ at that instant.

(ii) The current in the solenoid increases at a uniform rate from 0.0 A to 2.0 A in 0.45 s.

1. Calculate the magnitude of the e.m.f. induced in the flat coil during this period.

e.m.f. = V [2]

2. Hence, calculate the amount of charge that flows through the resistor during this period.

charge = C [2]

3. Determine the amount of charge that passes through the solenoid during this period.

charge = C [2]

- (c) A potential difference is applied between two horizontal plates, each 15.0 cm long and separated by 4.0 cm.

A uniform magnetic field is directed perpendicularly into the page between the two plates.

A beam of electrons moves in a straight line between the plates along a horizontal path halfway between the two plates as shown in Fig. 9.2.

Each electron has a velocity of $4.50 \times 10^6 \text{ m s}^{-1}$.




Fig. 9.2 (not drawn to scale)

- (i) Determine the magnetic flux density of the magnetic field.

magnetic flux density = T [2]

The undeflected beam of electrons enters a second region of uniform magnetic field as shown by the shaded region in Fig. 9.3. A magnetic flux density of 0.50×10^{-3} T is directed perpendicularly into the page.




Fig. 9.3

- (ii) Explain why the electrons move in a uniform circular motion inside the second region of magnetic field.

.....

.....

.....

..... [2]

- (iii) Calculate the radius of the path.

radius = m [2]

- (iv) The beam of electrons is now replaced by a beam of protons. The velocities of the protons before going between the two parallel plates ranges from of $3.00 \times 10^6 \text{ m s}^{-1}$ to $6.00 \times 10^6 \text{ m s}^{-1}$.

Determine the speed of the protons that enters the second region of magnetic field.

speed = m s^{-1} [1]

- (v) Sketch and label the respective paths of the electrons and protons inside the second magnetic field in Fig. 9.3.

[3]

[Total: 20]

End of Paper