

Section B

Answer **one** question from this Section in the spaces provided.

- 8 Fig. 8.1 shows a thin iron strip of length 8.0 cm, width 2.0 cm and thickness 2.0 mm. As iron is a conductor of electricity, it contains free electrons, one of which is shown in Fig. 8.1.




Fig. 8.1

- (a) A small potential difference is applied to the iron strip such that face ABCD is at a potential 12 mV higher than face PQRS.
- (i) Draw, on Fig. 8.1, an arrow to show the direction of the force on the free electron. Label as F_1 . [1]
- (ii) Calculate the magnitude of the force in (a)(i).

force = N [2]

- (iii) Determine the change in the electric potential energy of the electron when it moves across the entire width of the strip.

change in electric potential energy = J [2]

(b) Iron has a resistivity of $9.7 \times 10^{-8} \Omega \text{ m}$ and an electron density of $8.8 \times 10^{28} \text{ m}^{-3}$.

- (i)** Calculate the current in the iron strip when there is a potential difference of 12 mV across faces ABCD and PQRS.

current = A [3]

- (ii)** Hence, show that the drift velocity of the electrons is $4.4 \times 10^{-4} \text{ m s}^{-1}$.

[1]

(c) A magnetic field of flux density 20 mT is now applied in the downward direction into face ABQP.

- (i)** Explain what is meant by a magnitude of 20 mT.

.....

 [1]

- (ii)** Draw, on Fig. 8.1, an arrow to show the direction of the force exerted by the magnetic field on the electron. Label as F_B . [1]

- (iii)** Calculate the magnitude of the force in **(c)(ii)**.

force = N [2]

(iii) Explain why a potential difference develops across faces APSD and BQRC.

.....

 [1]

(iv) Draw, on Fig. 8.1, an arrow to show the direction of the force due to the potential difference/electric field in **(c)(iii)** on the electron. Label as F_E . [1]

(v) The potential difference in **(c)(iii)** eventually reaches a steady value. State the magnitude of the resultant force on the electron.

resultant force = N [1]

(vi) Hence calculate the value of the potential difference in **(c)(iii)**.

potential difference = V [2]

(vii) This iron strip can be used in a device called the Hall probe to determine the magnitude and direction of flux density in a region of space.

Briefly explain how this iron strip can be used to determine the direction of another magnetic field.

.....

 [2]

[Total: 20]