

- 9 The radioactive isotope americium-241 ($^{241}_{95}\text{Am}$) emits an α -particle as it decays to neptunium-237 ($^{237}_{93}\text{Np}$).

(a) Data for the decay are given in Table 9.1.

Table 9.1

nucleus	mass of nucleus / 10^{-27} kg
americium-241	400.19774
neptunium-237	393.54304
helium-4	6.64466

- (i) Show that the energy released in the decay of an americium-241 nucleus is 9.0×10^{-13} J.

[2]

- (ii) State the principle of conservation of linear momentum.

.....

 [2]



- (iii) The kinetic energy of an α -particle emitted by the decay of a stationary americium-241 nucleus is $8.784 \times 10^{-13} \text{ J}$.

Show that the kinetic energy of the neptunium-237 nucleus after the decay is $1.483 \times 10^{-14} \text{ J}$.

[4]

- (iv) Suggest why the sum of the kinetic energies of the α -particle and the neptunium-237 nucleus after the decay is not equal to the energy released in the decay given in (a)(i).

.....
.....
..... [1]



- (b) Americium-241 has a decay constant of $5.08 \times 10^{-11} \text{ s}^{-1}$.

A sample of pure americium-241 has an activity of 8.13 kBq at time $t = 0$.

- (i) Calculate the number of nuclei in the sample at $t = 0$.

number of nuclei = [2]

- (ii) Calculate the half-life of americium-241.

half-life = s [1]

- (iii) On Fig. 9.1, draw a graph to show the variation of the number of americium-241 nuclei with t from $t = 0$ to $t = 6.0 \times 10^{10} \text{ s}$. Label this line **A**.




Fig. 9.1

[2]

- (iv) Draw a second line on Fig. 9.1 to show how the number of neptunium-237 nuclei present in the sample varies from $t = 0$ to $t = 6.0 \times 10^{10} \text{ s}$. Neptunium-237 does not decay significantly over this time. Label this line **N**. [2]



- (c) A sample of a pure beta emitter has the same initial number of nuclei as the sample of americium-241 in (b). The pure beta emitter decays to a stable isotope. The probability of decay per unit time of a nucleus of the beta emitter is much greater than that of a nucleus of americium-241.

Describe and explain, without calculation, any difference in:

- (i) the initial activity of the sample of the beta emitter compared with the initial activity of the sample of americium-241

.....
.....
..... [2]

- (ii) the time taken for the activity of the sample of the beta emitter to fall to 1 Bq compared with the time taken for the activity of the sample of americium-241 to fall to 1 Bq.

.....
.....
..... [2]

[Total: 20]



DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

