

# Solutions to 35th Singapore Physics Olympiad Theory Paper 1

## Question 1: Projectile Motion

### (a) Smooth Inclined Plane

#### 1. Analyze the Projectile Motion

Let  $v$  be the speed of the particle as it leaves the top of the inclined plane at an angle  $\theta = 30^\circ$ . The vertical displacement is  $y = -h = -50 \sin(30^\circ) = -25$  m. The horizontal displacement is  $x = 153.3$  m.

The equations of projectile motion are:

$$\begin{aligned}x &= (v \cos \theta)t \\y &= (v \sin \theta)t - \frac{1}{2}gt^2\end{aligned}$$

Substitute known values:

1.  $153.3 = (v \cos 30^\circ)t \implies t = \frac{153.3}{v \cos 30^\circ}$
2.  $-25 = (v \sin 30^\circ)t - \frac{1}{2}(9.81)t^2$

Substitute  $t$  from (1) into (2):

$$\begin{aligned}-25 &= (v \sin 30^\circ) \left( \frac{153.3}{v \cos 30^\circ} \right) - 4.905 \left( \frac{153.3}{v \cos 30^\circ} \right)^2 \\-25 &= 153.3 \tan 30^\circ - \frac{4.905 \times (153.3)^2}{v^2 (\cos 30^\circ)^2} \\-25 &\approx 88.51 - \frac{153258}{v^2} \\\frac{153258}{v^2} &\approx 113.51 \implies v^2 \approx 1350.2 \implies v \approx 36.74 \text{ m/s}\end{aligned}$$

#### 2. Analyze Motion on the Inclined Plane

Using the kinematic equation  $v^2 = u^2 + 2aL$ :

- Final speed,  $v = 36.74$  m/s
- Distance,  $L = 50$  m
- Acceleration,  $a = -g \sin 30^\circ = -9.81 \times 0.5 = -4.905$  m/s<sup>2</sup>

$$\begin{aligned}(36.74)^2 &= u^2 + 2(-4.905)(50) \\1350.2 &= u^2 - 490.5 \implies u^2 = 1840.7 \implies u \approx 42.9 \text{ m/s}\end{aligned}$$

The initial speed is  $u \approx 43$  m/s.

## (b) Rough Inclined Plane

The speed at the top of the plane must still be  $v = 36.74$  m/s. With a coefficient of kinetic friction  $\mu_k = 0.25$ , the new acceleration  $a'$  is:

$$a' = -(g \sin 30^\circ + \mu_k g \cos 30^\circ)$$

$$a' = -9.81(\sin 30^\circ + 0.25 \cos 30^\circ) \approx -7.029 \text{ m/s}^2$$

Using  $v^2 = (u')^2 + 2a'L$ :

$$(36.74)^2 = (u')^2 + 2(-7.029)(50)$$

$$1350.2 = (u')^2 - 702.9 \implies (u')^2 = 2053.1 \implies u' \approx 45.31 \text{ m/s}$$

The percentage change in  $u$  is:

$$\% \text{ Change} = \frac{u' - u}{u} \times 100\% = \frac{45.31 - 42.9}{42.9} \times 100\% \approx 5.62\%$$

## Question 2: Waves and Oscillations

### (a) Tidal Motion

We model the sea depth  $d(t)$  as a sinusoidal function.

- Mean Depth:  $d_{\text{mean}} = \frac{30+20}{2} = 25$  m.
- Amplitude:  $A = \frac{30-20}{2} = 5$  m.
- Period:  $T = 12$  hours.
- Angular Frequency:  $\omega = \frac{2\pi}{T} = \frac{2\pi}{12} = \frac{\pi}{6}$  rad/hr.

Since depth is minimum at  $t = 0$  (12 noon), the model is  $d(t) = d_{\text{mean}} - A \cos(\omega t)$ .

$$d(t) = 25 - 5 \cos\left(\frac{\pi}{6}t\right)$$

We need to find the times when the depth is  $d(t) = 22.5$  m.

$$22.5 = 25 - 5 \cos\left(\frac{\pi}{6}t\right) \implies \cos\left(\frac{\pi}{6}t\right) = 0.5$$

The principal value is  $\frac{\pi}{6}t = \frac{\pi}{3}$ , which gives  $t = 2$  hours (2:00 pm). By symmetry, the second time is  $t_2 = 12 - 2 = 10$  hours (10:00 pm). The ship can stay in the harbor from **2:00 pm to 10:00 pm**.

**Assumption:** The tidal motion is sinusoidal (Simple Harmonic Motion).

### (b) Doppler Effect and Beats

#### (i) Stationary Source, Moving Object

The frequency detected by the source after reflection,  $f_r$ , from an object moving away at speed  $v_{\text{obj}}$  is:

$$f_r = f_s \left( \frac{v_{\text{sound}} - v_{\text{obj}}}{v_{\text{sound}} + v_{\text{obj}}} \right)$$

Given  $f_s = 256$  Hz,  $v_{\text{sound}} = 340$  m/s, and  $f_{\text{beat}} = 7$  Hz. The reflected frequency is  $f_r = f_s - f_{\text{beat}} = 256 - 7 = 249$  Hz.

$$249 = 256 \left( \frac{340 - v_{\text{obj}}}{340 + v_{\text{obj}}} \right)$$

Solving for  $v_{\text{obj}}$  gives:  $v_{\text{obj}} \approx 4.71$  m/s.

### (ii) Moving Source, Moving Object

The source moves towards the object with  $v_s = 5.0$  m/s. The reflected frequency  $f'_r$  is:

$$f'_r = f_s \left( \frac{v_{\text{sound}} - v_{\text{obj}}}{v_{\text{sound}} - v_s} \right) \left( \frac{v_{\text{sound}} + v_s}{v_{\text{sound}} + v_{\text{obj}}} \right)$$

Using  $v_{\text{obj}} = 4.71$  m/s from our calculation:

$$f'_r = 256 \left( \frac{340 - 4.71}{340 - 5} \right) \left( \frac{340 + 5}{340 + 4.71} \right) \approx 256.44 \text{ Hz}$$

The new beat frequency is  $f'_{\text{beat}} = |f'_r - f_s| = |256.44 - 256| = 0.44$  Hz.

## Question 3: Electrostatics

### (a) Time to Reach the Center

The total charge on the ring is  $Q = \lambda(2\pi R) = (10 \times 10^{-9}) \times (2\pi \times 0.5) = 10\pi \times 10^{-9}$  C. For small displacements,  $x_0 \ll R$ , the force is approximately  $F(x) \approx -kx$ , indicating Simple Harmonic Motion.

$$k = \frac{|q|Q}{4\pi\epsilon_0 R^3} = \frac{(5 \times 10^{-9})(10\pi \times 10^{-9})}{4\pi(8.85 \times 10^{-12})(0.5)^3} \approx 1.129 \times 10^{-5} \text{ N/m}$$

With mass  $m = 1 \times 10^{-6}$  kg, the angular frequency is:

$$\omega = \sqrt{\frac{k}{m}} = \sqrt{\frac{1.129 \times 10^{-5}}{1 \times 10^{-6}}} \approx 3.36 \text{ rad/s}$$

The period is  $T = \frac{2\pi}{\omega} \approx 1.87$  s. The time to reach the center is one-quarter of the period:

$$t = \frac{T}{4} \approx 0.467 \text{ s}$$

### (b) Kinetic Energy at the Center

By conservation of energy,  $K_f = U_i - U_f = qV(x_0) - qV(0)$ .

$$K_f = \frac{qQ}{4\pi\epsilon_0} \left( \frac{1}{\sqrt{R^2 + x_0^2}} - \frac{1}{R} \right)$$

$$K_f = (9 \times 10^9)(-5 \times 10^{-9})(10\pi \times 10^{-9}) \left( \frac{1}{\sqrt{0.5^2 + 0.005^2}} - \frac{1}{0.5} \right)$$

$$K_f \approx 1.414 \times 10^{-10} \text{ J}$$

## Question 4: Thermodynamics

### (a) Thermal Conduction and Entropy

#### (i) Thermal Conductivity

The rate of heat flow  $P$  is:

$$P = \frac{dm}{dt} L_f = \left( \frac{0.1683}{60} \right) (3.36 \times 10^4) = 94.248 \text{ W}$$

From the heat conduction equation,  $P = kA \frac{\Delta T}{L}$ :

$$k = \frac{PL}{A\Delta T} = \frac{(94.248)(0.2)}{(\pi(0.01)^2)(150)} \approx 400 \text{ W m}^{-1}\text{K}^{-1}$$

### (ii) Rate of Entropy Change

$$\frac{dS_{\text{total}}}{dt} = \frac{-P}{T_H} + \frac{P}{T_C} = 94.248 \left( \frac{1}{273.15} - \frac{1}{423.15} \right) \approx 0.1223 \text{ J K}^{-1}\text{s}^{-1}$$

### (b) Radiation Equilibrium

At equilibrium, power absorbed equals power emitted:  $I(\pi R^2) + \sigma(4\pi R^2)T_{\text{env}}^4 = \sigma(4\pi R^2)T_{\text{eq}}^4$ .

$$T_{\text{eq}}^4 = \frac{I}{4\sigma} + T_{\text{env}}^4$$

$$T_{\text{eq}} = \left( \frac{2400}{4(5.67 \times 10^{-8})} + (300.15)^4 \right)^{1/4} \approx 369.7 \text{ K}$$

## Question 5: Photoelectric Effect and Atomic Physics

### (a) Linear Momentum of Photoelectrons

Maximum kinetic energy is  $K_{\text{max}} = eV_s = (1.60 \times 10^{-19})(4.25) = 6.80 \times 10^{-19} \text{ J}$ . The momentum is  $p = \sqrt{2m_e K_{\text{max}}}$ .

$$p = \sqrt{2(9.11 \times 10^{-31})(6.80 \times 10^{-19})} \approx 1.114 \times 10^{-24} \text{ Ns}$$

### (b) Ratio of Emitted Electrons to Incident Photons ( $N_e/N_p$ )

Rate of photoelectron emission:

$$N_e = \frac{I_{\text{sat}}}{e} = \frac{21.7 \times 10^{-9}}{1.60 \times 10^{-19}} = 1.356 \times 10^{11} \text{ electrons/s}$$

Rate of photon incidence:

$$N_p = \frac{P_{\text{in}}}{E_p} = \frac{I \times A}{hc/\lambda} = \frac{(0.7)(10 \times 10^{-4})}{(6.63 \times 10^{-34})(3.00 \times 10^8)/(60 \times 10^{-9})} = 2.112 \times 10^{14} \text{ photons/s}$$

The ratio is:

$$\frac{N_e}{N_p} = \frac{1.356 \times 10^{11}}{2.112 \times 10^{14}} \approx 6.42 \times 10^{-4}$$

### (c) Collision with Hydrogen Atom

The electron's kinetic energy is  $K_{\text{max}} = 4.25 \text{ eV}$ . The minimum energy to excite a hydrogen atom from its ground state ( $n = 1$ ) to the first excited state ( $n = 2$ ) is:

$$\Delta E = E_2 - E_1 = \left( -\frac{13.6}{2^2} \right) - \left( -\frac{13.6}{1^2} \right) = 10.2 \text{ eV}$$

Since the electron's energy (4.25 eV) is less than the required excitation energy (10.2 eV), the electron cannot excite the hydrogen atom. Therefore, no photons will be emitted.