

Problem O. Migratory Birds

OS Linux

Given an array of bird sightings where every element represents a bird type id, determine the id of the most frequently sighted type. If more than 1 type has been spotted that maximum amount, return the smallest of their ids.

Example

$arr = [1, 1, 2, 2, 3]$

There are two each of types **1** and **2**, and one sighting of type **3**. Pick the lower of the two types seen twice: type **1**.

Function Description

Complete the *migratoryBirds* function in the editor below.

migratoryBirds has the following parameter(s):

- $int\ arr[n]$: the types of birds sighted

Returns

- int : the lowest type id of the most frequently sighted birds

Input Format

The first line contains an integer, n , the size of arr .

The second line describes arr as n space-separated integers, each a type number of the bird sighted.

Constraints

- $5 \leq n \leq 2 \times 10^5$
- It is guaranteed that each type is **1**, **2**, **3**, **4**, or **5**.

Sample Input 0

```
6
1 4 4 4 5 3
```

Sample Output 0

```
4
```

Explanation 0

The different types of birds occur in the following frequencies:

- Type **1**: 1 bird
- Type **2**: 0 birds
- Type **3**: 1 bird
- Type **4**: 3 birds
- Type **5**: 1 bird

The type number that occurs at the highest frequency is type **4**, so we print **4** as our answer.

Sample Input 1

```
11
1 2 3 4 5 4 3 2 1 3 4
```

Sample Output 1

```
3
```

Explanation 1

The different types of birds occur in the following frequencies:

- Type **1**: 2
- Type **2**: 2
- Type **3**: 3
- Type **4**: 3
- Type **5**: 1

Two types have a frequency of **3**, and the lower of those is type **3**.