



Swiss Institute of  
Bioinformatics


SINGLE-CELL TRANSCRIPTOMICS WITH R

# Differential gene expression and enrichment analysis

Deepak Tanwar

March 18-20, 2026

Adapted from previous year courses



# Learning objectives

## Differential gene expression analysis

- Finding markers
- Pseudo-bulk analysis

## Enrichment analysis

- What is Enrichment analysis?
- Distinguish between different ways to do it
- Challenges and Limitations of methods

# Differential gene expression analysis

DGE can be divided into 2 sub-groups:

## a. Single-Cell-Level DGE Analysis (finding markers)

**Goal:** Identify genes that are differentially expressed between cell populations (e.g., cell types, clusters, or conditions) while accounting for the single-cell nature of the data (e.g., sparsity, dropout events).

# Differential gene expression analysis

DGE can be divided into 2 sub-groups:

## a. Single-Cell-Level DGE Analysis (finding markers)

**Goal:** Identify genes that are differentially expressed between cell populations (e.g., cell types, clusters, or conditions) while accounting for the single-cell nature of the data (e.g., sparsity, dropout events).

## b. Pseudo-Bulk DGE Analysis (differential analysis)

**Goal:** Aggregate single-cell data into pseudo-bulk profiles to perform DGE analysis using bulk RNA-seq methods, reducing noise and leveraging biological replicates.

# Marker gene identification

Genes overexpressed by each cell type or cluster with the dataset => It can help in cell type annotation

# Marker gene identification

## Methods:

- **Log-Fold Change (LFC) Analysis:** Calculate the log-fold change in expression between a target cluster and all other cells (e.g., in Seurat or Scanpy).
- **Wilcoxon Rank-Sum Test:** Test for genes with significantly higher expression in one group compared to others (e.g., in Seurat).

# Marker gene identification

## Use Cases:

- Annotating cell types (e.g., identifying CD3 as a marker for T cells)
- Discovering novel cell states or subpopulations

# Typical Output

| Gene Symbol | avg_log2FC | pct.1 | pct.2 | p_val     | p_val_adj | cluster |
|-------------|------------|-------|-------|-----------|-----------|---------|
| CHI3L1      | 5.61       | 0.958 | 0.225 | 7.63E-255 | 1.97E-250 | 2       |
| HLA-DRA     | 3.41       | 0.978 | 0.215 | 2.84E-253 | 7.32E-249 | 2       |
| PTGFR       | 4.31       | 0.795 | 0.093 | 6.43E-244 | 1.66E-239 | 2       |
| HLA-DRB5    | 3.54       | 0.818 | 0.097 | 2.10E-243 | 5.41E-239 | 2       |
| GRIN2A      | 4.24       | 0.69  | 0.05  | 1.64E-235 | 4.22E-231 | 2       |
| CDHR3       | 3.34       | 0.892 | 0.159 | 2.56E-229 | 6.59E-225 | 2       |
| AKR1C3      | 3.16       | 0.955 | 0.254 | 5.75E-223 | 1.48E-218 | 2       |
| KCNK15      | 3.76       | 0.897 | 0.187 | 4.88E-217 | 1.26E-212 | 2       |
| HLA-DRB1    | 2.53       | 0.78  | 0.102 | 2.71E-212 | 6.98E-208 | 2       |
| PLPP3       | 3.34       | 0.965 | 0.322 | 3.54E-210 | 9.11E-206 | 2       |



# Marker gene identification

## Challenges:

- Marker genes may not be unique to a single cell type, requiring careful validation
- Dropout events can obscure marker gene detection

Research | [Open access](#) | Published: 26 February 2024

## A comparison of marker gene selection methods for single-cell RNA sequencing data


[Jeffrey M. Pullin](#) & [Davis J. McCarthy](#) 

[Genome Biology](#) **25**, Article number: 56 (2024) | [Cite this article](#)


**16k** Accesses | **32** Altmetric | [Metrics](#)

Methods based on logistic regression, Student's  $t$ -test and the Wilcoxon rank-sum test all have strong performance


# Highly-significant $p$ -value



# Highly-significant $p$ -value




# $P$ -value inflation



The probability of observing a test statistic at least as extreme as the one observed, assuming that the null hypothesis is true.

# Pseudo-Bulk DGE Analysis



# Pseudo-Bulk DGE Analysis

Genes whose expression is modulated (up or down) by experimental condition within a cell type or cluster



# What is the ideal method?

Analysis | Published: 26 February 2018

## **Bias, robustness and scalability in single-cell differential expression analysis**

[Charlotte Soneson](#)  & [Mark D Robinson](#) 

Many methods have been used to determine differential gene expression from single-cell RNA (scRNA)-seq data. We evaluated 36 approaches using experimental and synthetic data and found considerable differences in the number and characteristics of the genes that are called differentially expressed. Prefiltering of lowly expressed genes has important effects, particularly for some of the methods developed for bulk RNA-seq data analysis. However, we found that bulk RNA-seq analysis methods do not generally



# What is the ideal method?




Figure 5 | Summary of DE method performance across all major evaluation criteria. Criteria and cutoff values for performance categories are available in the Online Methods. Methods are ranked by their average performance across the criteria, with the numerical encoding good = 2, intermediate = 1, poor = 0.

# Limma/edgeR: old but gold

Methods designed for microarray and bulk RNAseq analysis

- Can be used to include batch effects in model as covariates
- Compare more than 2 groups: e.g. ANOVA(F-test)

Analysis with limma and example of model with covariate:

<https://ucdavis-bioinformatics-training.github.io/2018-June-RNA-Seq-Workshop/thursday/DE.html>

# Pseudo-Bulk DGE Analysis

## Methods:

- Aggregate gene expression counts within groups (e.g., by cell type and sample) and use bulk RNA-seq tools like DESeq2, edgeR, or limma
- Tools like muscat in Bioconductor are specifically designed for pseudo-bulk DGE analysis in scRNA-seq

# Pseudo-Bulk DGE Analysis

## Advantages:

- Reduces noise and dropout effects
- Leverages well-validated bulk RNA-seq tools

# Pseudo-Bulk DGE Analysis

**Limitations:** Loses single-cell resolution and cannot detect cell-to-cell variability

**Single Cell-Level: Finding markers**

**Pseudo-Bulk: Differential expression**


# Quiz

**What is a key limitation of pseudo-bulk analysis compared to single cell-level analysis?**


- A) It is computationally more intensive.**
- B) It loses single-cell resolution and cannot detect cell-to-cell variability.**
- C) It cannot be used for differential expression analysis.**
- D) It is less robust to dropout events.**

# Pseudo-bulk DE analysis: muscat


## Healthy donor A




## Patient A



## Healthy donor B




## Patient B






**Goal:** To gain biologically meaningful insights from long gene lists

# Overview of functional analysis: ORA



# Overview of functional analysis: FCS


## Functional Pathway Analysis






# Overview of functional analysis: PT

## Functional Pathway Analysis



# Overview of functional analysis: ORA & FCS



# What is a gene set?

A gene set is an unordered collection of genes that are functionally related.

- Genes located in the same compartment in a cell (e.g. all proteins located in the cell nucleus)
- Proteins that are all regulated by a same transcription factor
- Custom gene list that comes from a publication and that are down-regulated in a mutant
- List of genes that contain SNPs associated with a disease
- ...etc!
- Several gene sets are grouped into Knowledge bases
- A pathway can be interpreted as a gene set by ignoring functional relationships among genes

# Over-representation analysis (ORA)

Statistically evaluates the fraction of genes in a particular pathway found among the set of genes showing changes in expression.

# Over-representation analysis (ORA)

Statistically evaluates the fraction of genes in a particular pathway found among the set of genes showing changes in expression.

1. Select a list of genes with certain threshold ( $\text{FDR} \leq 0.05$ )



# Over-representation analysis (ORA)

Statistically evaluates the fraction of genes in a particular pathway found among the set of genes showing changes in expression.

1. Select a list of genes with certain threshold ( $\text{FDR} \leq 0.05$ )
2. For each pathway, count input genes that are part of the pathway

# Over-representation analysis (ORA)

Statistically evaluates the fraction of genes in a particular pathway found among the set of genes showing changes in expression.

1. Select a list of genes with certain threshold ( $\text{FDR} \leq 0.05$ )
2. For each pathway, count input genes that are part of the pathway
3. Repeat for an appropriate background list of genes

# Over-representation analysis (ORA)

Statistically evaluates the fraction of genes in a particular pathway found among the set of genes showing changes in expression.

1. Select a list of genes with certain threshold ( $\text{FDR} \leq 0.05$ )
2. For each pathway, count input genes that are part of the pathway
3. Repeat for an appropriate background list of genes
4. Every pathway is tested for over- or under-representation in the list of input genes

# Over-representation analysis (ORA)

Statistically evaluates the fraction of genes in a particular pathway found among the set of genes showing changes in expression.

1. Select a list of genes with certain threshold ( $\text{FDR} \leq 0.05$ )
2. For each pathway, count input genes that are part of the pathway
3. Repeat for an appropriate background list of genes
4. Every pathway is tested for over- or under-representation in the list of input genes

The most commonly used tests are based on the [hypergeometric](#), [chi-square](#), or [binomial distribution](#)

# Over-representation analysis (ORA)

|         |         |
|---------|---------|
| Gene1   | 0.051   |
| Gene2   | 0.05001 |
| Gene 3  | 0.049   |
| Gene 4  | 0.001   |
| Gene 5  | 0.023   |
| Gene 6  | 0.04    |
| Gene 7  | 0.01    |
| Gene 8  | 0.0501  |
| Gene 9  | 0.2     |
| Gene 10 | 0.051   |
| Gene 11 | 0.05    |
| Gene 12 | 0.49    |
| Gene 13 | 0.03    |
| Gene 14 | 0.01    |
| Gene 15 | 0.052   |
| Gene 16 | 0.9     |

# Over-representation analysis (ORA)

|         |         |
|---------|---------|
| Gene1   | 0.051   |
| Gene2   | 0.05001 |
| Gene 3  | 0.049   |
| Gene 4  | 0.001   |
| Gene 5  | 0.023   |
| Gene 6  | 0.04    |
| Gene 7  | 0.01    |
| Gene 8  | 0.0501  |
| Gene 9  | 0.2     |
| Gene 10 | 0.051   |
| Gene 11 | 0.05    |
| Gene 12 | 0.49    |
| Gene 13 | 0.03    |
| Gene 14 | 0.01    |
| Gene 15 | 0.052   |
| Gene 16 | 0.9     |

*pvalue* ≤ 0.05

|        |       |
|--------|-------|
| Gene 3 | 0.049 |
| Gene 4 | 0.001 |
| Gene 5 | 0.023 |
| Gene 6 | 0.04  |
| Gene 7 | 0.01  |

|         |      |
|---------|------|
| Gene 11 | 0.05 |
| Gene 12 | 0.49 |
| Gene 13 | 0.03 |
| Gene 14 | 0.01 |

# Over-representation analysis (ORA)


|         |         |
|---------|---------|
| Gene1   | 0.051   |
| Gene2   | 0.05001 |
| Gene 3  | 0.049   |
| Gene 4  | 0.001   |
| Gene 5  | 0.023   |
| Gene 6  | 0.04    |
| Gene 7  | 0.01    |
| Gene 8  | 0.0501  |
| Gene 9  | 0.2     |
| Gene 10 | 0.051   |
| Gene 11 | 0.05    |
| Gene 12 | 0.49    |
| Gene 13 | 0.03    |
| Gene 14 | 0.01    |
| Gene 15 | 0.052   |
| Gene 16 | 0.9     |

*pvalue* ≤ 0.05

|        |       |
|--------|-------|
| Gene 3 | 0.049 |
| Gene 4 | 0.001 |
| Gene 5 | 0.023 |
| Gene 6 | 0.04  |
| Gene 7 | 0.01  |

|         |      |
|---------|------|
| Gene 11 | 0.05 |
| Gene 12 | 0.49 |
| Gene 13 | 0.03 |
| Gene 14 | 0.01 |

Fisher's test



$H_0$ : The proportion of genes in the gene set is the same for both groups

$H_a$ : The proportion of genes in the gene set is higher in the differentially expressed group



# Problems with ORA

Cutoff? 0.051?

# Problems with ORA

Cutoff? 0.051?

Treat all genes equally

# Problems with ORA

Cutoff? 0.051?

Treat all genes equally

Each gene is independent of other

# Problems with ORA

Cutoff? 0.051?

Treat all genes equally

Each gene is independent of other

Each pathway is independent of each other

# Gene set enrichment analysis (GSEA)

|         |         |     |
|---------|---------|-----|
| Gene1   | 0.051   | 10  |
| Gene2   | 0.05001 | 12  |
| Gene 3  | 0.049   | 11  |
| Gene 4  | 0.001   | 8   |
| Gene 5  | 0.023   | 2   |
| Gene 6  | 0.04    | 3   |
| Gene 7  | 0.01    | 1   |
| Gene 8  | 0.0501  | 3   |
| Gene 9  | 0.2     | -10 |
| Gene 10 | 0.051   | -3  |
| Gene 11 | 0.05    | -8  |
| Gene 12 | 0.49    | -19 |
| Gene 13 | 0.03    | -3  |
| Gene 14 | 0.01    | -2  |
| Gene 15 | 0.052   | -1  |
| Gene 16 | 0.9     | -4  |

# Gene set enrichment analysis (GSEA)

|         |         |     |
|---------|---------|-----|
| Gene1   | 0.051   | 10  |
| Gene2   | 0.05001 | 12  |
| Gene 3  | 0.049   | 11  |
| Gene 4  | 0.001   | 8   |
| Gene 5  | 0.023   | 2   |
| Gene 6  | 0.04    | 3   |
| Gene 7  | 0.01    | 1   |
| Gene 8  | 0.0501  | 3   |
| Gene 9  | 0.2     | -10 |
| Gene 10 | 0.051   | -3  |
| Gene 11 | 0.05    | -8  |
| Gene 12 | 0.49    | -19 |
| Gene 13 | 0.03    | -3  |
| Gene 14 | 0.01    | -2  |
| Gene 15 | 0.052   | -1  |
| Gene 16 | 0.9     | -4  |

Research Article | [Open access](#) | Published: 12 May 2017

# Ranking metrics in gene set enrichment analysis: do they matter?

Joanna Zyla, Michal Marczyk✉, January Weiner & Joanna Polanska

*BMC Bioinformatics* **18**, Article number: 256 (2017) | [Cite this article](#)

## Genes ranked by test statistic

or

$$\log_2(\text{FC}) * t\text{-value}$$


Upregulated

$H_0$ : Genes in set are randomly distributed over ranked list

$H_a$ : Genes in set are not randomly distributed over the ranked list

## Downregulated

# Functional class scoring (FCS)



# Problems with FCS

Each gene is independent of other



# Problems with FCS

Each gene is independent of other

Each pathway is independent of each other

## Databases

- GO: BP, MF, CC
- KEGG
- Reactome
- DOSE
- DisGeNET
- MSigDb
- KEGG module
- WikiPathways
- TF
- miRNA
- "user input"
- PathGuide

## Methods

- ORA
- GSEA
- SAFE
- PADOG
- ROAST
- CAMERA
- GSA
- GSVA/ssGSEA
- GlobalTest
- EBM
- MGSA
- GOSeq
- QUSAGE
- Pathview
- GOSemSim
- GGEA
- SPIA
- PathNet
- DEGraph
- TopologyGSA
- GANPA
- CePa
- NetGSA
- WGCNA

# Databases and methods

## Databases

- GO: BP, MF, CC
- KEGG
- Reactome
- DOSE
- DisGeNET
- MSigDb
- KEGG module
- WikiPathways
- TF
- miRNA
- "user input"
- PathGuide


## Methods

- ORA
- GSEA
- SAFE
- PADOG
- ROAST
- CAMERA
- GSA
- GSVA/ssGSEA
- GlobalTest
- EBM
- MGSA
- GOSeq
- QUSAGE
- Pathview
- GOSemSim
- GGEA
- SPIA
- PathNet
- DEGraph
- TopologyGSA
- GANPA
- CePa
- NetGSA
- WGCNA

Problems with databases:  
Low resolution


Databases and methods

# Gene Ontology: the world's largest source of information on the functions of genes



The GO contains many terms that are highly similar or overlapping in meaning (e.g., "cell cycle" and "mitosis").

# Semantic Similarity Measurement Based on *Exclusively Inherited* Shared Information for Gene Ontology



"exclusively inherited" refers to the subset of shared information that is **unique to the two terms being compared** ( $GO_{Term_5}$  and  $GO_{Term_6}$ ) and **not inherited by other unrelated terms.**




Illustration of Semantic Similarity Measurement for Gene Ontology Terms Using Exclusively Inherited Shared Information

# Making your own database

database\_seeds

\$paper1\_day1

Gene1, Gene2, Gene3, Gene4

\$paper2\_day2


Gene3, Gene4, Gene5, Gene6

# clusterProfiler

## A universal enrichment tool for interpreting omics data

platforms all rank 36 / 2300 support 1 5 / 1 8 in Bioc 13.5 years build ok updated < 3 months dependencies 132

DOI: [10.18129/B9.bioc.clusterProfiler](https://doi.org/10.18129/B9.bioc.clusterProfiler)



Wu T, et al. (2021). "clusterProfiler 4.0: A universal enrichment tool for interpreting omics data."

*The Innovation*, 2(3), 100141. [doi:10.1016/j.xinn.2021.100141](https://doi.org/10.1016/j.xinn.2021.100141).

# Quiz

**1. Single cell-level pathway analysis can provide insights into cell-to-cell variability in pathway activity, while pseudo-bulk analysis cannot.**

- A) True
- B) False

**2. Using "exclusively inherited" shared information in semantic similarity calculations helps reduce the impact of redundant GO terms.**

- A) True
- B) False



# Summary

Three types of methods for enrichment analysis:

1. ORA
2. FCS
3. Pathway Topology

Databases problem

GO semantic similarity

