



Deterministic OR Models

CO 370



Bertrand Guenin

Preface

Disclaimer Much of the information on this set of notes is transcribed directly/indirectly from the lectures of CO 370 during Fall 2021 as well as other related resources. I do not make any warranties about the completeness, reliability and accuracy of this set of notes. Use at your own risk.

What is operations research (OR)? There's no standard definitions for it. One particular definition: use of mathematical models to make complex decisions for real life problems. The origin is British military in WW2. OR is actually everywhere today. Key milestone: Simplex algorithm (1947).

Recall optimization problem is of the form:

$$\begin{array}{ll}\max & f(x) \\ \text{s.t.} & \text{a set of constraints}\end{array}$$

There are some applications: mail delivery, machine scheduling, inventory problem, network design, facility location, class scheduling, portfolio optimization, surgery planning, sensor location.

For any questions, send me an email via <https://notes.sibeliusp.com/contact>.

You can find my notes for other courses on <https://notes.sibeliusp.com/>.

Sibeliusp Peng

Contents

Preface	1
I Formulations	3
1 LP formulations	4
1.1 Production problem	4
1.2 Minimax	5
2 Flows	7
2.1 Max st-flow model	7
2.2 Min cost flow model	8
3 IP formulations	10
3.1 IP tricks	10
3.2 Modeling piece-wise linear function	10
3.3 Modeling union of polyhedra	11
3.4 Perfect formulations	11



PART I:

FORMULATIONS

LP formulations

1.1 Production problem

Products $J = \{1, \dots, n\}$

Resources $I = \{1, \dots, m\}$

Data:

- $\forall j \in J : c_j = \text{value of unit of product } j$
- $\forall i \in I : b_i = \text{number of units of resource } i \text{ available}$
- $\forall i \in I, \forall j \in J : a_{ij} = \text{number of units of resource } i \text{ going to product } j$

Goal: maximize values of product made subject to available resources

Var: $x_j = \text{number of units of product } j \text{ produced}$

Then problem is

$$\begin{array}{ll} \max & \sum_{j \in J} c_j x_j \\ \text{s.t.} & \sum_{j \in J} a_{ij} x_j \leq b_i \quad (i \in I) \\ & x_j \geq 0 \quad (j \in J) \end{array}$$

Now let's generalize this problem to have more than one period.

Products $J = \{1, \dots, n\}$

Resources $I = \{1, \dots, m\}$

Periods $K = \{1, \dots, p\}$

Then we have **data**

- $\forall j \in J, k \in K : c_{jk} = \text{unit value of product } j \text{ in period } k$
- $\forall i \in I, k \in K : w_{ik} = \text{unit price for resource } i \text{ in period } k$
- $\forall i \in I, j \in J : a_{ij} = \text{number of units of resource } i \text{ going into product } j$

and the **goal**: decide how much of each resource to buy & how much of each product to make during each period, to maximize total profit. Unused resources are available at next time period.

Var:

- p_{ik} = number of units of resource i , purchased at start of period k
- x_{jk} = number of units of product j made in period k
- z_{ik} = number of units of resource i at the end of period k

$$\text{Profit} = \sum_{k \in K} \left[\sum_{j \in J} c_{jk} x_{jk} - \sum_{i \in I} w_{ik} p_{ik} \right] \quad (1.1)$$

Then we keep track of resources: for $i \in I, k \in K$

$$z_{ik} = z_{i(k-1)} + p_{ik} - \sum_{j \in J} a_{ij} x_{jk} \quad (1.2)$$

and we define for $i \in I$,

$$z_{i0} = 0 \quad (1.3)$$

Thus the optimization problem is

$$\begin{aligned} \max \quad & (1.1) \\ \text{s.t.} \quad & (1.2), (1.3) \\ & p, x, z \geq 0 \end{aligned} \quad (P)$$

Remark:

If (P) has a feasible solution of value that is bigger than 0, then (P) is unbounded. So we are missing some assumptions, maybe? For example, b_{ik} = amount of resource i that can be bound during period k . Then we can add constraints: $p_{ik} \leq b_{ik}$.

1.2 Minimax


Consider the problem of the form

$$\begin{aligned} \min_x \max \{f_1(x), \dots, f_k(x)\} &:= g(x) \\ \text{s.t.} \quad & \dots \end{aligned}$$

where $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$.

Example:

$f_1(x) = -x - 1, f_2(x) = 0, f_3(x) = x - 1$. Then $\max\{f_1(x), f_2(x), f_3(x)\}$ is as follows



A motivation

- $\forall i \in [k], f_i(x) = \text{completion time for task } i.$
- Project consists of task $1, \dots, k.$
- $g(x) = \text{completion time of entire project}$

Note that minimax is not an optimization problem as we defined it. We can revise it as follows

$$\begin{array}{ll}
 \min & y \\
 \text{s.t.} & y \geq f_1(x) \\
 & y \geq f_2(x) \\
 & \vdots \\
 & y \geq f_k(x) \\
 & \dots
 \end{array}$$

An application minimize a piece-wise linear convex function using linear programming.

Flows

digraph

A directed graph (digraph) is a pair (V, E) where

- V is a set of vertices,
- E is a set of ordered pairs of vertices called arcs.

Notation Let $q \in V$, then

$$\delta^+(q) = \{e \in E \mid e \text{ leaves } q\}$$

$$\delta^-(q) = \{e \in E \mid e \text{ arrives at } q\}$$

2.1 Max st-flow model

Given

1. digraph $G = (V, E)$,
2. two vertices $s, t \in V$, and $s \neq t$,
3. $\forall e \in E$, arc e has capacity $u_e \geq 0$.

Now we construct an LP.

For every arc e , we will have a variable x_e , and x_e will be called the **flow** on arc e .

Notation Let $q \in V$:

$$f_x(q) := \sum_{e \in \delta^+(q)} x_e - \sum_{e \in \delta^-(q)} x_e$$

The maximization problem is then

$$\begin{aligned} \max \quad & f_x(s) \\ \text{s.t.} \quad & f_x(q) = 0 \quad (q \in V, q \neq s, q \neq t) \\ & 0 \leq x_e \leq u_e \quad (e \in E) \end{aligned} \tag{P}$$

A feasible solution to (P) is a flow. An optimal solution to (P) is a maximum flow. The value of a flow x is $f_x(S)$.

Remark:

(P) is always feasible. It is not unbounded, so there always exists a maximum flow.

Application computer network. Suppose we have

- computers s and t ($s \neq t$)
- capacity u_e (gb/s) for every link e

The goal is to computer number gb that can be sent from s to t across network. Then x_e is the amount of information across e . $f_x(q) = 0$ means no information lost.

Magic property

If u is integer, then there exists an optimal solution to (P) that is integer.

Remark:

We need the condition “ u is integer” so that the property is still true. Also, an optimal solution to (P) is not necessarily integers.

Generalize max st-flows

We can add lower bounds to arcs: $\ell_e \forall e \in E$

$$\begin{aligned} \max \quad & f_x(s) \\ \text{s.t.} \quad & f_x(q) = 0 \quad (q \in V, q \neq s, q \neq t) \\ & \ell_e \leq x_e \leq u_e \quad (e \in E) \end{aligned} \tag{Q}$$

Magic property - revised

If ℓ, u is integer, and there exists an optimal solution to (Q), then there exists an optimal solution to (Q) that is integer.

Example: Consistent rounding

The goal is to round all entries to nearest up/down integer, so that row sums & column sums still hold.

Any feasible solutions give consistent rounding.

2.2 Min cost flow model

Given

- Digraph $G = (V, E)$
- Capacities $u_e \geq 0$ ($e \in E$)
- Costs c_e ($e \in E$)
- Supply/demands b_q ($q \in V$)

Then the model is

$$\begin{array}{ll} \min & \sum_{e \in E} c_e x_e \\ \text{s.t.} & f_x(q) = b_q \quad (q \in V) \\ & 0 \leq x_e \leq u_e \quad (e \in E) \end{array} \quad (P)$$

Similarly, feasible solution to (P) is a flow. An optimal solution to (P) is a min cost flow.

Magic property - min cost flows

Suppose u, b are integer, then if there exists a min cost flow, then there exists a min cost flow that is integer.

Similarly, we can add a lower bound:

$$\begin{array}{ll} \min & \sum_{e \in E} c_e x_e \\ \text{s.t.} & f_x(q) = b_q \quad (q \in V) \\ & \ell_e \leq x_e \leq u_e \quad (e \in E) \end{array} \quad (P)$$

Magic property - min cost flows (revised)

Suppose u, b, ℓ are integer, then if there exists a min cost flow, then there exists a min cost flow that is integer.

What is a necessary condition for b_q so that there exists a flow? $\sum_{q \in V} b_q = 0$.

Example: Staffing problem

hours	# of employees required
1-2	2
2-3	3
3-4	3
4-5	2

and we have cost of hiring a single employee between

hours	cost
1-5	6
1-4	4
3-5	5
2-4	3

The goal is to minimize cost of hiring employees while meeting staff needs.

IP formulations

3.1 IP tricks

Imaging we are forcing variable to take some prescribed set of values: $x \in \{5, 9, 13, 36\}$. Then we can introduce variables $z_1, z_2, z_3, z_4 \in \{0, 1\}$ so that we have two constraints:

$$\begin{aligned} 1 &= z_1 + z_2 + z_3 + z_4 \\ x &= 5z_1 + 9z_2 + 13z_3 + 36z_4 \end{aligned}$$

3.2 Modeling piece-wise linear function

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be piecewise linear. Given a_1, \dots, a_k and f_1, \dots, f_k such that $f(a_i) = f_i$. The goal is to write IP constraints with variables x, y such that $y = f(x)$, $x \in [a_1, a_k]$.

To generalize,

\uparrow
dom(f)

1. $\lambda_1, \dots, \lambda_k \geq 0, \sum_{i=1}^k \lambda_i = 1$
2. $x = \sum_{i=1}^k \lambda_i a_i$
3. $y = \sum_{i=1}^k \lambda_i f_i$
4. $z_0, \dots, z_k \in \{0, 1\}, z_0 = z_k = 0$
5. $\sum_{i=0}^k z_i = 1$
6. $\forall p \in [k], \lambda_p \leq z_{p-1} + z_p$

By (4) and (5), we may assume that $z_p = 1$ and $z_j = 0 \forall j \neq p$.

Claim λ_p, λ_{p+1} are the only non-zero λ variables.

Proof:

Pick $j \neq p, j \neq p+1$. As

$$0 \leq \lambda_j \leq z_{j-1} + z_j = 0 + 0 = 0$$

□

With this claim, we can simplify (2) and (3).

3.3 Modeling union of polyhedra

Let

$$P_1 = \{x \mid A^1 x \leq b^1\}$$

$$P_2 = \{x \mid A^2 x \leq b^2\}$$

Goal: write condition: $x \in P_1 \cup P_2$ as part of IP.

Hypothesis: If $x \in P_1 \cup P_2$, then $0 \leq x \leq U$ for some U .

Constraints:

1. $y_1, y_2 \in \{0, 1\}$
2. $y_1 + y_2 = 1$
3. $x = x^1 + x^2$
4. $A^i x^i \leq y_i b^i, i = 1, 2$
5. $0 \leq x^i \leq y_i U, i = 1, 2$

To show that $x \in P_1 \cup P_2 \iff \exists x^1, x^2, y_1, y_2$ such that all 5 conditions hold.

Proof:

First let's assume (1) - (5) hold. We may assume $y_1 = 1, y_2 = 0$. Then (5) tells us $x^2 = 0$. (3) tells us $x = x^1$. (4) implies

$$A^1 x^1 \leq y_1 b^1 \implies A^1 x \leq b^1 \implies x \in P_1 \subseteq P_1 \cup P_2$$

Now suppose $x \in P_1 \cup P_2$. We may assume $x \in P_1$. Then we set $y_1 = 1, y_2 = 0, x^1 = x, x^2 = 0$. Then we can verify that all 5 conditions hold. \square

3.4 Perfect formulations

basis

Let A be a matrix with column indices $\{1, \dots, n\}$. Then $B \subseteq \{1, \dots, n\}$ is a basis if

1. A_B square,
2. A_B non-singular.

Remark:

A has a basis \iff rows of A are independent.

basic solution

Let A be a matrix with column indices $1, \dots, n$. Consider

$$Ax = b \tag{*}$$

Pick B as basis of A . Then x is a basic solution of (*) if

1. $Ax = b$.
2. $x_j = 0$ if $j \notin B$.

x is a basic solution for (*) if x is a basic solution for some basis B .

standard equality form

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned}$$

and rows of A are independent.

The correctness of simplex algorithm implies the following theorem:

Theorem 3.1

If an LP is SEF has an optimal solution, then it has an optimal solution that is basic.

Let A be matrix, b vector with same number of entries as rows of A . Then $A \leftarrow_j b$ denotes matrix obtained from A by replacing column j by b .

Theorem 3.2: Cramer's rule

Let M be a non-singular matrix and consider $Mx = b$.

$$\bar{x}_j = \frac{\det(M \leftarrow_j b)}{\det(M)} \quad \forall j$$

then $M\bar{x} = b$.

Proposition 3.3

Let M be square matrix with $\det(M) = \pm 1$, and M, b are integer. Then there exists a unique solution to $Mx = b$ is integer.

Proof:

Directly from Cramer's rule. □

Totally unimodular matrix

A matrix A is totally unimodular if every square submatrix N of A , $\det(N) \in \{0, +1, -1\}$.