



# *Graph Theory*

CO 442



Luke Postle

# Preface

---

**Disclaimer** Much of the information on this set of notes is transcribed directly/indirectly from the lectures of CO 442 during Fall 2020 as well as other related resources. I do not make any warranties about the completeness, reliability and accuracy of this set of notes. Use at your own risk.

Here is the notation used in this course.

- $\chi(G)$ : chromatic number,  $k$  vertex coloring
- $\Delta(G)$ : max degree of vertices
- $\delta(G)$ : min degree of vertices
- $\omega(G)$ : max size of a clique
- $\chi'(G)$ : chromatic index, edge chromatic number,  $k$  edge coloring
- $L(G)$ : line graph of  $G$
- $\chi_\ell(G)$ : list chromatic number
- $\chi'_\ell(G)$ : list chromatic index
- $\mu(G)$ : multiplicity of a multigraph  $G$

For any questions, send me an email via <https://notes.sibeliusp.com/contact/>.

You can find my notes for other courses on <https://notes.sibeliusp.com/>.

---

*Sibelius Peng*

# Contents

---

<b>Preface</b>	<b>1</b>
<b>1 Colorings</b>	<b>4</b>
1.1 Coloring and Brooks' Theorem . . . . .	4
1.2 Edge Coloring . . . . .	12
1.2.1 Vizing's Theorem . . . . .	14
1.2.2 List Edge Coloring . . . . .	16
1.2.3 Edge Coloring Multigraphs . . . . .	18
1.3 Thomassen's Theorem . . . . .	21
1.4 Coloring and List Coloring Planar Graphs . . . . .	23
1.5 Discharging . . . . .	24
1.5.1 Common Discharging Setups for Planar Graphs . . . . .	25

First let's look at a proof example.

### Theorem

Every two longest paths in a connected graph  $G$  intersect.

#### Proof:

Suppose not. That is, there exist two longest paths  $P_1$  and  $P_2$  of  $G$  such that  $V(P_1) \cap V(P_2) = \emptyset$ . For each  $i \in \{1, 2\}$ , let  $v_{i,1}$  and  $v_{i,2}$  be the ends of  $P_i$ . Since  $G$  is connected, there exists a shortest path  $P$  from  $V(P_1)$  to  $V(P_2)$ . Since  $P$  is shortest, we have that  $|V(P_i) \cap V(P)| = 1$  for each  $i \in \{1, 2\}$ .

For each  $i \in \{1, 2\}$ , let  $u_i$  be the end of  $P$  in  $V(P_i)$ . For each  $i, j \in \{1, 2\}$ , let  $Q_{i,j}$  be the subpath of  $P_i$  from  $u_i$  to  $v_{i,j}$ . We assume without loss of generality that for each  $i \in \{1, 2\}$ , we have that  $|E(Q_{i,1})| \geq |E(Q_{i,2})|$  and hence

$$|E(Q_{i,1})| \geq |E(P_i)|/2.$$

Let  $P' = v_{1,1}Q_{1,1}u_1Pu_2Q_{2,1}v_{2,1}$ . Note that  $P'$  is a path in  $G$  and

$$|E(P')| = |E(Q_{1,1})| + |E(P)| + |E(Q_{2,1})| \geq |E(P)| + |E(P_1)| > |E(P_1)|.$$

Hence  $P'$  is a longer path than  $P_1$ , contradicting that  $P_1$  is a longest path.  $\square$

Things to remember:

1. Correctness
2. Clarity/Precision
3. Ease of Reading

# Colorings

---

## 1.1 Coloring and Brooks' Theorem

### coloring

A **coloring** of a graph  $G$  is an assignment of colors to vertices of  $G$  such that no two adjacent vertices receive the same color.

### k-coloring

Let  $G$  be a graph. We say  $\phi : V(G) \rightarrow [k]$  is a **k-coloring** of  $G$  if  $\phi(u) \neq \phi(v)$  for every  $uv \in E(G)$ .

Since every graph  $G$  has a  $|V(G)|$ -coloring, we are interested in the minimum numbers of colors needed to color  $G$ .

### chromatic number

The **chromatic number** of a graph  $G$ , denoted  $\chi(G)$ , is the minimum number  $k$  such that  $G$  has a  $k$ -coloring.

Then why coloring?

- Coloring is a foundational problem in graph labeling, wherein we study functions on  $V(G)$  according to constraints imposed on the graph (e.g. non-adjacent vertices are labeled differently)
- Coloring is a foundational problem in graph decomposition, wherein we seek to decompose  $V(G)$  into certain kinds of subgraphs (e.g. independent sets)
- Applications to maps, scheduling, job processing, frequency assignment (e.g. cell networks)
- Applications to algorithms (e.g. distributed computing)

However, coloring is hard.

A graph being an **independent set** is by definition equivalent to being **1-colorable**.

A graph being **bipartite** is by definition equivalent to being **2-colorable**. (Indeed coloring is a generalization of partite)

**Proposition 1.1**

$G$  is 2-colorable if and only if  $G$  does not contain an odd cycle.

Moreover, there exists a poly-time algorithm to decide if  $G$  is 2-colorable.

**Theorem: Karp (1972)**

For each  $k \geq 3$ , deciding if a graph  $G$  has a  $k$ -coloring is NP-complete.

Indeed, 3-coloring is NP-complete even for planar graphs. Any constant factor approximation is also NP-complete.

Then what about the bounds on chromatic number?

As mentioned  $\chi(G) \leq |V(G)|$ .

**Greedy Upper bound:**  $\chi(G) \leq \Delta(G) + 1$ , where  $\Delta(G)$  denotes the maximum degree of vertices in  $G$ . Why? By a greedy algorithm:

- Order the vertices of  $G$  arbitrarily,  $v_1, \dots, v_{|V(G)|}$ .
- Color the vertices in order avoiding the colors of previously colored neighbors.
- Since each vertex has at most  $\Delta(G)$  neighbors, there is always at least one color for the current vertex.

**Lower bound:**  $\chi(G) \geq \omega(G)$ , where  $\omega(G)$  denotes the clique number of  $H$ , that is the maximum size of a clique in  $G$ .

*Can we do better than the greedy upper bound?*

No! The bound is tight for complete graphs:  $\omega(K_n) = \chi(K_n) = (n - 1) + 1 = \Delta(K_n) + 1$ .

*Can we do better if the graph is not complete?*

No! The graph could have a component that is complete.

*Can we do better if the graph is connected and not complete?*

No! The bound is tight for odd cycles:  $\chi(C_{2k+1}) = 3 = 2 + 1 = \Delta(C_{2k+1}) + 1$ .

Can we do better if the graph is connected and neither complete nor an odd cycle? **Yes!**

**Theorem 1.2: Brooks 1941**

If  $G$  is connected, then  $\chi(G) \leq \Delta(G)$  if and only if  $G$  is neither complete nor an odd cycle.

**An Informal Proof of Brooks' Theorem**

How to prove Brook's Theorem?

Actually there are 8 to 10 distinct ways to prove Brooks' Theorem. See the nice survey *Brooks' Theorem and Beyond* by Cranston and Rabern from 2014 for more details. Here are some of those methods: Greedy Coloring, Kempe Chains, List Coloring, Alon-Tarsi Theorem, Kernel Perfection, Potential Method.


Today we give an informal proof sketch via the Greedy Coloring Method - arguably the most direct, brute-force of the approaches. (See Diestel for the Kempe Chain proof).

The idea is to try a method (greedy coloring) we know works for a similar problem ( $\Delta + 1$ -coloring), and ask under what conditions can we use this to get the desired outcome (a  $\Delta$ -coloring).

In the other cases we cannot apply greedy, we instead do **reductions**: that is, we show how to inductively color or to show that the graph is one of the exceptional outcomes (clique or odd cycle).

Alternatively, we could have built up a suite/library of reductions that work, and then tried to find a method to deal a finishing blow (i.e. to handle the cases we could not reduce).

**First Reduction**  $G$  has a cutvertex  $v$ . Then  $v$  separates  $G$  into two smaller graphs  $G_1$  and  $G_2$ .




By minimality of  $G$ ,  $G_i$  has a  $\Delta$ -coloring  $\phi_i$ ,  $i \in 1, 2$ .

This only works if neither graph is  $K_{\Delta+1}$  or odd cycle when  $\Delta = 2$ .

Now permute the colors in  $\phi_2$  so that  $\phi_1(v) = \phi_2(v)$ . Then  $\phi_1 \cup \phi_2$  yields a  $\Delta$ -coloring of  $G$ , a contradiction.


**Second Reduction**  $G$  has a cutset  $\{u, v\}$ .

Try the same trick. Say  $\{u, v\}$  separates  $G$  into two smaller graphs  $G_1$  and  $G_2$ . By induction or minimum counterexample, each of  $G_1, G_2$  has a  $\Delta$ -coloring  $\phi_i$ ,  $i \in 1, 2$ .




If  $uv \in E(G)$ , then we can permute the colorings so that  $\phi_1(u) = \phi_2(u)$  and  $\phi_1(v) = \phi_2(v)$ .

This fails if  $uv \notin E(G)$ . Because we may have  $u, v$  colored the same in one coloring and different in the other and no permuting will fix this! So we can add the edge  $uv$  to both  $G_1$  and  $G_2$ !




Have to show  $\Delta(G_1 + uv), \Delta(G_2 + uv) \leq \Delta(G)$ . We also have to ensure that neither  $G_1$  nor  $G_2$  is complete (or odd cycle in  $\Delta(G) = 2$  case).

Then we assume  $G$  is 3-connected. We now turn to the finishing blow (greedy). The greedy *fails* when a vertex has  $\Delta(G)$  earlier neighbors in the ordering, each with a different color from  $\{1, \dots, \Delta(G)\}$ .




Can we find an ordering where most of the vertices have at most  $\Delta(G) - 1$  earlier neighbors? Yes for all but the last vertex in the ordering! We can fix a root, then take a depth-first search tree ordering from the root. Reverse it!




Now all vertices but the last will be fine in greedy.

If  $\deg(v) \leq \Delta(G) - 1$ , then we can ensure greedy does not fail at the last vertex  $v$ . Otherwise, we ensure that two of its neighbors  $x$  and  $y$  are colored the same (and hence there is a color left for  $v$  when it is  $v$ 's turn). These two are two non-adjacent neighbors, which guaranteed to exist as  $G$  is not  $K_{\Delta+1}$ .





We can put  $x, y$  first in the ordering to guarantee  $x$  and  $y$  are colored the same. Then we can color them as we desire (since non-adjacent), say both with color 1.



Use the reverse of a depth-first search tree ordering of  $G - \{x, y\}$  with root  $v$ , then we finish the ordering so every vertex in  $V(G) \setminus \{x, y, v\}$  has at most  $\Delta(G) - 1$  earlier neighbors. Since  $G - \{x, y\}$  is connected as  $G$  is 3-connected, then this ordering exist.

### A Formal Proof of Brooks' Theorem

Let us codify our ordering fact as a proposition.

#### Proposition 1.3: Ordering Proposition

If  $G$  is a connected graph on  $n$  vertices and  $v \in V(G)$ , then there exists an ordering  $v_1, \dots, v_n = v$  of  $V(G)$  such that  $|N(v_i) \cap \{v_{i+1}, \dots, v_n\}| \geq 1$  for all  $i \in [n - 1]$ .

#### Proof:

Reverse a depth-first search tree ordering from root  $v$ . Or more formally:

We proceed by induction on  $|V(G)|$ . If  $|V(G)| = 1$ , then the ordering  $v$  is as desired. So we assume that  $|V(G)| \geq 2$ . Let  $G_1, \dots, G_k$  be the components of  $G - v$ . As  $G$  is connected, there exists neighbors  $u_1, \dots, u_k$  of  $v$  such that  $u_i \in V(G_i)$  for each  $i \in [k]$ . For each  $i \in [k]$ , there exists by induction applied to  $G_i$  and  $u_i$ , an ordering  $\sigma_i$  of  $V(G_i)$  as prescribed by the proposition. Let  $\sigma$  be the ordering of  $V(G)$  obtained by concatenating the  $\sigma_i$  and finally  $v$ . Then  $\sigma$  is as desired.  $\square$

Now we are ready to prove Brooks' Theorem:

Suppose not. Let  $G$  a counterexample with  $|V(G)|$  minimized. If  $\Delta(G) \leq 2$ , the result is standard. So we assume that  $\Delta(G) \geq 3$ .

**Claim 1** There does not exist a cutvertex of  $G$ .

#### Proof:

Suppose not. That is, there exists a cutvertex  $v$  of  $G$  and two connected subgraphs  $G_1, G_2$  of  $G$  such that  $G_1 \cap G_2 = \{v\}$ ,  $G_1 \cup G_2 = G$  and  $|V(G_i)| < |V(G)|$  for each  $i \in [2]$ .

As  $G_1$  and  $G_2$  are subgraphs of  $G$ , we have that  $\Delta(G_i) \leq \Delta(G)$  for each  $i \in [2]$ . Moreover, as  $G$  is connected, we have for each  $i \in [2]$  that  $\deg_{G_i}(v) \geq 1$  and hence  $\deg_{G_i}(v) \leq \Delta(G) - 1$ . Hence  $G_i \neq K_{\Delta(G)+1}$  for each  $i \in [2]$ . Thus by the minimality of  $G$ , there exist  $\Delta(G)$ -colorings  $\phi_i$  of  $G_i$  for each  $i \in [2]$ .

By permuting the colors of  $\phi_2$  as necessary, we assume without loss of generality that  $\phi_1(v) = \phi_2(v)$ . But then  $\phi_1 \cup \phi_2$  is a  $\Delta(G)$ -coloring of  $G$ , a contradiction.  $\square$

**Claim 2** There does not exist a 2-cut of  $G$ , or, there exists a vertex  $v \in V(G)$  with  $\deg_G(v) \leq \Delta(G) - 1$ .

#### Proof:

Suppose not. Now let us suppose there exists a 2-cut  $\{v_1, v_2\}$  of  $G$  and two connected subgraphs  $G_1, G_2$  of  $G$  such that  $G_1 \cap G_2 = \{v_1, v_2\}$ ,  $G_1 \cup G_2 = G$  and  $|V(G_i)| < |V(G)|$  for each  $i \in [2]$ .

Choose  $v_1, v_2, G_1, G_2$  such that neither  $G_1 + v_1v_2$  nor  $G_2 + v_1v_2$  is equal to  $K_{\Delta(G)+1}$  if possible.

As  $G$  is connected and  $G$  does not have a cutvertex by Claim 1, we have for all  $i, j \in [2]$  that  $\deg_{G_i}(v_j) \geq 1$  and hence  $\deg_{G_i}(v_j) \leq \Delta(G) - 1$ . Thus  $\Delta(G_i + v_1v_2) \leq \Delta(G)$  for all  $i \in [2]$ .

Next suppose that there exists  $i \in [2]$  such that  $G_i + v_1v_2 = K_{\Delta(G)+1}$ . Without loss of generality, we assume that  $i = 1$ . Let  $v'_1$  be the neighbor of  $v_1$  in  $G_2 - v_2$ . Let  $G_1 = G_1 + v_1v'_1$  and  $G'_2 = G_2 \setminus \{v_1\}$ . Now with  $\deg_G(v'_1) \leq \Delta(G) - 1$ , a contradiction, or we find that  $G'_i + v'_1v_2 \neq K_{\Delta(G)+1}$  for each  $i \in [2]$ . But then  $v'_1, v_2, G'_1, G'_2$  contradict the choice of  $v_1, v_2, G_1, G_2$ .

So we assume that  $G_1 + v_2v_2, G_2 + v_1v_2 \neq K_{\Delta(G)+1}$ . Thus by the minimality of  $G$ , there exist  $\Delta(G)$ -colorings  $\phi_i$  of  $G_i$  for each  $i \in [2]$ . By permuting the colors of  $\phi_2$  as necessary, we assume without loss of generality that  $\phi_1(v_j) = \phi_2(v_j)$  for each  $j \in [2]$ . But then  $\phi_1 \cup \phi_2$  is a  $\Delta(G)$ -coloring of  $G$ , a contradiction.  $\square$

Let  $v \in V(G)$  with  $\deg_G(v)$  minimized.

First suppose that  $\deg_G(v) \leq \Delta(G) - 1$ . By the Ordering Proposition, there exists an ordering  $v_1, \dots, v$  of  $V(G)$  such that  $|N(v_i) \cap \{v_{i+1}, \dots, v\}| \geq 1$  for all  $i \in [|V(G)| - 1]$ . Now greedily color  $V(G)$  in that order. This yields a  $\Delta(G)$ -coloring of  $G$ , a contradiction.

So we assume that  $\deg_G(v) = \Delta(G)$ . Since  $G \neq K_{\Delta+1}$ , there exist distinct  $x, y \in N(v)$  such that  $xy \notin E(G)$ . By Claims 1 and 2, it follows that  $G$  is 3-connected and hence  $G - \{x, y\}$  is connected. Hence by the Ordering Proposition, there exists an ordering  $v_1, \dots, v$  of  $V(G) - \{x, y\}$  such that  $|N(v_i) \cap \{v_{i+1}, \dots, v\}| \geq 1$  for all  $i \in [|V(G)| - 3]$ . Now color  $x, y$  with color 1. Then greedily color  $V(G) - \{x, y\}$  in that order. This yields a  $\Delta(G)$ -coloring of  $G$ , a contradiction.

## Beyond Brooks' Theorem

Can we go further? Can we save more colors? Under what conditions?

### Question ( $\omega, \Delta, \chi$ paradigm)

What is the maximum chromatic number of graphs with  $\omega(G) \leq \omega$  and  $\Delta(G) \leq \Delta$ ?

### Brooks' Reformulated

If  $G$  is a graph with  $\Delta(G) \geq 3$  and  $\omega(G) \leq \Delta(G)$ , then  $\chi(G) \leq \Delta(G)$ .

### Borodin-Kostochka Conjecture (1977)

If  $G$  is a graph with  $\Delta(G) \geq 9$  and  $\omega(G) \leq \Delta(G) - 1$ , then  $\chi(G) \leq \Delta(G) - 1$ .

Why  $\Delta \geq 9$ ?

Let  $G = C_5 \boxtimes K_3$ . (the blowup of every vertex in  $C_5$  to a triangle  $K_3$ ) Then  $\Delta(G) = 8$ ,  $\omega(G) = 6$ , and yet  $\chi(G) = 8$ .


### Theorem (Reed 1999)

True for  $\Delta(G) \geq 10^{14}$ .

## Reed's conjecture

### Reed's Conjecture (1998)

$$\chi(G) \leq \left\lceil \frac{\Delta(G) + 1 + \omega(G)}{2} \right\rceil.$$



5-cycle blowup

$$\begin{aligned}\Delta &= 3t - 1 \\ \omega &= 2t\end{aligned}$$

$$\left\lceil \frac{1}{2}(\Delta + 1 + \omega) \right\rceil = \left\lceil \frac{5t}{2} \right\rceil$$

$$\alpha = 2$$

### Theorem (Reed 1998)

The conjecture holds when  $\Delta(G)$  is sufficiently large and

$$\omega(G) \geq (1 - 7 \cdot 10^{-7})\Delta(G).$$

### Corollary (Reed)

There exists  $\varepsilon > 0$  such that for every graph  $G$ ,

$$\chi(G) \leq (1 - \varepsilon)(\Delta(G) + 1) + \varepsilon\omega(G).$$

Reed's value of  $\varepsilon$  was  $10^{-8}$ .

Can we improve the  $\varepsilon$  for large enough  $\Delta$ ? Can we get closer to  $\varepsilon = 1/2$ ?

For Large enough  $\Delta$ , the following  $\varepsilon$  suffices:

- $\frac{1}{320e^6}$  (King and Reed 2012)
- $\frac{1}{26}$  (Bonamy, Perrett, Postle 2016+)
- $\frac{1}{13}$  (Delcourt and Postle 2017+)
- $\frac{1}{8.4}$  (Hurley, de Joannis de Verclos, Kang 2020+)

## Large Girth

The **girth** of a graph  $G$  is the length of a shortest cycle in  $G$ .

### Theorem (Erdős 1959)

$\forall g, k \geq 1$ , there exists graphs of girth at least  $g$  and chromatic number at least  $k$ .

**Theorem (Frieze and Luczak 1992)**

Random  $d$ -regular graphs have chromatic number  $(1 - o(1)) \frac{d}{2 \ln d}$  with high probability.

**Corollary**

$\forall g, d \geq 1$ , there exists a  $d$ -regular graph  $G$  of girth at least  $g$  with

$$\chi(G) \geq (1 - o(1)) \frac{d}{2 \ln d}.$$

**Girth-Five and Triangle-Free****Theorem (Kim 1995)**

If  $G$  is a graph of girth five, then

$$\chi(G) \leq (1 + o(1)) \frac{\Delta(G)}{\ln \Delta(G)}.$$

**Theorem (Johansson 1996)**

If  $G$  is a triangle-free graph, then

$$\chi(G) \leq O\left(\frac{\Delta(G)}{\ln \Delta(G)}\right).$$

**Theorem (Molloy 2017)**

If  $G$  is a triangle-free graph, then

$$\chi(G) \leq (1 + o(1)) \frac{\Delta(G)}{\ln \Delta(G)}.$$

**Small Clique Number****Theorem (Johansson 1999)**

For every fixed  $r$ : if  $G$  is a graph with  $\omega(G) \leq r$ , then

$$\chi(G) \leq O\left(\frac{\Delta(G)}{\ln \Delta(G)} \cdot \ln \ln \Delta(G)\right).$$

**Theorem (Molloy 2017)**

$$\chi(G) \leq 200 \cdot \omega(G) \cdot \frac{\Delta(G)}{\ln \Delta(G)} \cdot \ln \ln \Delta(G).$$

Good for  $\omega(G) \leq \frac{\ln \Delta(G)}{\ln \ln \Delta(G)}$ . What if  $\omega(G)$  is larger?

**Question**

For  $k \geq 2$ , what value of  $\omega(G)$  guarantees  $\chi(G) \leq \frac{\Delta(G)}{k}$ ?

**Theorem (Bonamy, Kelly, Nelson, Postle 2018+)**

$$\chi(G) \leq O\left(\Delta(G) \cdot \sqrt{\frac{\ln \omega(G)}{\ln \Delta(G)}}\right).$$

**Corollary**

$\forall k \geq 2$ , if  $\omega(G) \leq \Delta(G)^{\frac{1}{(192k)^2}}$ , then

$$\chi(G) \leq \frac{\Delta(G)}{k}.$$

Ramsey theory constructions show that we cannot extend this beyond  $\Delta(G)^{\frac{2}{k-1}}$ .

## 1.2 Edge Coloring

**edge coloring**

An **edge-coloring** of a graph  $G$  is an assignment of colors to edges of  $G$  such that no two incident edges receive the same color.

**k edge coloring**

Let  $G$  be a graph. We say  $\phi : E(G) \rightarrow [k]$  is a **k-edge-coloring** of  $G$  if  $\phi(e) \neq \phi(f)$  for every  $e, f \in E(G)$  with  $e \sim f$ .

Here  $e \sim f$  means  $e, f$  share a common endpoint ("are adjacent") in  $G$ .

**chromatic index**

The **chromatic index** of a graph  $G$  (also known as **edge chromatic number**), denoted  $\chi'(G)$ , is the minimum number  $k$  such that  $G$  has a  $k$ -edge-coloring.

**line graph**

The **line graph** of a graph  $G$ , denoted by  $L(G)$ , is the graph where  $V(L(G)) := E(G)$  and  $E(L(G)) := \{ef : e, f \in E(G), e \sim f\}$ .

Edge colorings of  $G$  are equivalent to vertex colorings of  $L(G)$ . Hence  $\chi'(G) = \chi(L(G))$ .

What are some natural upper and lower bounds on  $\chi'(G)$ ?

**Proposition 1.4**

$$\Delta(L(G)) \leq 2\Delta(G) - 2$$

Hence by greedy,

$$\chi'(G) \leq 2\Delta(G) - 1$$

**Proposition 1.5**

$$\omega(L(G)) \geq \Delta(G)$$

Hence

$$\chi'(G) \geq \Delta(G)$$

Moreover,  $\omega(L(G)) = \Delta(G)$  if  $\Delta(G) \geq 3$ .

Note that for even cycles, namely  $K_n$ ,  $n$  even,  $\chi'(G) = \Delta(G)$ . For odd cycles, we have  $\chi'(G) > \Delta(G)$ .

**Theorem 1.6: König (1916)**

If  $G$  is a bipartite graph, then  $\chi'(G) = \Delta(G)$ .

**Proof (first):**

It suffices to prove the theorem when  $G$  is  $\Delta(G)$ -regular since every bipartite graph  $G$  is a subgraph of some  $\Delta(G)$ -regular graph  $H$ .

Prove by induction on  $\Delta(G)$ . If  $\Delta(G) = 0$ , then the statement holds trivially. So assume  $\Delta(G) \geq 1$ .

Let  $S \subseteq A$ . By double counting  $E(G(S, N(S)))$ , it follows that

$$\Delta(G)|S| = |E(G(S, N(S)))| \leq \Delta(G)|N(S)|,$$

and thus  $|S| \leq |N(S)|$ . Hence by Hall's theorem, there exists a perfect matching  $M$  of  $G$ .

By induction,  $G - M$  has a  $(\Delta(G) - 1)$ -coloring  $\phi$ . Let  $\phi(e) = \Delta(G)$  for each  $e \in M$ . Then  $\phi$  is a  $\Delta(G)$ -coloring of  $G$  as desired.  $\square$

**Kempe chain**

Let  $\phi$  be a partial  $k$ -edge-coloring of a graph  $G$ . If  $a, b \in [k]$  and  $v \in V(G)$ , then  $(a, b)$ -chain at  $v$  in  $\phi$ , denoted  $P_v(a, b, \phi)$  is the maximal path/cycle of  $a$  and  $b$  colored edges containing  $v$ .

**switching**

The coloring  $\phi'$  obtained from switching (aka **recoloring**)  $P_v(a, b, \phi)$  is defined as:

- $\phi'(e) = \{a, b\} \setminus \phi(e)$  if  $e \in P_v(a, b, \phi)$ , and
- $\phi'(e) = \phi(e)$  otherwise.

## missing colors

Let  $\phi$  be a partial  $k$ -edge coloring of a graph  $G$ .

- A coloring  $a \in [k]$  is missing at  $v$  in  $\phi$  if  $a \notin \{\phi(e) : e \sim v\}$ .
- We let  $\phi(v)$  denote the set of missing colors at  $v$ .

## Proof (second proof of König):

We proceed by induction on  $|E(G)|$ . If  $E(G) = \emptyset$ , there is nothing to show. So we assume that  $E(G) \neq \emptyset$ .

let  $e = uv \in E(G)$ . By induction, there exists a  $\Delta(G)$ -edge-coloring of  $G - e$ . Let  $\phi$  be a  $\Delta(G)$ -edge-coloring of  $G - e$  such that  $|\phi(u) \cap \phi(v)|$  is maximized.

Note that  $\phi(u), \phi(v) \neq \emptyset$  since  $\deg_{G-e}(u), \deg_{G-e}(v) \leq \Delta(G) - 1$ .

If  $\phi(u) \cap \phi(v) \neq \emptyset$ , then let  $\phi(e) \in \phi(u) \cap \phi(v)$  and hence  $\phi$  is a  $\Delta(G)$ -edge-coloring of  $G$  as desired.

Now assume that  $\phi(u) \cap \phi(v) = \emptyset$ . Let  $a \in \phi(u), b \in \phi(v)$ . Note that  $P := P_u(a, b, \phi)$  is a path. If  $v \in V(P)$ , then it follows that  $P$  has even length and hence  $P + e$  is an odd cycle in  $G$ , contradicting that  $G$  is bipartite.

So we assume that  $v \notin V(P)$ . But then switching  $P$  yields a coloring  $\phi'$  such that  $b \in \phi'(u) \cap \phi'(v)$ , contradicting the choice of  $\phi$ .  $\square$

## 1.2.1 Vizing's Theorem

## Theorem 1.7: Vizing (1964)

If  $G$  is a graph, then  $\chi'(G) \leq \Delta(G) + 1$ .


A graph  $G$  is called **class 1** if  $\chi'(G) = \Delta(G)$ , or **class 2** if  $\chi'(G) = \Delta(G) + 1$ . Note that deciding if a graph is class 1 is NP-complete!

## Vizing fan

Suppose that  $G$  is a graph,  $e = v_0v_1 \in E(G)$ , and  $\phi$  is a partial  $k$ -edge-coloring of  $G - e$  for some integer  $k$ . We say  $T = (v_0e_1v_1e_2 \dots e_nv_n)$  is a **Vizing fan** with respect to the edge  $e$ , vertex  $v_0$  and the coloring  $\phi$  if

- $v_0, v_1, v_2, \dots, v_n$  are all disjoint, and
- $\forall j, 1 \leq j \leq n, e_j = v_jv_0$ , and
- $\forall j, 2 \leq j \leq n, \phi(e_j) \in \bigcup_{i < j} \phi(v_i)$ .

Here is a depiction of Vizing fan.



Idea is no coloring implies disjoint missing colors in Vizing fan.

**Lemma 1.8: Disjoint missing colors**

Let  $G$  be a graph and  $e = v_0v_1 \in E(G)$  such that for some integer  $k \geq \Delta(G) + 1$ ,  $G - e$  has a  $k$ -edge-coloring, but  $G$  does not.


If  $\phi$  is a  $k$ -edge-coloring of  $G - e$ , and  $T = (v_0e_1v_1e_2v_2 \dots e_nv_n)$  is a Vizing fan with respect  $e, v_0$  and  $\phi$ , then  $\phi(v_i) \cap \phi(v_j) = \emptyset$  for all distinct  $v_i, v_j \in V(T)$ .

**Proof:**

Suppose not. That is  $\exists i < j \in \{0, \dots, n\}$  such that  $\phi(v_i) \cap \phi(v_j) \neq \emptyset$ .


Let  $\phi, T, i, j$  be chosen such that  $j$  is minimized, and subject to that condition,  $i$  is minimized. Let  $\alpha \in \phi(v_i) \cap \phi(v_j)$ . Three cases:  $i = 0$  and  $j = 1$ ;  $i = 0$  and  $j > 1$ ;  $i > 0$ .

Case 1:  $i = 0, j = 1$ .



Let  $\phi(e) = \alpha$  and hence  $\phi$  is a  $k$ -edge-coloring of  $G$ , a contradiction.


Case 2: Let  $\beta = \phi(v_0v_j)$ . Since  $T$  is a Vizing fan,  $\exists j' < j$  such that  $\beta \in \phi(v_{j'})$ .



Let  $\phi'$  be obtained from  $\phi$  switching  $P_{v_0}(\alpha, \beta, \phi) = v_0v_j$ . Then  $\beta \in \phi'(v_0) \cap \phi'(v_{j'})$ . Now  $T' := T[\{v_0, \dots, v_{j'}\}]$  and  $\phi'$  contradict minimality of  $T$  and  $\phi$ .

Case 3:  $i > 0$ . Let  $\beta \in \phi(v_0)$ . By minimality of  $T$ ,  $\beta \neq \alpha$ . Let  $j' := i$ .


(a)  $v_0 \notin V(P_{v_i}(\alpha, \beta, \phi))$ .





Let  $\phi'$  be obtained from  $\phi$  by switching  $P_{v_i}(\alpha, \beta, \phi)$ . Then  $\beta \in \phi'(v_0) \cap \phi'(v_i)$ . Now  $T' := T[\{v_0, \dots, v_i\}]$  and  $\phi'$  contradict the minimality of  $T$  and  $\phi$ .

(b)  $v_0 \in V(P_{v_i}(\alpha, \beta, \phi))$ .



Let  $\phi'$  be obtained from  $\phi$  by switching  $P_{v_j}(\alpha, \beta, \phi)$ . Then  $\beta \in \phi'(v_0) \cap \phi'(v_j)$ . Now  $T' := T[\{v_0, \dots, v_j\}]$  and  $\phi'$  contradict the minimality of  $T$  and  $\phi$ . □

Now we can take maximum Vizing fan and apply lemma for contradiction.

#### Proof of Vizing's Theorem:

Let  $G$  be a counterexample with  $|V(G)|$  minimized. Hence  $E(G) \neq \emptyset$ . Let  $e = v_0v_1 \in E(G)$ . By the minimality of  $G$ ,  $G - e$  has  $(\Delta(G) + 1)$ -edge-coloring  $\phi$  of  $G - e$ .

Let  $T = (v_0e_1v_1e_2v_2 \dots e_nv_nv_n)$  be a Vizing fan with respect to  $e, v_0$  and  $\phi$  such that  $n$  is maximized. By lemma,  $\phi(v_i) \cap \phi(v_j) = \emptyset$  for all distinct  $i, j \in \{0, \dots, n\}$ .

Let  $X := \bigcup_{i \in [n]} \phi(v_i)$ . By Lemma,  $|X| \geq n$ . So  $\exists \alpha \in X$  such that  $\exists f = v_0v_j, j \in [n]$  with  $\phi(f) = \alpha$ . Since  $\alpha \notin \phi(v_0)$ ,  $\exists e_{n+1} = v_0v_{n+1}$  with  $\phi(e_{n+1}) = \alpha$ . But then  $T + e_{n+1}$  is a larger Vizing fan with respect to  $e, v_0, \phi$ , contradicting the maximality of  $T$ . □

## 1.2.2 List Edge Coloring

List Coloring:

- list-assignment  $L$ : an assignment of lists  $L(v)$  for  $v \in V(G)$ .
- $k$ -list-assignment  $L$ :  $|L(v)| \geq k$  for all  $v \in V(G)$ .
- $L$ -coloring: a coloring  $\phi$  where  $\phi(v) \in L(v)$  for all  $v \in V(G)$ .
- A graph  $G$  is  $k$ -list-colorable if  $G$  has an  $L$ -coloring for every  $k$ -list-assignment  $L$ .

### list chromatic number

The **list chromatic number**, denoted  $\chi_\ell(G)$ , is the minimum  $k$  such that  $G$  has an  $L$ -coloring for every  $k$ -list-assignment  $L$ .

### Proposition 1.9

For all integer  $k \geq 0$ , there exists a bipartite graph  $G$  with  $\chi_\ell(G) = k$ .

### Theorem (Alon 2000)

If  $G$  has average degree  $d$ , then  $\chi_\ell(G) \geq \Omega(\log d)$ .

For edge-coloring, we have something similar.

#### list chromatic index

The **list chromatic index**, denoted  $\chi'_\ell(G)$ , is  $\chi_\ell(L(G))$ .

#### List Coloring Conjecture (Various authors, 1970s/80s)

If  $G$  is a graph, then  $\chi'_\ell(G) = \chi'(G)$ .

#### Theorem 1.10: Galvin, 1995

If  $G$  is a bipartite graph, the  $\chi'_\ell(G) = \chi'(G)$ .

#### Theorem (Kahn, 1996)

If  $G$  is a graph, then  $\chi'_\ell(G) = (1 + o(1))\chi'(G)$ .

#### latin square

A **latin square** is an  $n$  by  $n$  array such that each of the numbers 1 to  $n$  appears exactly once in each row and exactly once in each column.

Equivalently, a latin square is an  $n$ -edge-coloring of  $K_{n,n}$ . Such always exist by König's theorem which shows that  $\chi'(K_{n,n}) = \Delta(K_{n,n}) = n$ .


#### Dinitz Conjecture (1978)

Given an  $n$  by  $n$  array and an assignment of  $n$  symbols to each square, there exists a choice of symbol for each square such that each symbol appears at most once in each row and each column.

Or equivalently, by Galvin's Theorem,  $\chi'_\ell(K_{n,n}) = n$ .

#### kernel

A **kernel** of a digraph  $D$  is an independent set  $U$  such that every vertex in  $D \setminus U$  has an out-neighbor in  $U$ .



#### kernel perfect

We say an orientation  $D$  of a graph  $G$  is **kernel perfect** if every induced subgraph  $D'$  of  $D$  has a kernel.

**Lemma 1.11: Kernel Perfect Lemma**

Let  $G$  be a graph and  $L$  a list assignment. If  $G$  has a kernel perfect orientation  $D$  such that  $d^+(v) < |L(v)|$  for every  $v$  in  $D$ , then  $G$  has an  $L$ -coloring.

**Proof:**


By induction on  $|V(G)|$ . If  $|V(G)| = 0$ , we are done.

Let  $v \in V(G)$ ,  $\alpha \in L(v)$  and  $D' := G[\{u \in V(G) : \alpha \in L(u)\}]$ . Since  $D$  is kernel perfect,  $D'$  has a kernel  $U$ . Let  $\phi(u) = \alpha$  for all  $u \in U$ , and  $L'(x) := L(x) \setminus \{\alpha\}$  for all  $x \in G \setminus U$ . Now  $d_{G \setminus U}^+(v) < |L'(v)|$  for all  $v \in D \setminus U$ . By induction,  $G \setminus U$  has an  $L'$ -coloring and hence  $G$  has an  $L$ -coloring.  $\square$

**Proof of Galvin's Theorem:**

Let  $G = (A, B)$ . Let  $k = \chi'(G) = \Delta(G)$ . Let  $\phi$  be a  $k$ -edge-coloring of  $G$ .

Let  $D$  be an orientation of  $L(G)$  where  $e \sim e' \in E(G)$  with  $\phi(e) < \phi(e')$ , we orient  $ee'$ :  $e' \rightarrow e$  if  $e \cap e' \in A$ ;  $e \rightarrow e'$  if  $e \cap e' \in B$ .



Let  $e = uv$ ,  $u \in A, v \in B$ . Let  $\phi(e) = i$ . Then

$$\begin{aligned} d^+(e) &\leq |\{e' : e' \cap e = u, \phi(e') < \phi(e)\}| + |\{e' : e' \cap e = v, \phi(e') > \phi(e)\}| \\ &\leq (i-1) + (\Delta(G) - i) = \Delta(G) - 1 = k - 1 \end{aligned}$$

Let  $D'$  be an induced subgraph of  $D$ . Then  $D'$  has a kernel  $U$ , namely a stable matching as guaranteed by the Stable Marriage Theorem (where  $v$  prefers  $u$  to  $u'$  if  $u'v$  is directed towards  $uv$ ). Hence  $D$  is kernel perfect. By Kernel Perfect Lemma,  $L(G)$  has an  $L$ -coloring.  $\square$

**Proof Ideas for Kahn's Theorem:**

Randomly color edges; Uncolor incident edges with same color; iterate; Finish with a well chosen reserve of colors.  $\square$

Molly and Reed:  $\chi'_\ell(G) = \Delta(G) + O(\Delta(G)^{1/2} \log^4 \Delta(G))$

Kahn: Also holds for edge-coloring  $k$ -uniform linear hypergraphs. (A hypergraph is linear if any two vertices are contained in a most one hyperedge.)

**1.2.3 Edge Coloring Multigraphs**


If  $G$  is a multigraph, let  $\chi'(G)$  denote  $\chi(L(G))$ . Note that  $L(G)$  is well-defined for multigraphs. Also note that we allow parallel edges but not loops, so that  $L(G)$  is simple. We have a trivial bound:

$$\Delta(G) \leq \omega(L(G)) \leq \chi'(G) \leq \Delta(L(G)) + 1 \leq 2\Delta(G) - 1$$

**Theorem 1.12: König, 1916**

If  $G$  is a bipartite multigraph, then  $\chi'(G) = \Delta(G)$ .

Same proof works. However, Vizing's Theorem does not hold.

**Theorem 1.13: Shannon, 1949**

If  $G$  is a multigraph, then  $\chi'(G) \leq \left\lfloor \frac{3\Delta(G)}{2} \right\rfloor$ .

We can't do better because it is tight for triangle.

**multiplicity**

The **multiplicity** of a multigraph  $G$ , denoted  $\mu(G)$ , is the maximum number of pairwise parallel edges in  $G$ .

**Theorem 1.14: Vizing, 1964**

If  $G$  is a multigraph, then  $\chi'(G) \leq \Delta(G) + \mu(G)$ .

**Proof:**


We carry over all definitions to multigraphs: Kempe chains, switching, missing colors, Vizing fan. Same proof works for disjoint missing color lemma in the context of multigraphs. Then we fix the finish as follows:

Let  $k = \Delta(G) + \mu(G)$ . Let  $X := \bigcup_{i \in [n]} \phi(v_i)$ . By lemma, we have  $|X| \geq (k - \Delta(G))n = \mu(G)n$ . There are at most  $\mu(G)n - 1$  colored edges incident with  $v_0$  and another vertex in  $T$ . So  $\exists \alpha \in X$  such that  $\nexists f = v_0 v_j, j \in [n]$  with  $\phi(f) = \alpha$ . Since  $\alpha \notin \phi(v_0)$ ,  $\exists e_{n+1} = v_0 v_{n+1}$  with  $\phi(e_{n+1}) = \alpha$ . But then  $T + e_{n+1}$  is a larger Vizing fan with respect to  $e, v_0$  and  $\phi$ , contradicting the maximality of  $T$ .  $\square$

**Proof of Shannon's Theorem:**

We proceed by induction. If  $\mu(G) \leq \left\lfloor \frac{\Delta(G)}{2} \right\rfloor$ , then desired outcome follows from Vizing's Theorem for multigraphs.

So we assume  $\mu(G) > \frac{\Delta(G)}{2}$ . Let  $u, v \in V(G)$  such that there exist  $\mu(G)$  parallel edges with ends  $u, v$ . Let  $G'$  be the multigraph obtained from  $G$  by identifying  $u$  and  $v$  to a new vertex  $w$  and deleting all loop incident with  $w$ .



Now  $\deg_{G'}(w) \leq 2(\Delta - \mu) \leq \Delta$ . Hence  $\Delta(G') \leq \Delta(G)$ . By induction,  $G'$  has a  $\left\lfloor \frac{3\Delta(G)}{2} \right\rfloor$ -coloring  $\phi$ . Extend  $\phi$  to  $G$  as desired. Note this is possible as

$$\mu(G) + \deg_{G'}(w) \leq 2\Delta - \mu(G) \leq \left\lfloor \frac{3\Delta(G)}{2} \right\rfloor$$

□

Let

$$p(G) := \max \left\{ \left\lfloor \frac{2|E(G[X])|}{|X| - 1} \right\rfloor \mid X \subseteq V(G) \right\}$$

Note that  $\chi'(G) \geq p(G)$ .

**Goldberg 1979, indep. Seymour 1977**

$$\chi'(G) \leq \max\{\Delta(G) + 1, p(G)\}$$

Edmonds (matching polytope theorem) shows that the fractional chromatic index satisfies

$$\chi'_f(G) = \max\{\Delta(G) + 1, p(G)\}.$$

So the Goldberg-Seymour conjecture is equivalent to saying that  $\chi'(G) = \chi'_f(G)$ .

#### Kierstead Path

Suppose  $G$  is a graph,  $e_1 = v_0v_1 \in E(G)$ ,  $S$  is a set of colors, and  $\phi$  is an  $S$ -coloring of  $G - e_1$ . We say  $T = (v_0e_1v_1e_2v_2 \dots e_nv_n)$  is a **Kierstead Path** with respect to the edge  $e_1$  and the coloring  $\phi$  if

- $v_0, v_1, v_2, \dots, v_n$  are all disjoint, and
- $\forall j, 1 \leq j \leq n, e_j = v_ju$  where  $u \in \bigcup_{i < j} \{v_i\}$  and  $u = v_{j-1}$ , and
- $\forall j, 2 \leq j \leq n, \phi(e_j) \in \bigcup_{i < j} \phi(v_i)$ .

#### Theorem (Kierstead)

The Disjoint missing colors lemma holds for Kierstead Paths.

#### Tashkinov Tree

Suppose  $G$  is a graph,  $e_1 = v_0v_1 \in E(G)$ ,  $S$  is a set of colors, and  $\phi$  is an  $S$ -coloring of  $G - e_1$ . We say  $T = (v_0e_1v_1e_2v_2 \dots e_nv_n)$  is a **Tashkinov Tree** with respect to the edge  $e_1$  and the coloring  $\phi$  if

- $v_0, v_1, v_2, \dots, v_n$  are all disjoint, and
- $\forall j, 1 \leq j \leq n, e_j = v_ju$  where  $u \in \bigcup_{i < j} \{v_i\}$ , and
- $\forall j, 2 \leq j \leq n, \phi(e_j) \in \bigcup_{i < j} \phi(v_i)$ .

**Theorem (Tashkinov 2000)**

The Disjoint missing colors lemma holds for Tashkinov Trees.

As time goes by, we have several upper bounds. The latest is

**Theorem (Chen, Gao, Kim, Postle, Shan 2018)**

$$\chi'(G) \leq \max \left\{ \Delta(G) + \left\lceil (\Delta(G)/2)^{1/3} \right\rceil, p(G) \right\}$$

### 1.3 Thomassen's Theorem

What is the maximum list chromatic number of planar graphs? It is at least 4 since  $K_4$  is planar, and at most 6 since planar graphs are 5-degenerate.

**k-degenerate**

A graph is **k-degenerate** if its vertices can be successively deleted so that when deleted, each has degree at most  $k$ . The degeneracy of a graph is the smallest  $k$  such that it is  $k$ -degenerate.

or

A  $k$ -degenerate graph is a graph in which every induced subgraph has a vertex with degree at most  $k$ .

**Conjecture (Erdős, Rubin and Taylor 1979)**

$\exists$  a planar graph with list chromatic number at least 5.

**Theorem (Voigt 1993)**

$\exists$  a planar graph with list chromatic number at least 5.

**Conjecture (Erdős, Rubin and Taylor 1979)**

Every planar graph has list chromatic number at most 5.

**Theorem (Thomassen 1994)**

Every planar graph has list chromatic number at most 5.

How to prove there exists a 5-list-coloring? Identification and Kempe chains do not work for list coloring, but we can prove something stronger.


**Theorem 1.15: Thomassen's Stronger Theorem**

Let  $G$  be a connected plane graph,  $C$  be the boundary walk of the infinite face of  $G$ , and  $P$  be a path on at most two vertices in  $C$ .

If  $L$  is a list assignment of  $G$  such that

- $|L(p)| = 1 \ \forall p \in V(P)$ ,
- $|L(v)| \geq 3 \ \forall v \in V(C) \setminus V(P)$ ,
- $|L(w)| \geq 5 \ \forall w \in V(G) \setminus V(C)$ , and
- $G[V(P)]$  has an  $L$ -coloring,

then  $G$  has an  $L$ -coloring.




**Proof:**

Let  $G$  be a counterexample with  $|V(G)|$  minimized. Assume WLOG that  $|V(P)| = 2$ . Let  $P = p_1 p_2$ .


**Global reduction: cutvertex/chord**

Assume  $G$  has a cutvertex:  $G_1 \cap G_2 = \{v\}$ ,  $G_1 \cup G_2 = G$ . Assume WLOG that  $V(P) \subseteq V(G_1)$ .



By minimality,  $G_1$  has an  $L$ -coloring  $\phi$ . By minimality,  $\phi$  extends to an  $L$ -coloring of  $G_2$ .


Assume  $G$  has a chord:  $G_1 \cap G_2 = \{uv\}$ ,  $G_1 \cup G_2 = G$ . Assume WLOG  $V(P) \subseteq V(G_1)$ .



By minimality,  $G_1$  has an  $L$ -coloring  $\phi$ . By minimality,  $\phi$  extends to an  $L$ -coloring of  $G_2$ .

**Local Reduction**


Let  $v \neq p_2$  be neighbor of  $p_1$  in  $C$ . Let  $\{c_1, c_2\} \in L(v) \setminus L(p_1)$ .



Let  $G' = G - v$ , let  $v' \neq p_1$  be the neighbor of  $v$  in  $C$  and

$$L'(w) = \begin{cases} L(w) \setminus \{c_1, c_2\} & w \in N(v) \setminus \{v'\} \\ L(w) & \text{otherwise} \end{cases}$$

Let  $C'$  be the outer face of  $G'$ .



Since  $G$  is 2-connected,  $G'$  is connected. Note that  $|L'(w)| \geq 3 \forall w \in V(C') \setminus V(P)$  because if  $w \in N(v) \setminus \{p_1, v'\}$ , then  $w \in V(G) \setminus V(C)$  since  $C$  has no chord.  $|L'(w)| \geq 5 \forall w \in V(G') \setminus V(C')$  since if  $w \in N(v) \setminus V(C)$ , then  $w \in V(C')$ . By minimality of  $G$ ,  $\exists L'$ -coloring  $\phi$  of  $G'$ . Now let  $\phi(v) \in \{c_1, c_2\} \setminus \phi(v')$ .  $\square$

## 1.4 Coloring and List Coloring Planar Graphs

In 1852, Guthrie proposed Four Color Conjecture: Every planar graph is 4-colorable. It's equivalent formulation: Every bridgeless cubic planar graph is 3-edge colorable. The four color theorem was proved in 1976 by Kenneth Appel and Wolfgang Haken after many false proofs and counterexamples. Uses the method of discharging to show that every minimum counterexample contains one of 1834 unavoidable configurations. Show that each of these configurations is reducible (i.e. does not occur in a minimum counterexample). Second shorter proof by Robertson, Sanders, Seymour and Thomas in 1997 (only 633 configurations). Latter proof verified by formal proof software Coq in 2005. Interestingly, we can use fewer colors by excluding certain subgraphs.

Grötzsch's Theorem (1959) states that every triangle-free graph is 3-colorable. This does not extend to list coloring: Voigt (1995) states that  $\exists$  a planar triangle-free graph that is not 3-list-colorable. In 1995, Thomassen's theorem states that every planar graph of girth at least five (i.e., no  $\triangle$  or 4-cycles) is 3-colorable.

In 1976, Steinberg proposed a conjecture: Every planar graph without 4-cycles or 5-cycles is 3-colorable. Erdős, in 1991, proposed a question: What is the smallest  $k$  such that planar graphs with no cycles of length 4 to  $k$  are 3-colorable? Borodin, Glebov, Raspaud, Salavatipour (2005) showed  $k = 7$  works. Cohen-Addad, Hebdige, Král', Li, Salgado (2016) showed Steinberg's conjecture is false. Dvořák, Postle, in 2018: planar and no 4 to 8 cycles is 3-list-colorable. But we have more questions on planar graphs:

- If a theorem guarantees one  $k$ -coloring, could we also prove a theorem guaranteeing many  $k$ -colorings?
- If a theorem guarantees one  $k$ -coloring, could we also prove a theorem guaranteeing a  $k$ -coloring if we add a few anomalies (e.g. precolored vertices, crossings, etc.)?

Not all planar graphs have (exponentially) many 4-colorings. Repeatedly adding new degree three



vertices inside facial triangles of  $K_4$  has only one 4-coloring up to permutation of colors. However, all planar graphs have (exponentially) many 5-colorings.

### Proposition 1.16

Every graph has at least  $2^{\frac{|V(G)|}{\chi(G)}} (\chi(G) + 1)$ -colorings.

#### Proof:

Take the largest color class in a  $\chi(G)$ -coloring and recolor any subset with the  $\chi(G) + 1$ -st color.  $\square$

These results can extend to list colorings. In 2007, Thomassen proved that If  $G$  is a planar graph and  $L$  is a 5-list-assignment of  $G$ , then  $G$  has at least  $2^{\frac{|V(G)|}{9}}$  distinct  $L$ -colorings. In 1974, Aksenov proved that every planar graph with at most 3 triangles is 3-colorable. Havel's conjecture (1969) which states that  $\exists D > 0$  s.t.: every planar graph where every pair of triangles is at distance  $\geq D$  apart is 3-colorable, is proved by Dvořák, Král's and Thomas.

In 1997, Thomassen raised a question: Does  $\exists D > 0$  s.t.: If  $G$  is a planar graph and  $X$  is a set of vertices  $\geq D$  pairwise far apart, then every 5-coloring of  $X$  extends to  $G$ ?

### Proposition (Alberston 1998)

If  $G$  is a graph and  $X$  is a set of vertices  $\geq 4$  far apart, then every  $\chi(G) + 1$ -coloring of  $X$  extends to  $G$ .

#### Proof:

Take a  $\chi(G)$ -coloring  $\phi$  of  $G$ . Recolor every vertex  $x_i$  in  $X$  to its preferred color  $c_i$ . Recolor the neighbors of  $x_i$  colored  $c_i$  to color  $\chi(G) + 1$ . This yields a  $\chi(G) + 1$ -coloring of  $G$ , because  $N(x_i)$  are disjoint/non-adjacent as distance  $\geq 4$ .  $\square$

What about list coloring? By Thomassen's theorem: planar graphs with at most 2 precolored adjacent vertices are 5-list-colorable. What if the vertices are pairwise far apart? Alberston's Conjecture (1998):  $\exists D > 0$  s.t.: every planar graph where every pair of precolored vertices is at distance  $\geq D$  apart is 5-list-colorable, was proved by Dvořák et al. in 2017.

## 1.5 Discharging

Discharging is a counting method wherein:

- We assign charges to objects (e.g. vertices, edges, faces of a graph) such that the sum of the charges is negative (resp. positive)
- We redistribute the charge according to a set of discharging rules such that the sum is unchanged
- We derive a contradiction by showing the sum of the new charges is non-negative (resp. non-positive) given the assumed properties.

We use charges for plane graphs, natural choices come from Euler's formula. Recall Euler's formula:

### Euler's formula

If  $G$  is a plane graph, then

$$|V(G)| - |E(G)| + |\mathcal{F}(G)| = 1 + |\mathcal{C}(G)|$$

where  $\mathcal{F}(G)$  denote the set of faces of  $G$  and  $\mathcal{C}(G)$  denotes the set of components of  $G$ .

### 1.5.1 Common Discharging Setups for Planar Graphs

Let  $G$  be a plane graph. Initial charges:

$$\begin{aligned} \text{ch}_0(v) &= \deg(v) - 6 & \forall v \in V(G) \\ \text{ch}_0(f) &= 2(|f| - 3) & \forall f \in \mathcal{F}(G) \end{aligned}$$

Total sum of charges:

$$\begin{aligned} & \sum_{v \in V(G)} \text{ch}_0(v) + \sum_{f \in \mathcal{F}(G)} \text{ch}_0(f) \\ &= \sum_{v \in V(G)} (\deg(v) - 6) + \sum_{f \in \mathcal{F}(G)} 2(|f| - 3) \\ &= \sum_{v \in V(G)} \deg(v) - 6|V(G)| + 2 \sum_{f \in \mathcal{F}(G)} |f| - 6|\mathcal{F}(G)| \\ &= 2|E(G)| - 6|V(G)| + 2(2|E(G)|) - 6|\mathcal{F}(G)| && \text{by handshaking} \\ &= 6(|E(G)| - |V(G)| - |\mathcal{F}(G)|) \\ &= -6(1 + |\mathcal{C}(G)|) && \text{by Euler's formula} \\ &= -12 && \text{if connected} \end{aligned}$$

#### 1. Vertex-centric Setup

- $\text{ch}_0(v) = \deg(v) - 6 \ \forall v \in V(G)$
- $\text{ch}_0(f) = 2(|f| - 3) \ \forall f \in \mathcal{F}(G)$
- Sum:  $-12$  if connected
- Good for: unrestricted plane graphs.

#### 2. Face-centric Setup

- $\text{ch}_0(v) = 2(\deg(v) - 3) \ \forall v \in V(G)$
- $\text{ch}_0(f) = |f| - 6 \ \forall f \in \mathcal{F}(G)$
- Sum:  $-12$  if connected
- Good for: cubic plane graphs

#### 3. Balanced Setup

- $\text{ch}_0(v) = \deg(v) - 4 \ \forall v \in V(G)$
- $\text{ch}_0(f) = |f| - 4 \ \forall f \in \mathcal{F}(G)$
- Sum:  $-8$  if connected
- Good for: triangle-free plane graphs or restrictions on triangles

### A First Example

Some notation:

- $k$ -vertex: degree is  $k$
- $k^+$ -vertex: degree is  $\geq k$
- $k^-$ -vertex: degree is  $\leq k$
- $k$ -face,  $k^+$ -face,  $k^-$ -face similar

**Proposition 1.17**

Every plane graph with  $\delta(G) \geq 5$  contains a 5-vertex adjacent to a  $7^-$ -vertex.

Vertex-centric Setup:  $\text{ch}_0(v) = \deg(v) - 6$ ,  $\text{ch}_0(f) = 2(|f| - 3)$ , Sum:  $\leq -12$ .

**Rule:** Every  $8^+$ -vertex sends  $+\frac{1}{4}$  charge to each neighbor.

Let  $\text{ch}$  denote the final charge after applying rule.

**Claim** All final charges are nonnegative.

- Faces:  $\text{ch}(f) = \text{ch}_0(f) \geq 0$

- $8^+$ -vertices:

$$\text{ch}(v) \geq \deg(v) - 6 - \frac{\deg(v)}{4} = \frac{3\deg(v) - 24}{4} \geq 0$$

since  $\deg(v) \geq 8$ .

- 6-vertex, 7-vertex:  $\text{ch}(v) \geq \text{ch}_0(v) \geq 0$

- 5-vertex:

$$\text{ch}(v) = -1 + \frac{1}{4}(5) \geq \frac{1}{4} > 0$$

since every neighbor of  $v$  is an  $8^+$  vertex and sends  $+\frac{1}{4}$  to  $v$  by Rule.

**A Second Example****Proposition 1.18**

Every plane graph with  $\delta(G) \geq 3$  contains:

- a 3-vertex incident with a  $5^-$ -face, or
- a  $5^-$ -vertex incident with a 3-face.

Balanced Setup:  $\text{ch}_0(v) = \deg(v) - 4$ ,  $\text{ch}_0(f) = |f| - 4$ , Sum:  $\leq -8$ .

**Rules:**

1. Every  $6^+$ -face sends  $+\frac{1}{3}$  charge to each incident vertex.
2. Every  $6^+$ -vertex sends  $+\frac{1}{3}$  charge to each incident face.

Let  $\text{ch}$  denote the final charge after applying both rules.

**Claim** All final charges are nonnegative.

- $6^+$ -vertices:

$$\text{ch}(v) \geq \deg(v) - 4 - \frac{\deg(v)}{3} = \frac{2\deg(v) - 12}{3} \geq 0$$

since  $\deg(v) \geq 6$ .

- 4-vertex, 5-vertex:  $\text{ch}(v) \geq \text{ch}_0(v) \geq 0$ .

- 3-vertex:

$$\text{ch}(v) = -1 + \frac{1}{3}(3) = 0$$

since every face incident with  $v$  is a  $6^+$ -face and sends  $+\frac{1}{3}$  to  $v$  by Rule 1.

- Symmetric for faces.

## A Final Example

### Proposition 1.19

Every plane graph with  $\delta(G) \geq 3$  contains:

- two adjacent 3-faces, or
- a  $j$ -face for some  $4 \leq j \leq 9$ , or
- a 10-face incident with only 3-vertices.

Here we use Face-centric Setup:  $\text{ch}_0(v) = 2(\deg(v) - 3)$ ,  $\text{ch}_0(f) = |f| - 6$ .

**Rules:**

1. Every  $10^+$ -face sends +1 to each adjacent 3-face.
2. Every  $4^+$ -vertex  $v$  sends +1 to each incident  $10^+$ -face  $f$  where  $v$  is contained in a triangle sharing an edge with  $f$ .

Let  $\text{ch}$  denote the final charge after applying both rules.

**Claim** All final charges are nonnegative.

- 3-vertices:  $\text{ch}(v) = \text{ch}_0(v) = 0$

- $4^+$ -vertex:

$$\text{ch}(v) \geq 2(\deg(v) - 3) - \left\lfloor \frac{2\deg(v)}{3} \right\rfloor = \left\lceil \frac{4\deg(v)}{3} \right\rceil - 6 \geq 0$$

since  $\deg(v) \geq 4$ .

- 3-face:

$$\text{ch}(f) = -3 + (+1)(3) = 0$$

since every face adjacent to  $f$  is a  $10^+$ -face which sends +1 to  $f$  by Rule 1.

- $10^+$ -faces:

– Loses 1 for every path along its boundary such that neighboring faces are triangles and ends are degree 3: Loses 1 to each triangle on the path by Rule 1, and Gains 1 for each interior vertex of path by Rule 2 (since these are  $4^+$ -vertices)

– Net loss is at most  $\lfloor \frac{|f|}{2} \rfloor$

- $11^+$ -face:

$$\text{ch}(f) \geq |f| - 6 - \left\lfloor \frac{|f|}{2} \right\rfloor = \left\lceil \frac{|f|}{2} \right\rceil - 6 \geq 0$$

- 10-face: if  $\text{ch}(f) < 0$ , then 5 such paths, implies  $f$  incident to only 3-vertices, contradiction.

### Corollary 1.20

Every planar graph with no 4 to 9-cycles is 3-colorable.

**Proof:**

By standard reduction, we may assume  $G$  is 2-connected and  $\delta(G) \geq 3$ . By proposition,  $\exists$  a 10-face  $C$  incident with only 3-vertices. Delete  $V(C)$ , color by induction, extend to  $C$ . (Works since even cycles are 2-list-colorable)  $\square$

# Index

---

## C

chromatic index ..... 12  
chromatic number ..... 4  
class 1 ..... 14  
class 2 ..... 14  
coloring ..... 4

## E

edge chromatic number ..... 12  
edge coloring ..... 12

## G

girth ..... 10

## K

k edge coloring ..... 12  
k-coloring ..... 4  
k-degenerate ..... 21  
Kempe chain ..... 13  
kernel ..... 17  
kernel perfect ..... 17  
Kierstead Path ..... 20

## L

latin square ..... 17  
line graph ..... 12  
list chromatic index ..... 17  
list chromatic number ..... 16

## M

missing colors ..... 14  
multiplicity ..... 19

## R

recoloring ..... 13

## S

switching ..... 13

## T

Tashkinov Tree ..... 20

## V

Vizing fan ..... 14