



Operating Characteristics Across Eight Scenarios (ncycle = 3, Carryover = 0)


A. Percentage of correct MTD selection




B. Percentage of overdosing selection




C. Percentage of administrations allocated to the MTD




D. Percentage of administrations overdosed




E. Percentage of administrations underdosed




F. Average total number of IPDE doses



G. Average sample size



H. Average trial duration




—▲— AIDE-alpha-CRM - - -◆- - - Oracle CRM

Scenario 8 note: all doses are above target; early stop is treated as correct MTD selection, and all non-stop selections are treated as overdosing.


Scenario 1 note: all doses are below target; dose 8 is treated as the true MTD. Patient percentages use dose-level administration counts divided by total administrations. Oracle duration is not plotted; oracle sample size is sum of dose-level treated counts.

Operating Characteristics Across Eight Scenarios (ncycle = 3, Carryover = 0.3)


A. Percentage of correct MTD selection




B. Percentage of overdosing selection




C. Percentage of administrations allocated to the MTD




D. Percentage of administrations overdosed




E. Percentage of administrations underdosed




F. Average total number of IPDE doses



G. Average sample size



H. Average trial duration




—▲— AIDE-alpha-CRM -◆- Oracle CRM

Scenario 8 note: all doses are above target; early stop is treated as correct MTD selection, and all non-stop selections are treated as overdosing.


Scenario 1 note: all doses are below target; dose 8 is treated as the true MTD. Patient percentages use dose-level administration counts divided by total administrations. Oracle duration is not plotted; oracle sample size is sum of dose-level treated counts.

Operating Characteristics Across Eight Scenarios (ncycle = 3, Carryover = 0.6)


A. Percentage of correct MTD selection




B. Percentage of overdosing selection




C. Percentage of administrations allocated to the MTD




D. Percentage of administrations overdosed




E. Percentage of administrations underdosed




F. Average total number of IPDE doses



G. Average sample size



H. Average trial duration




—▲— AIDE-alpha-CRM -◆- Oracle CRM

Scenario 8 note: all doses are above target; early stop is treated as correct MTD selection, and all non-stop selections are treated as overdosing.


Scenario 1 note: all doses are below target; dose 8 is treated as the true MTD. Patient percentages use dose-level administration counts divided by total administrations. Oracle duration is not plotted; oracle sample size is sum of dose-level treated counts.

Operating Characteristics Across Eight Scenarios (ncycle = 3, Carryover = 0.9)


A. Percentage of correct MTD selection




B. Percentage of overdosing selection




C. Percentage of administrations allocated to the MTD




D. Percentage of administrations overdosed




E. Percentage of administrations underdosed




F. Average total number of IPDE doses



G. Average sample size



H. Average trial duration



—▲— AIDE-alpha-CRM -◆- Oracle CRM

Scenario 8 note: all doses are above target; early stop is treated as correct MTD selection, and all non-stop selections are treated as overdosing.

Scenario 1 note: all doses are below target; dose 8 is treated as the true MTD. Patient percentages use dose-level administration counts divided by total administrations. Oracle duration is not plotted; oracle sample size is sum of dose-level treated counts.