

















Operating Characteristics Across Six Scenarios (ncycle = 2, Carryover = 0)



—●— AIDE-CRM fixed, $r = 0$
 —■— AIDE-IP-CRM
 —▼— AIDE-CRM level
—▲— AIDE-alpha-CRM
 —◆— AIDE-CRM random
 - - -◆- - - Oracle CRM

Scenario 6 note: early stop is treated as correct MTD selection; all non-stop selections are treated as overdosing. Patient-overdose percentages use dose-level treated counts above the true MTD divided by total treated counts. Oracle duration is not plotted; oracle sample size is sum of dose-level number treated.









Operating Characteristics Across Six Scenarios (ncycle = 2, Carryover = 0.3)



● AIDE-CRM fixed, $r = 0$
■ AIDE-IP-CRM
 ▼ AIDE-CRM level
 ▲ AIDE-alpha-CRM
 ◆ AIDE-CRM random
 ◆ Oracle CRM

Scenario 6 note: early stop is treated as correct MTD selection; all non-stop selections are treated as overdosing. Patient-overdose percentages use dose-level treated counts above the true MTD divided by total treated counts. Oracle duration is not plotted; oracle sample size is sum of dose-level number treated.









Operating Characteristics Across Six Scenarios (ncycle = 2, Carryover = 0.6)



—●— AIDE-CRM fixed, $r = 0$
—■— AIDE-IP-CRM
 —▼— AIDE-CRM level
 —▲— AIDE-alpha-CRM
 —◆— AIDE-CRM random
 - - * - - Oracle CRM

Scenario 6 note: early stop is treated as correct MTD selection; all non-stop selections are treated as overdosing. Patient-overdose percentages use dose-level treated counts above the true MTD divided by total treated counts. Oracle duration is not plotted; oracle sample size is sum of dose-level number treated.

Operating Characteristics Across Six Scenarios (ncycle = 2, Carryover = 0.9)



—●— AIDE-CRM fixed, $r = 0$
—■— AIDE-IP-CRM
 —▼— AIDE-CRM level
 —▲— AIDE-alpha-CRM
 —◆— AIDE-CRM random
 - - * - - Oracle CRM

Scenario 6 note: early stop is treated as correct MTD selection; all non-stop selections are treated as overdosing. Patient-overdose percentages use dose-level treated counts above the true MTD divided by total treated counts. Oracle duration is not plotted; oracle sample size is sum of dose-level number treated.