



Operating Characteristics Across Eight Scenarios (ncycle = 3, Carryover = 0)


A. Percentage of correct MTD selection




B. Percentage of overdosing selection




C. Percentage of patients allocated to the MTD




D. Percentage of patients overdosed




E. Percentage of patients underdosed




F. Average total number of IPDE



G. Average sample size



H. Average trial duration




Scenario 8 note: early stop is treated as correct MTD selection; all non-stop selections are treated as overdosing.
Patient-overdose percentages use dose-level treated counts above the true MTD divided by total treated counts.
Oracle duration is not plotted; oracle sample size is sum of dose-level number treated.


—●— AIDE-CRM fixed, $r = 0$ —■— AIDE-alpha-CRM —▲— AIDE-cumu-CRM —◆— AIDE-CRM random -+ - Oracle CRM

Operating Characteristics Across Eight Scenarios (ncycle = 3, Carryover = 0.3)


A. Percentage of correct MTD selection




B. Percentage of overdosing selection




C. Percentage of patients allocated to the MTD




D. Percentage of patients overdosed




E. Percentage of patients underdosed




F. Average total number of IPDE



G. Average sample size



H. Average trial duration




Scenario 8 note: early stop is treated as correct MTD selection; all non-stop selections are treated as overdosing.
Patient-overdose percentages use dose-level treated counts above the true MTD divided by total treated counts.
Oracle duration is not plotted; oracle sample size is sum of dose-level number treated.


—●— AIDE-CRM fixed, $r = 0$ —■— AIDE-alpha-CRM —▲— AIDE-cumu-CRM —◆— AIDE-CRM random -+ - Oracle CRM

Operating Characteristics Across Eight Scenarios (ncycle = 3, Carryover = 0.6)


A. Percentage of correct MTD selection




B. Percentage of overdosing selection




C. Percentage of patients allocated to the MTD




D. Percentage of patients overdosed




E. Percentage of patients underdosed




F. Average total number of IPDE



G. Average sample size



H. Average trial duration




Scenario 8 note: early stop is treated as correct MTD selection; all non-stop selections are treated as overdosing.
 Patient-overdose percentages use dose-level treated counts above the true MTD divided by total treated counts.
 Oracle duration is not plotted; oracle sample size is sum of dose-level number treated.




Operating Characteristics Across Eight Scenarios (ncycle = 3, Carryover = 0.9)


A. Percentage of correct MTD selection




B. Percentage of overdosing selection




C. Percentage of patients allocated to the MTD




D. Percentage of patients overdosed




E. Percentage of patients underdosed




F. Average total number of IPDE



G. Average sample size



H. Average trial duration



Scenario 8 note: early stop is treated as correct MTD selection; all non-stop selections are treated as overdosing.
Patient-overdose percentages use dose-level treated counts above the true MTD divided by total treated counts.
Oracle duration is not plotted; oracle sample size is sum of dose-level number treated.

—●— AIDE-CRM fixed, $r = 0$ —■— AIDE-alpha-CRM —▲— AIDE-cumu-CRM —◆— AIDE-CRM random -+ - Oracle CRM