

# Phase I/II Operating Characteristics

TITE versus new non-TITE design - N = 30, Cycle\_Max = 2, efficacy futility cutoff = 0.85

|                                     | Dose 1      | Dose 2      | Dose 3      | Dose 4      | Dose 5     | No OBD %    |
|-------------------------------------|-------------|-------------|-------------|-------------|------------|-------------|
| <b>Scenario 1</b>                   |             |             |             |             |            |             |
| DLT rate                            | 0.3         | 0.35        | 0.4         | 0.45        | 0.5        |             |
| Efficacy rate                       | 0.05        | 0.06        | 0.07        | 0.08        | 0.09       |             |
| <b>OBD selection (%)</b>            |             |             |             |             |            |             |
| TITE design                         | 18.6        | 8.8         | 0.9         | 0.1         | 0          | 71.6        |
| <b>New non-TITE design</b>          | <b>17.4</b> | <b>7.2</b>  | <b>2</b>    | <b>0.8</b>  | <b>0.1</b> | <b>72.5</b> |
| <b>Mean administrations by dose</b> |             |             |             |             |            |             |
| TITE design                         | 11.5        | 4.6         | 2.2         | 0.9         | 0.3        |             |
| <b>New non-TITE design</b>          | <b>11</b>   | <b>4.4</b>  | <b>2.3</b>  | <b>1</b>    | <b>0.3</b> |             |
| <b>Scenario 16</b>                  |             |             |             |             |            |             |
| DLT rate                            | 0.02        | 0.06        | 0.09        | 0.14        | 0.26       |             |
| Efficacy rate                       | 0.2         | 0.2         | 0.2         | 0.2         | 0.2        |             |
| <b>OBD selection (%)</b>            |             |             |             |             |            |             |
| TITE design                         | 31.4        | 24.3        | 19.3        | 15.1        | 9.8        | 0.1         |
| <b>New non-TITE design</b>          | <b>33.9</b> | <b>22.2</b> | <b>16.5</b> | <b>15.4</b> | <b>12</b>  | <b>0</b>    |
| <b>Mean administrations by dose</b> |             |             |             |             |            |             |
| TITE design                         | 7.3         | 7           | 5.9         | 5.5         | 4.3        |             |
| <b>New non-TITE design</b>          | <b>6.6</b>  | <b>6.1</b>  | <b>5.6</b>  | <b>6.1</b>  | <b>5.6</b> |             |

# Phase I/II Operating Characteristics

TITE versus new non-TITE design - N = 30, Cycle\_Max = 2, efficacy futility cutoff = 0.85

|                                     | Dose 1      | Dose 2      | Dose 3      | Dose 4      | Dose 5      | No OBD % |
|-------------------------------------|-------------|-------------|-------------|-------------|-------------|----------|
| <b>Scenario 20</b>                  |             |             |             |             |             |          |
| DLT rate                            | 0.07        | 0.14        | 0.31        | 0.41        | 0.52        |          |
| Efficacy rate                       | 0.05        | 0.25        | 0.25        | 0.25        | 0.25        |          |
| <b>OBD selection (%)</b>            |             |             |             |             |             |          |
| TITE design                         | 11          | 50.2        | 25.9        | 8.8         | 1.2         | 2.9      |
| <b>New non-TITE design</b>          | <b>10.4</b> | <b>47.7</b> | <b>25.6</b> | <b>10.6</b> | <b>2.7</b>  | <b>3</b> |
| <b>Mean administrations by dose</b> |             |             |             |             |             |          |
| TITE design                         | 5.4         | 11.1        | 8           | 3.8         | 1.4         |          |
| <b>New non-TITE design</b>          | <b>5.7</b>  | <b>9.5</b>  | <b>8</b>    | <b>4.6</b>  | <b>1.8</b>  |          |
| <b>Scenario 24</b>                  |             |             |             |             |             |          |
| DLT rate                            | 0.01        | 0.02        | 0.05        | 0.1         | 0.25        |          |
| Efficacy rate                       | 0.05        | 0.08        | 0.2         | 0.5         | 0.5         |          |
| <b>OBD selection (%)</b>            |             |             |             |             |             |          |
| TITE design                         | 1.3         | 1.8         | 4.8         | 56          | 36.1        | 0        |
| <b>New non-TITE design</b>          | <b>0.4</b>  | <b>0.8</b>  | <b>3.6</b>  | <b>56.6</b> | <b>38.6</b> | <b>0</b> |
| <b>Mean administrations by dose</b> |             |             |             |             |             |          |
| TITE design                         | 3.4         | 3.5         | 4.5         | 10.6        | 8           |          |
| <b>New non-TITE design</b>          | <b>3.6</b>  | <b>3.7</b>  | <b>4.9</b>  | <b>9.4</b>  | <b>8.4</b>  |          |

# Phase I/II Operating Characteristics

TITE versus new non-TITE design - N = 30, Cycle\_Max = 2, efficacy futility cutoff = 0.85

|                                     | Dose 1      | Dose 2      | Dose 3      | Dose 4      | Dose 5      | No OBD %   |
|-------------------------------------|-------------|-------------|-------------|-------------|-------------|------------|
| <b>Scenario 27</b>                  |             |             |             |             |             |            |
| DLT rate                            | 0.11        | 0.17        | 0.3         | 0.5         | 0.62        |            |
| Efficacy rate                       | 0.1         | 0.2         | 0.4         | 0.42        | 0.45        |            |
| <b>OBD selection (%)</b>            |             |             |             |             |             |            |
| TITE design                         | 16.4        | 27.8        | 42.4        | 6.8         | 0.5         | 6.1        |
| <b>New non-TITE design</b>          | <b>10.5</b> | <b>23.6</b> | <b>46.6</b> | <b>12.5</b> | <b>1.1</b>  | <b>5.7</b> |
| <b>Mean administrations by dose</b> |             |             |             |             |             |            |
| TITE design                         | 7.7         | 8.2         | 9           | 3.3         | 0.9         |            |
| <b>New non-TITE design</b>          | <b>7.4</b>  | <b>8.2</b>  | <b>8.4</b>  | <b>4</b>    | <b>1.1</b>  |            |
| <b>Scenario 38</b>                  |             |             |             |             |             |            |
| DLT rate                            | 0.05        | 0.1         | 0.12        | 0.15        | 0.3         |            |
| Efficacy rate                       | 0.05        | 0.08        | 0.12        | 0.15        | 0.5         |            |
| <b>OBD selection (%)</b>            |             |             |             |             |             |            |
| TITE design                         | 17.2        | 10          | 8.9         | 8.9         | 53.5        | 1.5        |
| <b>New non-TITE design</b>          | <b>11.7</b> | <b>6</b>    | <b>5.5</b>  | <b>8.5</b>  | <b>67.3</b> | <b>1</b>   |
| <b>Mean administrations by dose</b> |             |             |             |             |             |            |
| TITE design                         | 4.9         | 5           | 5.5         | 5.4         | 9           |            |
| <b>New non-TITE design</b>          | <b>4.7</b>  | <b>4.8</b>  | <b>5.3</b>  | <b>6.9</b>  | <b>8.2</b>  |            |