

Reproducible Research Rpubs Assignment

Shaaz

March 20, 2016

Link on [RPubs](#)

My [Repository](#) On Github

Impact of Severe Weather Events on Public Health and Economy in the United States

Synonpsis

In this report, we aim to analyze the impact of different weather events on public health and economy based on the storm database collected from the U.S. National Oceanic and Atmospheric Administration's (NOAA) from 1950 - 2011. We will use the estimates of fatalities, injuries, property and crop damage to decide which types of event are most harmful to the population health and economy. From these data, we found that excessive heat and tornado are most harmful with respect to population health, while flood, drought, and hurricane/typhoon have the greatest economic consequences.

Basic settings

```
echo = TRUE # Always make code visible
options(scipen = 1) # Turn off scientific notations for numbers
library(R.utils)
```

```
## Loading required package: R.oo
```

```
## Loading required package: R.methodsS3
```

```
## R.methodsS3 v1.7.1 (2016-02-15) successfully loaded. See ?R.methodsS3 for help.
```

```
## R.oo v1.20.0 (2016-02-17) successfully loaded. See ?R.oo for help.
```

```
##  
## Attaching package: 'R.oo'
```

```
## The following objects are masked from 'package:methods':  
##  
##      getClasses, getMethods
```

```
## The following objects are masked from 'package:base':  
##  
##      attach, detach, gc, load, save
```

```
## R.utils v2.2.0 (2015-12-09) successfully loaded. See ?R.utils for help.
```

```
##  
## Attaching package: 'R.utils'
```

```
## The following object is masked from 'package:utils':
```

```
##
##      timestamp
```

```
## The following objects are masked from 'package:base':
##
##      cat, commandArgs, getOption, inherits, isOpen, parse, warnings
```

```
library(ggplot2)
library(plyr)
require(gridExtra)
```

```
## Loading required package: gridExtra
```

Data Processing

First, we download the data file and unzip it.

```
if (!"repdata_data_StormData.csv.bz2" %in% dir("E:/Downloads/Compressed/")) {
  print("hhh")
  download.file("http://d396qusza40orc.cloudfront.net/repdata%2Fdata%2FStormData.csv.bz2", de
stfile = "E:/Downloads/Compressed/stormData.csv.bz2")
  bunzip2("E:/Downloads/Compressed/stormData.csv.bz2", overwrite=T, remove=F)
}
```

Then, we read the generated csv file. If the data already exists in the working environment, we do not need to load it again. Otherwise, we read the csv file.

```
if (!"stormData" %in% ls()) {
```

```
stormData <- read.csv("E:/Documents/stormData.csv", sep = ",")
}
dim(stormData)
```

```
## [1] 902297      37
```

```
head(stormData, n = 2)
```


```
##      STATE__      BGN_DATE BGN_TIME TIME_ZONE COUNTY COUNTYNAM STATE
## 1         1 4/18/1950 0:00:00    0130     CST     97    MOBILE    AL
## 2         1 4/18/1950 0:00:00    0145     CST      3    BALDWIN    AL
##      EVTYPE BGN_RANGE BGN_AZI BGN_LOCATI END_DATE END_TIME COUNTY_END
## 1 TORNADO          0                0
## 2 TORNADO          0                0
##      COUNTYENDN END_RANGE END_AZI END_LOCATI LENGTH WIDTH F MAG FATALITIES
## 1          NA          0          14    100 3   0          0
## 2          NA          0          2    150 2   0          0
##      INJURIES PROPDMG PROPDMGEXP CROPDMG CROPDMGEXP WFO STATEOFFIC ZONENAMES
## 1         15    25.0           K         0
## 2          0     2.5           K         0
##      LATITUDE LONGITUDE LATITUDE_E LONGITUDE_ REMARKS REFNUM
## 1      3040      8812      3051      8806          1
## 2      3042      8755          0          0          2
```

There are 902297 rows and 37 columns in total. The events in the database start in the year 1950 and end in November 2011. In the earlier years of the database there are generally fewer events recorded, most likely due to a lack of good records. More recent years should be considered more complete.

```
if (dim(stormData)[2] == 37) {
```

```
stormData$year <- as.numeric(format(as.Date(stormData$BGN_DATE, format = "%m/%d/%Y %H:%M:%S"), "%Y"))
}
hist(stormData$year, breaks = 30)
```

Histogram of stormData\$year



Based on the above histogram, we see that the number of events tracked starts to significantly increase around 1995. So, we use the subset of the data from 1990 to 2011 to get most out of good records.

```
storm <- stormData[stormData$year >= 1995, ]
```

```
dim(storm)
```

```
## [1] 681500    38
```

Now, there are 681500 rows and 38 columns in total.

Impact on Public Health

In this section, we check the number of **fatalities** and **injuries** that are caused by the severe weather events. We would like to get the first 15 most severe types of weather events.

```
sortHelper <- function(fieldName, top = 15, dataset = stormData) {  
  index <- which(colnames(dataset) == fieldName)  
  field <- aggregate(dataset[, index], by = list(dataset$EVTYPE), FUN = "sum")  
  names(field) <- c("EVTYPE", fieldName)  
  field <- arrange(field, field[, 2], decreasing = T)  
  field <- head(field, n = top)  
  field <- within(field, EVTYPE <- factor(x = EVTYPE, levels = field$EVTYPE))  
  return(field)  
}  
  
fatalities <- sortHelper("FATALITIES", dataset = storm)  
injuries <- sortHelper("INJURIES", dataset = storm)
```

Impact on Economy

We will convert the **property damage** and **crop damage** data into comparable numerical forms according to the meaning of units described in the code book ([Storm Events](#)). Both `PROPDMGEXP` and `CROPDMGEXP` columns record a multiplier for each observation where we have Hundred (H), Thousand (K), Million (M) and Billion (B).

```
convertHelper <- function(dataset = storm, fieldName, newFieldName) {
```

```

totalLen <- dim(dataset)[2]
index <- which(colnames(dataset) == fieldName)
dataset[, index] <- as.character(dataset[, index])
logic <- !is.na(toupper(dataset[, index]))
dataset[logic & toupper(dataset[, index]) == "B", index] <- "9"
dataset[logic & toupper(dataset[, index]) == "M", index] <- "6"
dataset[logic & toupper(dataset[, index]) == "K", index] <- "3"
dataset[logic & toupper(dataset[, index]) == "H", index] <- "2"
dataset[logic & toupper(dataset[, index]) == "", index] <- "0"
dataset[, index] <- as.numeric(dataset[, index])
dataset[is.na(dataset[, index]), index] <- 0
dataset <- cbind(dataset, dataset[, index - 1] * 10^dataset[, index])
names(dataset)[totalLen + 1] <- newFieldName
return(dataset)
}

storm <- convertHelper(storm, "PROPDMGEXP", "propertyDamage")

```

```

## Warning in convertHelper(storm, "PROPDMGEXP", "propertyDamage"): NAs
## introduced by coercion

```

```

storm <- convertHelper(storm, "CROPDMGEXP", "cropDamage")

```

```

## Warning in convertHelper(storm, "CROPDMGEXP", "cropDamage"): NAs introduced
## by coercion

```

```

names(storm)

```



```
## [1] "STATE__"      "BGN_DATE"      "BGN_TIME"      "TIME_ZONE"
## [5] "COUNTY"      "COUNTYNAME"   "STATE"          "EVTYPE"
## [9] "BGN_RANGE"    "BGN_AZI"       "BGN_LOCATI"     "END_DATE"
## [13] "END_TIME"     "COUNTY_END"   "COUNTYENDN"    "END_RANGE"
## [17] "END_AZI"      "END_LOCATI"    "LENGTH"         "WIDTH"
## [21] "F"           "MAG"           "FATALITIES"     "INJURIES"
## [25] "PROPDMG"      "PROPDMGEXP"    "CROPDMG"         "CROPDMGEXP"
## [29] "WFO"          "STATEOFFIC"    "ZONENAMES"       "LATITUDE"
## [33] "LONGITUDE"    "LATITUDE_E"    "LONGITUDE_"      "REMARKS"
## [37] "REFNUM"       "year"          "propertyDamage"  "cropDamage"
```

```
options(scipen=999)
property <- sortHelper("propertyDamage", dataset = storm)
crop <- sortHelper("cropDamage", dataset = storm)
```

Results

As for the impact on public health, we have got two sorted lists of severe weather events below by the number of people badly affected.

fatalities

```
##           EVTYPE FATALITIES
## 1  EXCESSIVE HEAT      1903
## 2      TORNADO      1545
## 3  FLASH FLOOD       934
## 4        HEAT       924
## 5  LIGHTNING       729
## 6      FLOOD       423
```

## 7	RIP CURRENT	360
## 8	HIGH WIND	241
## 9	TSTM WIND	241
## 10	AVALANCHE	223
## 11	RIP CURRENTS	204
## 12	WINTER STORM	195
## 13	HEAT WAVE	161
## 14	THUNDERSTORM WIND	131
## 15	EXTREME COLD	126


injuries

##	EVTTYPE	INJURIES
## 1	TORNADO	21765
## 2	FLOOD	6769
## 3	EXCESSIVE HEAT	6525
## 4	LIGHTNING	4631
## 5	TSTM WIND	3630
## 6	HEAT	2030
## 7	FLASH FLOOD	1734
## 8	THUNDERSTORM WIND	1426
## 9	WINTER STORM	1298
## 10	HURRICANE/TYPHOON	1275
## 11	HIGH WIND	1093
## 12	HAIL	916
## 13	WILDFIRE	911
## 14	HEAVY SNOW	751
## 15	FOG	718


And the following is a pair of graphs of total fatalities and total injuries affected by these severe weather events.

```
fatalitiesPlot <- qplot(EVTYPE, data = fatalities, weight = FATALITIES, geom = "bar") +  
  scale_y_continuous("Number of Fatalities") +  
  theme(axis.text.x = element_text(angle = 45,  
    hjust = 1)) + xlab("Severe Weather Type") +  
  ggtitle("Total Fatalities by Severe Weather\n Events in the U.S.\n from 1995 - 2011")  
injuriesPlot <- qplot(EVTYPE, data = injuries, weight = INJURIES, geom = "bar") +  
  scale_y_continuous("Number of Injuries") +  
  theme(axis.text.x = element_text(angle = 45,  
    hjust = 1)) + xlab("Severe Weather Type") +  
  ggtitle("Total Injuries by Severe Weather\n Events in the U.S.\n from 1995 - 2011")  
grid.arrange(fatalitiesPlot, injuriesPlot, ncol = 2)
```

Total Fatalities by Severe Weather Events in the U.S. from 1995 - 2011



Total Injuries by Severe Weather Events in the U.S. from 1995 - 2011



Based on the above histograms, we find that **excessive heat** and **tornado** cause most fatalities; **tornado** causes most injuries in the United States from 1995 to 2011.

As for the impact on economy, we have got two sorted lists below by the amount of money cost by damages.

property

```
##          EVTYPE  propertyDamage
## 1          FLOOD    144022037057
## 2  HURRICANE/TYPHOON    69305840000
## 3          STORM SURGE    43193536000
## 4          TORNADO    24935939545
## 5          FLASH FLOOD    16047794571
## 6          HAIL    15048722103
## 7          HURRICANE    11812819010
## 8    TROPICAL STORM    7653335550
## 9          HIGH WIND    5259785375
## 10         WILDFIRE    4759064000
## 11  STORM SURGE/TIDE    4641188000
## 12         TSTM WIND    4482361440
## 13         ICE STORM    3643555810
## 14 THUNDERSTORM WIND    3399282992
## 15    HURRICANE OPAL    3172846000
```

crop

```
##          EVTYPE  cropDamage
## 1    DROUGHT 13922066000
## 2    FLOOD  5422810400
## 3    HURRICANE 2741410000
## 4    HAIL  2614127070
## 5  HURRICANE/TYPHOON 2607872800
## 6    FLASH FLOOD 1343915000
## 7    EXTREME COLD 1292473000
```

## 8	FROST/FREEZE	1094086000
## 9	HEAVY RAIN	728399800
## 10	TROPICAL STORM	677836000
## 11	HIGH WIND	633561300
## 12	TSTM WIND	553947350
## 13	EXCESSIVE HEAT	492402000
## 14	THUNDERSTORM WIND	414354000
## 15	HEAT	401411500

And the following is a pair of graphs of total property damage and total crop damage affected by these severe weather events.

```
propertyPlot <- qplot(EVTYPE, data = property, weight = propertyDamage, geom = "bar") +
  theme(axis.text.x = element_text(angle = 45, hjust = 1)) + scale_y_continuous("Property Dam
age in US dollars")+
  xlab("Severe Weather Type") + ggtitle("Total Property Damage by\n Severe Weather Events in\
n the U.S. from 1995 - 2011")

cropPlot<- qplot(EVTYPE, data = crop, weight = cropDamage, geom = "bar") +
  theme(axis.text.x = element_text(angle = 45, hjust = 1)) + scale_y_continuous("Crop Damage
in US dollars") +
  xlab("Severe Weather Type") + ggtitle("Total Crop Damage by \nSevere Weather Events in\n th
e U.S. from 1995 - 2011")
grid.arrange(propertyPlot, cropPlot, ncol = 2)
```

Property Damage in US dollars

Disaster Type	Property Damage (US dollars)
HURRICANE/TYPHOON	145,000,000,000
FLOOD	70,000,000,000
STORM SURGE	45,000,000,000
TORNADO	25,000,000,000
FLASH FLOOD	18,000,000,000
HAIL	17,000,000,000
HURRICANE	14,000,000,000
TROPICAL STORM	8,000,000,000
HIGH WIND	5,000,000,000
WILDFIRE	4,000,000,000
STORM SURGE/TIDE	4,000,000,000
TSTM WIND	4,000,000,000
ICE STORM	3,000,000,000
THUNDERSTORM WIND	3,000,000,000
HURRICANE OPAL	3,000,000,000

Property Damage in US dollars

A bar chart titled 'Crop Damage in US dollars' showing the impact of various weather events. The y-axis represents the damage in US dollars, with major grid lines at 0, 500,000,000, and 1,000,000,000. The x-axis lists 18 weather events. The bars are dark gray. The damage decreases significantly from left to right, with DROUGHT being the most damaging event by a large margin.

Weather Event	Crop Damage (US dollars)
DROUGHT	~1,200,000,000
FLOOD	~550,000,000
HURRICANE	~280,000,000
HURRICANE/TYPHOON	~270,000,000
HAIL	~270,000,000
FLASH FLOOD	~150,000,000
EXTREME COLD	~140,000,000
FROST/FREEZE	~120,000,000
HEAVY RAIN	~70,000,000
TROPICAL STORM	~70,000,000
HIGH WIND	~60,000,000
TSTM WIND	~50,000,000
EXCESSIVE HEAT	~40,000,000
THUNDERSTORM WIND	~30,000,000
HEAT	~30,000,000

Crop Damage in US dollars

version Are you a developer? Try out the [HTML to PDF API](#)

Conclusion

From these data, we found that **excessive heat** and **tornado** are most harmful with respect to population health, while **flood**, **drought**, and **hurricane/typhoon** have the greatest economic consequences.