

Advanced Mathematics Exercises

1 Chapter 1

Exercise 1.1. Let f be the scalar function defined by $f(x, y) = xy$. Sketch the contour lines and the vector field ∇f .

Solution: TODO □

Exercise 1.2. Let

$$f(x, y) = \sin(\pi xy) \cdot e^{-\frac{x}{3}} \quad \text{and} \quad p = \left(1, \frac{1}{3}\right).$$

- (a) Compute ∇f and $(\nabla f)(p)$. Use the special values of \sin and \cos .
- (b) Find the directions of maximum increase and decrease at p . You can give approximate values.
- (c) Give the direction of the contour line at p .
- (d) The equation of the tangent plane of the graph of f at (x_0, y_0) is

$$z = f(x_0, y_0) + (\nabla f)(x_0, y_0) \cdot (x - x_0, y - y_0).$$

Determine the equation of the tangent plane of $z = f(x, y)$ at p . Give a normal vector of the plane.

- (e) Find the directional derivative of $f(x, y)$ at p along the vector

$$v = \frac{1}{\sqrt{2}}(1, 1).$$

Solution: TODO □

Exercise 1.3. Compute curl and divergence of the vector field

$$F(x, y, z) = (\sin(x)xy, ze^{-x}, yz).$$

Solution: TODO □

Exercise 1.4. Let

$$f(x, y, z) = x^2yz^3 \quad \text{and} \quad F(x, y, z) = (xz, -y^2, 2x^2y).$$

Give ∇f , $\nabla^2 f$, $\nabla \cdot F$, and $\nabla \times F$.

Solution: TODO □

Exercise 1.5. Compute differential forms:

(a) Let

$$f(x, y, z) = \frac{x}{yz}$$

be a scalar function. Give the differential form df .

Hint:

$$df = F_1(x, y, z) dx + F_2(x, y, z) dy + F_3(x, y, z) dz.$$

(b) Let

$$\omega = x^2 \sin(y) dx + 2^x \cos(y) dy$$

be a differential form on \mathbb{R}^2 . Give the 2-form $d\omega$.

Hint:

$$d\omega = \phi(x, y) dx \wedge dy.$$

(c) Let

$$\omega = x^2 \sin(y) dx + z^2 \cos(y) dy - xy^2 dz$$

be a differential form on \mathbb{R}^3 . Give the 2-form $d\omega$.

Hint:

$$d\omega = F_1(x, y, z) dy \wedge dz + F_2(x, y, z) dz \wedge dx + F_3(x, y, z) dx \wedge dy.$$

Solution: TODO

□