PHYS 3040: MODERN PHYSICS

Department of Physics and Astronomy York University

Homework Set #3

Due on Wednesday November 6, 2024. Please write solutions clearly and carefully. Submit a single pdf file through eClass.

- 1. Fermilab puts a muon into a uniform magnetic field $\vec{B}=B\hat{z}$ to measure the muon's magnetic moment, which is a way to search for new physics beyond the standard model. We can name the energy eigenvalues E_+ and E_- . Suppose a muon is in the state $|+\rangle_n$ at time t=0, where $\hat{n}=\frac{1}{\sqrt{2}}(\hat{y}+\hat{z})$.
- (a) Calculate an expression for the state of the muon at any time t, named $|\psi(t)\rangle$.
- (b) Calculate the probability that a measurement of S_x will give $+\frac{\hbar}{2}$ at time t.
- (c) Use your final expression from (b) to compute numerical values for the minimum and maximum probabilities as time passes.
- (d) Calculate the expectation value of S_x at time t.
- (e) Use your final expression from (d) to obtain simple algebraic answers for the minimum and maximum expectation values as time passes.
- **2.** For a hospital MRI machine, the magnetic field has both a longitudinal component and a transverse component. The Hamiltonian matrix and its eigenvectors are

$$H = \frac{\hbar}{2} \left(\begin{array}{cc} \omega \cos \theta & \omega \sin \theta \\ \omega \sin \theta & -\omega \cos \theta \end{array} \right), \qquad |+\rangle_B = \left(\begin{array}{c} \cos \frac{\theta}{2} \\ \sin \frac{\theta}{2} \end{array} \right), \qquad |-\rangle_B = \left(\begin{array}{c} \sin \frac{\theta}{2} \\ -\cos \frac{\theta}{2} \end{array} \right).$$

We know that H depends on time during normal operation, but for this homework question we will operate the machine with H held constant. Suppose the nucleus of a hydrogen atom in the patient is in the state $|-\rangle_z$ at time t=0.

- (a) Calculate the eigenvalues of H.
- (b) Calculate an expression for the state of the nucleus at any time t, named $|\psi(t)\rangle$.
- (c) Calculate the expectation value of S_x for the nucleus at time t.
- (d) Suppose your MRI machine has $\theta=45^\circ$. Use your final expression from (c) to compute algebraic values for the minimum and maximum expectation values as time passes.