
OCR MEI Notes

(Year 2)

Department of Mathematics



Author: Siddartha Nath

Date: 21st December 2023

Disclaimer

First Edition

Publisher: Published by PolyNath

Organisation Website: www.polynath.com

Organisation Email: polynathofficial@gmail.com

This publication is not affiliated with, nor has it been authorised, sponsored, or otherwise approved by any academic or industry organisation. The content within this book is meant for guidance only, and the publishers and authors offer no warranty regarding the use of this material for individual purposes. The authors and publishers disclaim any responsibility for any potential loss or damage arising from the use of this book. No part of this guidebook may be replicated, altered, or transmitted in any form (via electronic, photocopying, recording) or by any information retrieval system without the prior written permission of the publisher. While all efforts have been made to ensure the accuracy of the information in this book, the authors and publisher are not responsible for any errors or omissions or for the results obtained from the use of this information. Your statutory rights remain unaffected.

Author

Siddhartha Nath is the founder of *PolyNath* and the author of a plethora of top-selling books (*Quantitative Finance*, *MAT*, *NLP*, *TMUA/CTMUA*).

He has an undergraduate First-Class Honours BSc Mathematics with Statistics from Imperial College London (ICL) and a postgraduate Distinction MSc Computational Statistics and Machine Learning from University College London (UCL). He has taken his education and amassed over 2 years of industry experience, having worked as a Data Scientist for PayPal and DataSpartan.

In terms of academic teaching, he is an extremely dedicated and supportive tutor, having taught for over 5 years at public and private schools, for GCSE, A-Levels and University preparation. He possesses a wealth of knowledge in all UK-based Mathematics and Computer Science admission tests, with the following accomplishments:

- Achieved a perfect score of 9.0/9.0, in the Cambridge TMUA, resulting in a top 10% ranking.
- Achieved a commendable score of 63/100, in the Oxford MAT, resulting in a top 25% ranking.
- Achieved a high score of 1, 1, 1 in the Cambridge STEP I, II and III.

Outside of academia, he enjoys music, dance, watching sports and creating content.



Preface

This document includes a summary of notes of Chapter 1 - Proof (OCR MEI Year 2).

Happy Learning!

Chapter 1

Proof

Summary Checklist

1.
 - \Rightarrow means *if...then, implies, therefore* etc...
 - \Leftarrow means *only if..., is implied by, follows from* etc...
 - \Leftrightarrow means *implies and is implied by, is equivalent to* etc...
2. The converse of $A \Rightarrow B$ is $B \Rightarrow A$.
3.
 - If $A \Leftarrow B$, A is a **necessary** condition for B .
 - If $A \Rightarrow B$, A is a **sufficient** condition for B .
4. The methods of proof are:
 - Direct Argument
 - Exhaustion
 - Contradiction
 - Counter-Example

Below are 2 examples, demonstrating a complete *thought-process* breakdown of how to approach a *proof*:

Example

Consider the following example

"Prove that $n^3 - n$ is a multiple of 6."

We do this via **direct argument**.

1. **Firstly, is this true?**

When $n = 1$, $1^3 - 1 = 0$, which is divisible by 6. For $n = 2$, $2^3 - 2 = 6$, which is divisible by 6. For $n = 3$, $3^3 - 3 = 24$, which is divisible by 6. So far, it seems to hold. Note, for shorthand, we will use $|$ to denote *divides* i.e. $2|6$ means 2 divides 6.

2. **Secondly, how do we prove divisibility?**

To show divisibility, the number should be divisible by all the *prime* factors of the number i.e. prime factorisation tree. Hence, to be divisible by 6, we should prove that the number is divisible by 2 and 3.

3. **Thirdly, how do we use algebra to demonstrate these divisibility properties?**

- Since we are thinking about factors, we perhaps should do some *factorisation*...

$$\begin{aligned}n^3 - n &= n(n^2 - 1) \\ &= n(n - 1)(n + 1)\end{aligned}$$

- Logically, $2|n^3 - n$ because we have a product 2 consecutive integers between the 3 factors and thus one of them must be divisible by 2.
- Logically, $3|n^3 - n$ because we have a product 3 consecutive integers between the 3 factors and thus one of them must be divisible by 3.

4. **Finally, how do we conclude the proof?**

Hence, since $2|n^3 - n$ and $3|n^3 - n$, then $2 \cdot 3 = 6|n^3 - n$, a trivial property of divisibility and factors.

A follow up question is

"Prove that $n^3 + 11n$ is a multiple of 6."

1. **Firstly, is this true?**

When $n = 1$, $1^3 + 11(1) = 12$, which is divisible by 6. For $n = 2$, $2^3 + 11(2) = 30$, which is divisible by 6. For $n = 3$, $3^3 + 11(3) = 42$, which is divisible by 6. So far, it

seems to hold.

2. **Secondly, how do we prove divisibility?**

We already know from before that we should show this is divisible by both 2 and 3.

3. **Thirdly, how do we use algebra to demonstrate these divisibility properties?**

- Since we are thinking about factors, we perhaps should do some *factorisation*...

$$n^3 + 11n = n(n^2 + 11)$$

PROBLEM:

We cannot do the same trick as before because we do not seem to have consecutive factors i.e. we do not have a nice factorisation. This is usually the point where we stop and give up... But no! **There was a reason for why we did the first part of the question!**

4. **Fourthly, how can I use previous knowledge to help?**

- A trivial property of divisibility is that if $k|A$ and $k|B$, then $k|A + B$ i.e. if you add multiples of k together, the resulting number is also divisible by k .
- One should spot that to get $n^3 + 11n$, we must simply add $12n$.

5. **Finally, how do we conclude the proof?**

As $6|12n$ because 6 is a factor of 12, regardless of n , using the rule before, we can conclude that since $6|n^3 - n$ and $6|12n$, we must have $6|(n^3 - n) + (12n) = n^3 + 11n$.

What happens if we need to prove divisibility from first principles, where factorisation or prior knowledge base is not available?

To achieve this, we follow a 2-step approach.

- Obtain the prime factors of the divisor.
- Proceed via cases with the prime factors via exhaustion.

Take a look back at $n^3 + 11n$ - focusing on divisibility by 2 and 3, we do a case breakdown by covering all possibilities of what n could be

1. **Divisibility by 2:**

- $n = 2k$: $n^3 + 11n = 8k^3 + 22k = 2k(4k^2 + 11) \rightarrow$ **divisible by 2.**
- $n = 2k+1$: $n^3 + 11n = 8k^3 + 12k^2 + 6k + 1 + 22k + 11 = 2(4k^3 + 6k^2 + 14k + 6) + 12 \rightarrow$ **divisible by 2.**

This says that for all n , broken into multiples/non-multiples of 2, $n^3 + 11n$ is always divisible by 2.

2. **Divisibility by 3:**

- $n = 3k$: $n^3 + 11n = 27k^3 + 33k = 3k(9k^2 + 11) \rightarrow$ **divisible by 3.**
- $n = 3k + 1$: $n^3 + 11n = 27k^3 + 27k^2 + 42k + 12 = 3(9k^3 + 9k^2 + 14k + 4) \rightarrow$ **divisible by 3.**
- $n = 3k + 2$: $n^3 + 11n = 27k^3 + 54k^2 + 69k + 30 = 3(9k^3 + 18k^2 + 33k + 10) \rightarrow$ **divisible by 3.**

This says that for all n , broken into multiples/non-multiples of 3, $n^3 + 11n$ is always divisible by 3.

3. Hence, since $2|n^3 + 11n$ and $3|n^3 + 11n$, then $2 \cdot 3 = 6|n^3 + 11n$, a trivial property of divisibility and factors.

Note: We do not need to test say $2k+2$ or $3k+3$ because they are equivalent constructions i.e. $2k + 2 = 2(k + 1) \equiv 2p$ and $3k + 3 = 3(k + 1) \equiv 3q$. This links to an equivalent derivation, known as *modulo arithmetic*, which is not covered in the syllabus.

Example

Consider the following example

"Prove that $\sqrt{3}$ is irrational."

We do this via **contradiction**.

1. **Firstly, is this true?**

We know when taking square roots of numbers which are not square numbers themselves, that we end up with a decimal value, which typically is never ending i.e. infinite and thus at first glance, this seems true.

2. **Secondly, how do we prove irrationality?**

We prove irrationality by looking at the negation of the statement i.e. **assuming** rationality. We then follow algebraic steps and gather information, to which if we end up concluding something which is **false**, we then know that we were **wrong** to assume rationality in the first place.

3. **Thirdly, how do we use algebra to demonstrate rationality?**

- Since we assume $\sqrt{3}$ is rational, this means it can be expressed as a fraction $\frac{a}{b}$, where a and b are positive integers with no common factors i.e. $\text{HCF}(a, b) = 1$.

- Given that $\sqrt{3} = \frac{a}{b}$, we can arrange to obtain

$$3 = \left(\frac{a}{b}\right)^2 = \frac{a^2}{b^2}$$

$$\therefore a^2 = 3b^2$$

The right hand side is divisible by 3 and hence so must the left i.e. $3|a^2$. This also means that $3|a$ i.e. $a = 3k$. This step should also be proven rigorously but the main reason for this is because 3 is prime and thus if $3|a^2$, the prime factorisation of a^2 must contain 3^2 , which implies $3|a$.

- Substituting this back in gives us

$$(3k)^2 = 3b^2$$

$$\Rightarrow 9k^2 = 3b^2$$

$$\therefore 3k^2 = b^2$$

The left hand side is divisible by 3 and hence so must the right i.e. $3|b^2$. This also means that $3|b$ i.e. $b = 3p$.

- And voila! We have arrived at a contradiction since the $\text{HCF}(a, b) = \text{HCF}(3k, 3p) = 3$ but, by construction of the statement, we should have $\text{HCF}(a, b) = 1$.

Note: It is not so clear on an alternative method to prove this. If we tried to do this via *direct argument*, we would need more knowledge about square root representations, but this is not covered in the syllabus.