

Please check the examination details below before entering your candidate information

Candidate surname					Other names				
Centre Number					Candidate Number				
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	

Pearson Edexcel Level 3 GCE

Paper reference	8MA0/22
-----------------	----------------

Mathematics

Advanced Subsidiary

PAPER 22: Mechanics

<p>You must have: Mathematical Formulae and Statistical Tables (Green), calculator</p>	<p>Total Marks</p>
---	--------------------

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear.
Answers without working may not gain full credit.
- Unless otherwise indicated, wherever a value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- The total mark for this part of the examination is 30. There are 4 questions.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P68658A

©2022 Pearson Education Ltd.

Q:1/1/1/



Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 1 is 7 marks)

2. A train travels along a straight horizontal track from station P to station Q .

In a model of the motion of the train, at time $t = 0$ the train starts from rest at P , and moves with constant acceleration until it reaches its maximum speed of 25 m s^{-1}

The train then travels at this constant speed of 25 m s^{-1} before finally moving with constant deceleration until it comes to rest at Q .

The time spent decelerating is four times the time spent accelerating.

The journey from P to Q takes 700 s.

Using the model,

- (a) sketch a speed-time graph for the motion of the train between the two stations P and Q .
(1)

The distance between the two stations is 15 km.

Using the model,

- (b) show that the time spent accelerating by the train is 40 s,
(3)

- (c) find the acceleration, in m s^{-2} , of the train,
(1)

- (d) find the speed of the train 572 s after leaving P .
(2)

- (e) State one limitation of the model which could affect your answers to parts (b) and (c).
(1)



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 2 is 8 marks)

3. A fixed point O lies on a straight line.

A particle P moves along the straight line.

At time t seconds, $t \geq 0$, the distance, s metres, of P from O is given by

$$s = \frac{1}{3}t^3 - \frac{5}{2}t^2 + 6t$$

- (a) Find the acceleration of P at each of the times when P is at instantaneous rest.

(6)

- (b) Find the total distance travelled by P in the interval $0 \leq t \leq 4$

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 8 6 5 8 A 0 1 1 1 6

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 3 is 9 marks)

4.

**Figure 1**

A vertical rope PQ has its end Q attached to the top of a small lift cage.

The lift cage has mass 40 kg and carries a block of mass 10 kg , as shown in Figure 1.

The lift cage is raised vertically by moving the end P of the rope vertically upwards with constant acceleration 0.2 m s^{-2}

The rope is modelled as being light and inextensible and air resistance is ignored.

Using the model,

(a) find the tension in the rope PQ (3)

(b) find the magnitude of the force exerted on the block by the lift cage. (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 8 6 5 8 A 0 1 5 1 6

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 4 is 6 marks)**TOTAL FOR MECHANICS IS 30 MARKS**



Mark Scheme (Results)

Summer 2022

Pearson Edexcel GCE
AS Mathematics (8MA0)
Paper 22 Mechanics

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2022

Question Paper Log Number P68658A*

Publications Code 8MA0_22_2206_MS*

All the material in this publication is copyright

© Pearson Education Ltd 2022

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 80.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \checkmark will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 5. Where a candidate has made multiple responses and indicates which response they wish to submit, examiners should mark this response.
If there are several attempts at a question which have not been crossed out, examiners should mark the final answer which is the answer that is the most complete.

6. Ignore wrong working or incorrect statements following a correct answer.
7. Mark schemes will firstly show the solution judged to be the most common response expected from candidates. Where appropriate, alternative answers are provided in the notes. If examiners are not sure if an answer is acceptable, they will check the mark scheme to see if an alternative answer is given for the method used.

General Principles for Mechanics Marking

(But note that specific mark schemes may sometimes override these general principles)

- Rules for M marks: correct no. of terms; dimensionally correct; all terms that need resolving (i.e. multiplied by cos or sin) are resolved.
 - Omission or extra g in a resolution is an accuracy error not method error.
 - Omission of mass from a resolution is a method error.
 - Omission of a length from a moments equation is a method error.
 - Omission of units or incorrect units is not (usually) counted as an accuracy error.
 - DM indicates a dependent method mark i.e. one that can only be awarded if a previous specified method mark has been awarded.
 - Any numerical answer which comes from use of $g = 9.8$ should be given to 2 or 3 SF.
 - Use of $g = 9.81$ should be penalised once per (complete) question.
- N.B. Over-accuracy or under-accuracy of correct answers should only be penalised *once* per complete question. However, premature approximation should be penalised every time it occurs.
- Marks must be entered in the same order as they appear on the mark scheme.
 - In all cases, if the candidate clearly labels their working under a particular part of a question i.e. (a) or (b) or (c),.....then that working can only score marks for that part of the question.
 - Accept column vectors in all cases.
 - Misreads – if a misread does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, bearing in mind that after a misread, the subsequent A marks affected are treated as A ft

• Mechanics Abbreviations

M(A)	Taking moments about A.
N2L	Newton's Second Law (Equation of Motion)
NEL	Newton's Experimental Law (Newton's Law of Impact)
HL	Hooke's Law
SHM	Simple harmonic motion
PCLM	Principle of conservation of linear momentum
RHS, LHS	Right hand side, left hand side.

Question	Scheme	Marks	AOs
1(a)	Complete method to produce an equation in U only	M1	3.4
	e.g. $10^2 = U^2 + 2 \times g \times 1.8$ oe	A1	1.1b
	OR a complete method where they find T first and use it to find an equation in U only	M1	
	A correct equation in U only.	A1	
	$U = 8$ (<u>only</u> this answer)	A1	1.1b
		(3)	
(b)	Complete method to find an equation in T only: $10 = -8 + gT$ or $1.8 = 10T - \frac{1}{2}gT^2$ or $1.8 = \frac{(-8+10)}{2}T$ or $1.8 = -8T + \frac{1}{2}gT^2$ OR a complete method if they split the time. In both cases, the M1 is only earned on the final line when they try to add the two times to give an equation in T . ALT 1: time up + time down e.g. $0 = 8 - gt_{UP}$ ($\Rightarrow t_{UP} = 0.8$) $h_{UP} = \frac{(8+0)}{2} \times 0.8$ ($= 3.2$) $(h_{UP} + 1.8) = \frac{(0+10)}{2} \times t_{DOWN}$ ($\Rightarrow t_{DOWN} = 1$) $T = t_{UP} + t_{DOWN}$ ALT 2: time to A + time from A to ground e.g. $8 = -8 + gt_A$ ($\Rightarrow t_A = 1.6$) $1.8 = \frac{(8+10)}{2} \times t_{AG}$ ($\Rightarrow t_{AG} = 0.2$) $T = t_A + t_{AG}$	M1	3.4
	$T = 1.8$ oe e.g. $9/5$	A1	1.1b
		(2)	
	e.g. Use a more accurate (less rounded) value for g (or gravity), use $g = 9.8$ or $g = 9.81$, allow for wind effects, allow for the spin of the stone, include dimensions of stone (not a particle), shape and/or size of stone, allow for variable acceleration. If air resistance is mentioned as an extra, ignore it.	B1	3.5c

			(1)	
(d)		U would be greater. Allow without U , e.g it would be greater, or just 'greater' oe ISW	B1	3.5a
			(1)	
(7 marks)				
Notes:				
1a	M1	Use the model to obtain an equation in U only, condone sign errors, but M0 if using an incorrect formula.		
	A1	A correct equation in U only, g does not need to be substituted (so allow $g = 9.8$ or 9.81)		
	A1	cao (A0 if $g = 10$ has not been used)		
1b	M1	Use the model to obtain an equation in T only, g does not need to be substituted (so allow $g = 9.8$ or 9.81) condone sign errors, but M0 if using an incorrect formula. Follow through on their U where necessary		
	A1	cao (A0 if $g = 10$ has not been used) A0 if they give two answers.		
1c	B1	Any appropriate refinement. B0 if an incorrect extra is given e.g. the mass or weight is mentioned		
1d	B1	cao		

Question	Scheme	Marks	AOs
2(a)		B1	1.1b
		(1)	
(b)	<p>Using <i>total</i> area = 15000 to set up an <i>equation</i> in one unknown Or they may use <i>suvat</i> on one or more sections (but must still be considering <i>all</i> sections)</p> <p>Allow an attempt at a clear explicit verification using $t = 40$ e.g. the following would score M1A1A1*: $4 \times 40 = 160$ then $700 - 40 - 160 = 500$ $\frac{(700 + 500)}{2} \times 25 = 15000 = 15 \text{ km}$ Withhold A1* if they don't include $= 15 \text{ km}$</p> <p>N.B. M0 if a single <i>suvat</i> formula is used for the whole journey.</p>	M1	3.4
	$\frac{1}{2}(700 + 700 - t - 4t) \times 25 = 15000$ <p>OR $\frac{1}{2} \times 25 \times t + 25(700 - t - 4t) + \frac{1}{2} \times 25 \times 4t = 15000$</p>	A1	1.1b
	$t = 40 \text{ (s)}^*$	A1*	1.1b
		(3)	
(c)	0.63 or 0.625 or $\frac{5}{8}$ oe (m s ⁻²) isw	B1	1.1b/ (2.2a)
		(1)	
(d)	<p>Complete method to find the speed or velocity at $t = 572$ e.g. $\pm \left(25 - \left(32 \times \frac{5}{32} \right) \right)$ or $\pm \left(128 \times \frac{5}{32} \right)$ oe</p>	M1	3.1b
	20 (m s ⁻¹)	A1	1.1b
		(2)	
(e)	<p>e.g. (the train) cannot instantaneously change acceleration, (the train) won't move with <u>constant</u> acceleration, (the train) won't move with <u>constant</u> speed Allow negatives of these:</p>	B1	3.5b

	e.g. (The train) moving at constant speed, or just 'constant speed' or 'constant acceleration' (is a limitation of the model) Must be a limitation of the model, so friction or air resistance or size of train is B0. N.B. Ignore incorrect reasons following a correct answer.		
		(1)	
(8 marks)			

Notes:

2a	B1	Overall shape of graph, starting at the origin, with deceleration phase <i>longer</i> than the acceleration phase if nothing on the t -axis but ignore the relative lengths and allow if t (or 40) and $4t$ (or 160) are clearly and correctly marked. Ignore incorrect figs on the axes. This mark can be earned if the graph appears anywhere in qu 2.
2b	M1	Need <i>all</i> sections to be included, with <u>correct structure for each section</u> , with $\frac{1}{2}$'s where appropriate. <u>Allow</u> = 15 or 150 or 1500 etc instead of 15000
	A1	A correct equation in their t only, seen or implied (or with $t = 40$ for verification)
	A1*	cso. At least one line of working with brackets removed and t 's collected, or equivalent
2c	B1	cao
2d	M1	Any complete method, must have correct figs, but condone sign errors
	A1	cao. Must be positive and exact i.e must not come from rounding.
2e	B1	Any appropriate limitation <u>of the model</u> . B0 if any incorrect extra answers .

Question		Scheme	Marks	AOs
3(a)		Differentiate s wrt t	M1	3.1a
		$(v =) t^2 - 5t + 6$	A1	1.1b
		Equate their v to 0 and solve	M1	1.1b
		$t = 2$ or 3	A1	1.1b
		$(a =) 2t - 5$	B1ft	2.1
		$a = 1$ and -1 (m s^{-2}) isw (A0 if extras)	A1	1.1b
			(6)	
(b)		Attempt to find values of s for $t = 2, 3$ and 4 oe Correct values are $\left(s_2 = \frac{14}{3}, s_3 = \frac{9}{2} \text{ and } s_4 = \frac{16}{3}\right)$ Could be implied by correct values for: s_2 , $(s_3 - s_2)$ and $(s_4 - s_3)$ which are $\frac{14}{3}$, $(-\frac{1}{6})$ and $\frac{5}{6}$	DM1	1.1b
		Total distance travelled $= s_2 + (s_2 - s_3) + s_4 - s_3$ OR $s_2 - (s_3 - s_2) + s_4 - s_3$ OR $\left[\frac{1}{3}t^3 - \frac{5}{2}t^2 + 6t\right]_0^2 - \left[\frac{1}{3}t^3 - \frac{5}{2}t^2 + 6t\right]_2^3 + \left[\frac{1}{3}t^3 - \frac{5}{2}t^2 + 6t\right]_3^4$ OR $\frac{14}{3} - (-\frac{1}{6}) + \frac{5}{6}$ OR $s_2 + 2(s_2 - s_3) + s_4 - s_2$ $(= 2s_2 - 2s_3 + s_4)$ oe	M1	2.1
		$5\frac{2}{3}$ oe (m) Accept 5.7 or better	A1	1.1b
			(3)	
(9 marks)				
Notes:				
3a	M1	Differentiate, with at least 2 powers decreasing by 1		
	A1	Correct expression		
	M1	Must have attempted to differentiate s to find v and be solving a 3 term quadratic		
	A1	Both values needed		
	B1ft	Follow their v (must be differentiating)		

	A1	cao
3b	DM 1	<u>This mark is dependent on the 2nd M1 in part (a) and their t values are between 0 and 4.</u> Clear attempt to find all three s values (may integrate their v incorrectly) N.B. No penalty for extra values.
	M1	Complete method using their s values Do NOT condone sign errors.
	A1	Any equivalent fraction, 5.7 or better.
		S.C. Correct answer, with no working, scores all 3 marks, since $\int_0^4 t^2 - 5t + 6 dt$ entered on a calculator will give $\frac{17}{3}$

Question	Scheme		Marks	AOs
	N.B. Use the mass in the ‘ma’ term of an equation to determine which part of the system (cage and block, cage or block) it applies to.			
4(a)	Translate situation into the model and set up the equation of motion for the <u>cage and the block</u> to obtain an equation in T only.		M1	3.3
	$T - 40g - 10g = 50 \times 0.2$		A1	1.1b
	500 (N) Must be positive		A1	1.1b
	Some examples: $T - 50 = 50 \times 0.2$ and $T - 40g - 10g = 50g \times 0.2$ both score M1A0A0			
			(3)	
(b)	Use the model to set up the equation of motion for the <u>block</u> to obtain an equation in R only.		M1	3.4
	$R - 10g = 10 \times 0.2$ Allow - R instead of R		A1	1.1b
	100 (N) Must be positive.		A1	1.1b
	OR: Use the model to set up the equation of motion for the <u>cage</u> to obtain an equation in R only.		M1	3.4
	$T - 40g - R = 40 \times 0.2$ with their T substituted		A1	1.1b
	100 (N) Must be positive		A1	1.1b
			(3)	
(6 marks)				
Notes: N.B. Only penalise the use of an incorrect value of g ONCE for the whole question, so max (a) M1A1A0 (b) M1A1A1				
4a	M1	Correct number of terms, condone sign errors		
	A1	Correct equation in T only		
	A1	cao		
4b	M1	Correct number of terms, condone sign errors		
	A1	Correct equation in R only		
	A1	cao		