

①

Name - Siddhart B  
 Grade - X  
 Board - CISCE

Date - 24/7/20

Sub - Phy.  
 Chapters - Light

Page - 6

Q1.

i) Light is not absorbed where?

**Write complete sentences**

$$\frac{23}{30}$$

Ans:-

ii)

1) Angle of incidence ✓

2) Thickness of the glass ✗

3) Angle of the Prism ✓

iii)

1) There should be one polished ~~soft~~ surface.

2) The glass should be thick.

iv) The beam of light moves towards the base while it passes through the prism.

v) If the image appears magnified (i.e. place the lens ~~at~~ a little far from the printed page) then it is a convex lens. However, if it appears diminished, then it is a ~~concave~~ lens.

vi) Glass absorbs U.V rays due to which quartz envelope is preferred as it can easily pass through ~~heat~~.

vii) Upon bringing the blackened <sup>bulb</sup> end of a thermo-metro beyond the red light (once obtaining the spectrum through dispersion), the

FOR EDUCATIONAL USE

(2)

temperature of the thermometer rises. Hence the infrared rays can be detected (as they can cause sudden rise in temperature).

viii)

- i) If it is +, then the image is real.
- If it is -, then the image is virtual.

- ii) If it is +, then the image is erect.
- If it is -, then the image is inverted.

ix) Infrared waves ~~rays~~Micro-waves ~~X~~

x) Given:-

$$\lambda = 5500 \text{ Å}$$

$$v = 3 \times 10^8 \text{ ms}^{-1}$$

$$\therefore f_b = \frac{\lambda}{v}$$

$$= \frac{3 \times 10^8}{5500}$$

$$= \frac{3 \times 10^6}{55}$$

$$= 5.454 \times 10^4 \text{ Hz}$$

$$= 5.454 \times 10^4 \text{ Hz}$$

$$\therefore f_b = 5.454 \times 10^4 \text{ Hz}$$

$$\frac{3 \cdot 818}{5454}$$


(3)

Q2&gt;

Gamma rays, X-rays, infrared rays, micro-waves, radio waves


Q3&gt;

RED +  
BLUE +  
YELLOW →



2

Q4&gt;



2

Object placed at  $2 F_1$ .

Q5>

- 1> It is absorbed by glass.
- 2> Can cause skin burns
- 3> It is not visible.

2

Q6>

$$u = -45\text{ cm}$$

a) Convex lens.

b)  $u = -45\text{ cm}$

$$v = 90\text{ cm}$$

$$\therefore \frac{1}{f} = \frac{1}{v} - \frac{1}{u}$$

$$\begin{aligned}\therefore &= \frac{1}{90} - \left(-\frac{1}{45}\right) \\ &= \frac{1+2}{90}\end{aligned}$$

$$\frac{1}{f} = \frac{-3}{90} = -\frac{1}{30}$$

3

f

$\therefore f = 30\text{ cm}$


$\therefore b = 30\text{ cm}$

c)  $m = \frac{v}{u} = \frac{90}{-45} = -2$

$\therefore m = -2$

Q7>

i)



Sundaram®

FOR EDUCATIONAL USE

2021/07/24 17:13

(5)

ii)  $\angle i = \angle e + 60^\circ \text{ } \textcircled{1}$

$$(\angle i + \angle i) - (\angle r + \angle r) = s$$

However,

$$\begin{aligned} \angle A &= \angle i + \angle i - 60 - (180 - (\angle r + \angle r)) \\ &= -120 + \angle r + \angle r \end{aligned}$$

$$\begin{aligned} s + \angle A &= 60 - \angle i + \angle i - 120 \\ &= -60^\circ + 60^\circ - 120^\circ \end{aligned}$$


$$s + \angle A = 0$$

$$s + \angle A = \angle i + \angle i$$

2/2


Q8>

a)



3

b)




FOR EDUCATIONAL USE

Sundaram®

2021/07/24 17:15

Q9>

i)



ii) L.L - Optical Centre Line.

O -  $F_2$

O' -  $F_1$

iii)

- 1) Virtual.
- 2) Erect.
- 3) Diminished.

3