

MA 205 Tutorial Batch 3

Recap-5

Siddhant Midha

10th September 2022

Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$.

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let γ be any positively oriented simple closed contour around p lying in the annulus.

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let γ be any positively oriented simple closed contour around p lying in the annulus. Then,

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let γ be any positively oriented simple closed contour around p lying in the annulus. Then,

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

holds for all $z \in \mathcal{A}(p, r_1, r_2)$.

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let γ be any positively oriented simple closed contour around p lying in the annulus. Then,

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

holds for all $z \in \mathcal{A}(p, r_1, r_2)$. Where,

$$a_n := \frac{1}{2\pi i} \int_{\gamma} \frac{f(\eta)}{(\eta - p)^{n+1}} \quad \forall n \in \mathbb{Z}$$

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let γ be any positively oriented simple closed contour around p lying in the annulus. Then,

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

holds for all $z \in \mathcal{A}(p, r_1, r_2)$. Where,

$$a_n := \frac{1}{2\pi i} \int_{\gamma} \frac{f(\eta)}{(\eta - p)^{n+1}} \quad \forall n \in \mathbb{Z}$$

- The **principal part** of the Laurent Series of f at p is

$$\sum_{n \leq -1} a_n (z - p)^n$$

Question

Give the Laurent Series for,

Question

Give the Laurent Series for,

① $f(z) = 1/z^{10231}$ in $\{z : 0 < |z| < 1\}$

Question

Give the Laurent Series for,

- ① $f(z) = 1/z^{10231}$ in $\{z : 0 < |z| < 1\}$
- ② $f(z) = 1/z(z+1)$ in $\{z : 0 < |z| < 1\}$,

Question

Give the Laurent Series for,

① $f(z) = 1/z^{10231}$ in $\{z : 0 < |z| < 1\}$

② $f(z) = 1/z(z+1)$ in $\{z : 0 < |z| < 1\}, \{z : |z| > 1\}$

Question

Give the Laurent Series for,

- ① $f(z) = 1/z^{10231}$ in $\{z : 0 < |z| < 1\}$
- ② $f(z) = 1/z(z+1)$ in $\{z : 0 < |z| < 1\}, \{z : |z| > 1\}$
- ③ $f(z) = 1/(z^2 + 1)$ in $\{z : |z| < 1\}$,

Question

Give the Laurent Series for,

- ① $f(z) = 1/z^{10231}$ in $\{z : 0 < |z| < 1\}$
- ② $f(z) = 1/z(z+1)$ in $\{z : 0 < |z| < 1\}, \{z : |z| > 1\}$
- ③ $f(z) = 1/(z^2 + 1)$ in $\{z : |z| < 1\}, \{z : |z - i| < 1\}$

Question

Give the Laurent Series for,

- ① $f(z) = 1/z^{10231}$ in $\{z : 0 < |z| < 1\}$
- ② $f(z) = 1/z(z+1)$ in $\{z : 0 < |z| < 1\}, \{z : |z| > 1\}$
- ③ $f(z) = 1/(z^2 + 1)$ in $\{z : |z| < 1\}, \{z : |z - i| < 1\}$
- ④ $f(z) = e^{1/z}$

Question

Give the Laurent Series for,

- ① $f(z) = 1/z^{10231}$ in $\{z : 0 < |z| < 1\}$
- ② $f(z) = 1/z(z+1)$ in $\{z : 0 < |z| < 1\}, \{z : |z| > 1\}$
- ③ $f(z) = 1/(z^2 + 1)$ in $\{z : |z| < 1\}, \{z : |z - i| < 1\}$
- ④ $f(z) = e^{1/z}$

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$.

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series.

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series. Then,

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series. Then,

- ① f has a pole at $p \Leftrightarrow a_n = 0$ for all but finitely many $n < 0$, and $a_k \neq 0$ for some $k < 0$.

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series. Then,

- 1 f has a pole at $p \Leftrightarrow a_n = 0$ for all but finitely many $n < 0$, and $a_k \neq 0$ for some $k < 0$.
- 2 f has an essential singularity at $p \Leftrightarrow a_n \neq 0$ for **infinitely many** $n < 0$.

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series. Then,

- 1 f has a pole at $p \Leftrightarrow a_n = 0$ for all but finitely many $n < 0$, and $a_k \neq 0$ for some $k < 0$.
- 2 f has an essential singularity at $p \Leftrightarrow a_n \neq 0$ for **infinitely many** $n < 0$.

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series. Then,

- ① f has a pole at $p \Leftrightarrow a_n = 0$ for all but finitely many $n < 0$, and $a_k \neq 0$ for some $k < 0$.
- ② f has an essential singularity at $p \Leftrightarrow a_n \neq 0$ for **infinitely many** $n < 0$.

Thus,

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series. Then,

- 1 f has a pole at $p \Leftrightarrow a_n = 0$ for all but finitely many $n < 0$, and $a_k \neq 0$ for some $k < 0$.
- 2 f has an essential singularity at $p \Leftrightarrow a_n \neq 0$ for **infinitely many** $n < 0$.

Thus,

- 1 The singularity is removable iff the principal part is zero.

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series. Then,

- ① f has a pole at $p \Leftrightarrow a_n = 0$ for all but finitely many $n < 0$, and $a_k \neq 0$ for some $k < 0$.
- ② f has an essential singularity at $p \Leftrightarrow a_n \neq 0$ for **infinitely many** $n < 0$.

Thus,

- ① The singularity is removable iff the principal part is zero.
- ② The singularity is a pole iff the principal part has a finite (not zero) number of non-zero terms.

A consequence of the Laurent Series

Theorem

Suppose f is analytic in the annulus $\mathcal{A}(p, r_1, r_2) := \{z : r_1 < |z - p| < r_2\}$. Let a_n be the coefficients of its Laurent Series. Then,

- 1 f has a pole at $p \Leftrightarrow a_n = 0$ for all but finitely many $n < 0$, and $a_k \neq 0$ for some $k < 0$.
- 2 f has an essential singularity at $p \Leftrightarrow a_n \neq 0$ for **infinitely many** $n < 0$.

Thus,

- 1 The singularity is removable iff the principal part is zero.
- 2 The singularity is a pole iff the principal part has a finite (not zero) number of non-zero terms.
- 3 The singularity is essential iff the principal part has an infinite number of non-zero terms.

Residue Theorem

Residue Theorem

Definition (Residue at a point)

Residue Theorem

Definition (Residue at a point)

Suppose f is analytic in the annulus $\mathcal{A}(p, 0, r_2) := \{z : 0 < |z - p| < r_2\}$.

Residue Theorem

Definition (Residue at a point)

Suppose f is analytic in the annulus $\mathcal{A}(p, 0, r_2) := \{z : 0 < |z - p| < r_2\}$.
Let

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

Residue Theorem

Definition (Residue at a point)

Suppose f is analytic in the annulus $\mathcal{A}(p, 0, r_2) := \{z : 0 < |z - p| < r_2\}$.
Let

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

be the Laurent Series expansion of f in this annulus.

Residue Theorem

Definition (Residue at a point)

Suppose f is analytic in the annulus $\mathcal{A}(p, 0, r_2) := \{z : 0 < |z - p| < r_2\}$.
Let

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

be the Laurent Series expansion of f in this annulus. Then, we define,

$$\text{Res}(f, p) := a_{-1}$$

Residue Theorem

Definition (Residue at a point)

Suppose f is analytic in the annulus $\mathcal{A}(p, 0, r_2) := \{z : 0 < |z - p| < r_2\}$.
Let

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

be the Laurent Series expansion of f in this annulus. Then, we define,

$$\text{Res}(f, p) := a_{-1}$$

Residue Theorem

Definition (Residue at a point)

Suppose f is analytic in the annulus $\mathcal{A}(p, 0, r_2) := \{z : 0 < |z - p| < r_2\}$.
Let

$$f(z) = \sum_{n \in \mathbb{Z}} a_n (z - p)^n$$

be the Laurent Series expansion of f in this annulus. Then, we define,

$$\text{Res}(f, p) := a_{-1}$$

As an immediate consequence, see that

$$\int_C f(z) dz = 2\pi i \times \text{Res}(f, p)$$

Residues of Meromorphic functions

Theorem

If f is analytic in $D^*(p, r)$ for some r and has a pole of order k at p , then,

Residues of Meromorphic functions

Theorem

If f is analytic in $D^*(p, r)$ for some r and has a pole of order k at p , then,

$$\operatorname{Res}(f, p) = \lim_{z \rightarrow p} \frac{1}{(k-1)!} \frac{d^{k-1}}{dz^{k-1}} \left((z-p)^k f(z) \right)$$

Questions

- 1 Find residue at 0 of $1/(z + z^2)$.

Questions

- 1 Find residue at 0 of $1/(z + z^2)$.
- 2 Find residue at 0 of $1/(z^2(1 - z))$ using *both* methods.

Questions

- 1 Find residue at 0 of $1/(z + z^2)$.
- 2 Find residue at 0 of $1/(z^2(1 - z))$ using *both* methods.
- 3 $\int_{|z|=1} e^{1/z^2} dz$

Questions

- 1 Find residue at 0 of $1/(z + z^2)$.
- 2 Find residue at 0 of $1/(z^2(1 - z))$ using *both* methods.
- 3 $\int_{|z|=1} e^{1/z^2} dz$
- 4 $\int_{|z|=1} \frac{1}{\sin 1/z} dz$

Questions

- 1 Find residue at 0 of $1/(z + z^2)$.
- 2 Find residue at 0 of $1/(z^2(1 - z))$ using *both* methods.
- 3 $\int_{|z|=1} e^{1/z^2} dz$
- 4 $\int_{|z|=1} \frac{1}{\sin 1/z} dz$ Uh? What? CRT? No!

The Cauchy Residue Theorem

This is a formal restatement of the earlier conclusion.

Theorem (CRT)

Let γ be a simple closed contour.

The Cauchy Residue Theorem

This is a formal restatement of the earlier conclusion.

Theorem (CRT)

Let γ be a simple closed contour. Let f be analytic inside and on γ except for a finite number of isolated singularities p_k ,

The Cauchy Residue Theorem

This is a formal restatement of the earlier conclusion.

Theorem (CRT)

Let γ be a simple closed contour. Let f be analytic inside and on γ except for a finite number of isolated singularities p_k , then,

$$\int_{\gamma} f(z) dz = 2\pi i \sum_k \text{Res}(f, p_k)$$

Orders and Multiplicities

(Recall)

Theorem (The Order of a Pole)

Let f have a pole at $p \in \mathbb{C}$.

(Recall)

Theorem (The Order of a Pole)

Let f have a pole at $p \in \mathbb{C}$. Then, there exists some $k \in \mathbb{N}$ such that for some $r > 0$ we have,

(Recall)

Theorem (The Order of a Pole)

Let f have a pole at $p \in \mathbb{C}$. Then, there exists some $k \in \mathbb{N}$ such that for some $r > 0$ we have,

$$f(z) = (z - p)^{-k} H(z)$$

(Recall)

Theorem (The Order of a Pole)

Let f have a pole at $p \in \mathbb{C}$. Then, there exists some $k \in \mathbb{N}$ such that for some $r > 0$ we have,

$$f(z) = (z - p)^{-k} H(z)$$

where, H is holomorphic on $D(p, r)$ **and** $H(p) \neq 0$.

(Recall)

Theorem (The Order of a Pole)

Let f have a pole at $p \in \mathbb{C}$. Then, there exists some $k \in \mathbb{N}$ such that for some $r > 0$ we have,

$$f(z) = (z - p)^{-k} H(z)$$

where, H is holomorphic on $D(p, r)$ **and** $H(p) \neq 0$. We say that k is the **order of the pole**.

(Recall)

Theorem (The Order of a Pole)

Let f have a pole at $p \in \mathbb{C}$. Then, there exists some $k \in \mathbb{N}$ such that for some $r > 0$ we have,

$$f(z) = (z - p)^{-k} H(z)$$

where, H is holomorphic on $D(p, r)$ **and** $H(p) \neq 0$. We say that k is the **order of the pole**.

Orders and Multiplicities

And similarly, we define,

Orders and Multiplicities

And similarly, we define,

Definition (Multiplicity of a zero)

Orders and Multiplicities

And similarly, we define,

Definition (Multiplicity of a zero)

Given a function f ,

Orders and Multiplicities

And similarly, we define,

Definition (Multiplicity of a zero)

Given a function f , we say that $z_0 \in \mathbb{C}$ is a zero of multiplicity m of f , if

Orders and Multiplicities

And similarly, we define,

Definition (Multiplicity of a zero)

Given a function f , we say that $z_0 \in \mathbb{C}$ is a zero of multiplicity m of f , if

$$f^{(n)}(z_0) = 0 \quad \forall n \leq m-1 \quad \textbf{and} \quad f^{(m)}(z_0) \neq 0$$

Orders and Multiplicities

And similarly, we define,

Definition (Multiplicity of a zero)

Given a function f , we say that $z_0 \in \mathbb{C}$ is a zero of multiplicity m of f , if

$$f^{(n)}(z_0) = 0 \quad \forall n \leq m-1 \quad \textbf{and} \quad f^{(m)}(z_0) \neq 0$$

We have that,

Orders and Multiplicities

And similarly, we define,

Definition (Multiplicity of a zero)

Given a function f , we say that $z_0 \in \mathbb{C}$ is a zero of multiplicity m of f , if

$$f^{(n)}(z_0) = 0 \quad \forall n \leq m-1 \quad \textbf{and} \quad f^{(m)}(z_0) \neq 0$$

We have that,

$$f(z) = (z - z_0)^m g(z)$$

for some holomorphic g which does not vanish in a neighbourhood of z_0 .

Orders and Multiplicities

And similarly, we define,

Definition (Multiplicity of a zero)

Given a function f , we say that $z_0 \in \mathbb{C}$ is a zero of multiplicity m of f , if

$$f^{(n)}(z_0) = 0 \quad \forall n \leq m-1 \quad \textbf{and} \quad f^{(m)}(z_0) \neq 0$$

We have that,

$$f(z) = (z - z_0)^m g(z)$$

for some holomorphic g which does not vanish in a neighbourhood of z_0 .

The Argument Principle

This is a nice application of the CRT.

The Argument Principle

This is a nice application of the CRT.

Theorem

The Argument Principle

This is a nice application of the CRT.

Theorem

Let f be a meromorphic function on and inside some closed contour γ ,

The Argument Principle

This is a nice application of the CRT.

Theorem

Let f be a meromorphic function on and inside some closed contour γ ,
and has no poles or zeros on γ .

The Argument Principle

This is a nice application of the CRT.

Theorem

Let f be a meromorphic function on and inside some closed contour γ , **and** has no poles or zeros on γ . Then,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = Z - P$$

The Argument Principle

This is a nice application of the CRT.

Theorem

Let f be a meromorphic function on and inside some closed contour γ , **and** has no poles or zeros on γ . Then,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = Z - P$$

The Argument Principle

This is a nice application of the CRT.

Theorem

Let f be a meromorphic function on and inside some closed contour γ , **and** has no poles or zeros on γ . Then,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = Z - P$$

where, Z is the number of zeros of f inside γ **counted with multiplicities**

The Argument Principle

This is a nice application of the CRT.

Theorem

Let f be a meromorphic function on and inside some closed contour γ , **and** has no poles or zeros on γ . Then,

$$\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = Z - P$$

where, Z is the number of zeros of f inside γ **counted with multiplicities** and P is the number of poles of f inside γ **counted with order**.

- 1 Why did the mathematician name his dog Cauchy?

- 1 Why did the mathematician name his dog Cauchy? –?

- 1 Why did the mathematician name his dog Cauchy? –? Because he left a residue at each pole!

- 1 Why did the mathematician name his dog Cauchy? –? Because he left a residue at each pole!
- 2 What is the contour integral over the Western Europe?

- 1 Why did the mathematician name his dog Cauchy? –? Because he left a residue at each pole!
- 2 What is the contour integral over the Western Europe?



- 1 Why did the mathematician name his dog Cauchy? –? Because he left a residue at each pole!
- 2 What is the contour integral over the Western Europe?



Zero! All the poles are in Eastern Europe.

credit: MA 205 2021 Lectures

And that is all

And that is all

... for the recap's of MA 205!