

MA 109 D2 T1

Week Five Recap

Siddhant Midha

<https://siddhant-midha.github.io/>

December 7, 2022

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$.

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)
- Introduce a *partition* of $[a, b]$, viz.

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)
- Introduce a *partition* of $[a, b]$, viz.

$$\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$$

- Define the global quantities,

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)
- Introduce a *partition* of $[a, b]$, viz.

$$\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$$

- Define the global quantities,

$$m(f) := \inf\{f(x) \mid x \in [a, b]\} \quad ; \quad M(f) := \sup\{f(x) \mid x \in [a, b]\}$$

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)
- Introduce a *partition* of $[a, b]$, viz.

$$\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$$

- Define the global quantities,

$$m(f) := \inf\{f(x) \mid x \in [a, b]\} \quad ; \quad M(f) := \sup\{f(x) \mid x \in [a, b]\}$$

- Define the local quantities for all $i = 1, 2, \dots, n$

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)
- Introduce a *partition* of $[a, b]$, viz.

$$\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$$

- Define the global quantities,

$$m(f) := \inf\{f(x) \mid x \in [a, b]\} \quad ; \quad M(f) := \sup\{f(x) \mid x \in [a, b]\}$$

- Define the local quantities for all $i = 1, 2, \dots, n$

$$m_i(f) := \inf\{f(x) \mid x \in [x_{i-1}, x_i]\} \quad ; \quad M_i(f) := \sup\{f(x) \mid x \in [x_{i-1}, x_i]\}$$

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)
- Introduce a *partition* of $[a, b]$, viz.

$$\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$$

- Define the global quantities,

$$m(f) := \inf\{f(x) \mid x \in [a, b]\} \quad ; \quad M(f) := \sup\{f(x) \mid x \in [a, b]\}$$

- Define the local quantities for all $i = 1, 2, \dots, n$

$$m_i(f) := \inf\{f(x) \mid x \in [x_{i-1}, x_i]\} \quad ; \quad M_i(f) := \sup\{f(x) \mid x \in [x_{i-1}, x_i]\}$$

- Note $m(f) \leq m_i(f) \leq M_i(f) \leq M(f) \quad \forall i$,

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)
- Introduce a *partition* of $[a, b]$, viz.

$$\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$$

- Define the global quantities,

$$m(f) := \inf\{f(x) \mid x \in [a, b]\} \quad ; \quad M(f) := \sup\{f(x) \mid x \in [a, b]\}$$

- Define the local quantities for all $i = 1, 2, \dots, n$

$$m_i(f) := \inf\{f(x) \mid x \in [x_{i-1}, x_i]\} \quad ; \quad M_i(f) := \sup\{f(x) \mid x \in [x_{i-1}, x_i]\}$$

- Note $m(f) \leq m_i(f) \leq M_i(f) \leq M(f) \quad \forall i$, and define the sums,

The Riemann Integral – Setting

The motivation: Rigorously define the notion of ‘area’.

- Identify a **bounded nonnegative** function $f : [a, b] \rightarrow \mathbb{R}$. (Draw Picture)
- Introduce a *partition* of $[a, b]$, viz.

$$\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$$

- Define the global quantities,

$$m(f) := \inf\{f(x) \mid x \in [a, b]\} \quad ; \quad M(f) := \sup\{f(x) \mid x \in [a, b]\}$$

- Define the local quantities for all $i = 1, 2, \dots, n$

$$m_i(f) := \inf\{f(x) \mid x \in [x_{i-1}, x_i]\} \quad ; \quad M_i(f) := \sup\{f(x) \mid x \in [x_{i-1}, x_i]\}$$

- Note $m(f) \leq m_i(f) \leq M_i(f) \leq M(f) \quad \forall i$, and define the sums,

$$L(f, \mathcal{P}) := \sum_{i=1}^n m_i(f)(x_i - x_{i-1}) \quad ; \quad U(f, \mathcal{P}) := \sum_{i=1}^n M_i(f)(x_i - x_{i-1}) \quad (\text{Draw!})$$

The Riemann Integral – Setting

- Define the *main players*

$$L(f) := \sup\{L(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

$$U(f) := \inf\{U(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

The Riemann Integral – Setting

- Define the *main players*

$$L(f) := \sup\{L(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

$$U(f) := \inf\{U(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

Need the lower sums to pump up, and the upper sums to power down in order to ‘approximate’. Now, $L(f) \leq U(f)$? Obvious?

The Riemann Integral – Setting

- Define the *main players*

$$L(f) := \sup\{L(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

$$U(f) := \inf\{U(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

Need the lower sums to pump up, and the upper sums to power down in order to ‘approximate’. Now, $L(f) \leq U(f)$? Obvious?

- A helpful definition: \mathcal{P}^* is said to be a refinement of \mathcal{P} if $\mathcal{P} \subseteq \mathcal{P}^*$. That is, \mathcal{P}^* is *finer* than \mathcal{P} .

The Riemann Integral – Setting

- Define the *main players*

$$L(f) := \sup\{L(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

$$U(f) := \inf\{U(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

Need the lower sums to pump up, and the upper sums to power down in order to ‘approximate’. Now, $L(f) \leq U(f)$? Obvious?

- A helpful definition: \mathcal{P}^* is said to be a refinement of \mathcal{P} if $\mathcal{P} \subseteq \mathcal{P}^*$. That is, \mathcal{P}^* is *finer* than \mathcal{P} .

Theorem

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function.

The Riemann Integral – Setting

- Define the *main players*

$$L(f) := \sup\{L(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

$$U(f) := \inf\{U(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

Need the lower sums to pump up, and the upper sums to power down in order to ‘approximate’. Now, $L(f) \leq U(f)$? Obvious?

- A helpful definition: \mathcal{P}^* is said to be a refinement of \mathcal{P} if $\mathcal{P} \subseteq \mathcal{P}^*$. That is, \mathcal{P}^* is *finer* than \mathcal{P} .

Theorem

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function.

- If \mathcal{P} is partition of $[a, b]$, and \mathcal{P}^* is a refinement of \mathcal{P} , then

$$L(f, \mathcal{P}) \leq L(f, \mathcal{P}^*) \quad \text{and} \quad U(f, \mathcal{P}^*) \leq U(f, \mathcal{P}),$$

- If \mathcal{P}_1 and \mathcal{P}_2 are partitions of $[a, b]$, then $L(f, \mathcal{P}_1) \leq U(f, \mathcal{P}_2)$.

The Riemann Integral – Setting

- Define the *main players*

$$L(f) := \sup\{L(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

$$U(f) := \inf\{U(f, \mathcal{P}) \mid \mathcal{P} \text{ is a partition of } [a, b]\}$$

Need the lower sums to pump up, and the upper sums to power down in order to ‘approximate’. Now, $L(f) \leq U(f)$? Obvious?

- A helpful definition: \mathcal{P}^* is said to be a refinement of \mathcal{P} if $\mathcal{P} \subseteq \mathcal{P}^*$. That is, \mathcal{P}^* is *finer* than \mathcal{P} .

Theorem

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function.

- If \mathcal{P} is partition of $[a, b]$, and \mathcal{P}^* is a refinement of \mathcal{P} , then

$$L(f, \mathcal{P}) \leq L(f, \mathcal{P}^*) \quad \text{and} \quad U(f, \mathcal{P}^*) \leq U(f, \mathcal{P}),$$

- If \mathcal{P}_1 and \mathcal{P}_2 are partitions of $[a, b]$, then $L(f, \mathcal{P}_1) \leq U(f, \mathcal{P}_2)$.
- $L(f) \leq U(f)$.

The Riemann Integral

Definition (The RI)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function.

The Riemann Integral

Definition (The RI)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then, f is said to be Riemann Integrable on $[a, b]$ if $U(f) = L(f)$.

The Riemann Integral

Definition (The RI)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then, f is said to be Riemann Integrable on $[a, b]$ if $U(f) = L(f)$. Further, we say that the quantity $U(f)$ ($= L(f)$) is the *Riemann Integral* of f on $[a, b]$, and denote it as

$$U(f) = L(f) = \int_a^b f(x) dx$$

The Riemann Condition

Theorem (The RC)

Given a bounded real function f on $[a, b]$,

The Riemann Condition

Theorem (The RC)

Given a bounded real function f on $[a, b]$, it holds that f is RI as per the previous definition **if and only if** the following holds,

The Riemann Condition

Theorem (The RC)

Given a bounded real function f on $[a, b]$, it holds that f is RI as per the previous definition **if and only if** the following holds,

$\forall \epsilon > 0$, there exists a partition \mathcal{P} of $[a, b]$

such that $U(f, \mathcal{P}) - L(f, \mathcal{P}) < \epsilon$

Two Funny Functions

- The **Dirichlet Function**, denoted $1_{\mathbb{Q}} : [0, 1] \rightarrow \mathbb{R}$, defined as,

$$1_{\mathbb{Q}} := \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$$

Two Funny Functions

- The **Dirichlet Function**, denoted $1_{\mathbb{Q}} : [0, 1] \rightarrow \mathbb{R}$, defined as,

$$1_{\mathbb{Q}} := \begin{cases} 1 & \text{if } x \in \mathbb{Q} \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$$

- The **Thomae Function**, denoted $T : [0, 1] \rightarrow \mathbb{R}$, defined as,

$$T(x) := \begin{cases} \frac{1}{q} & \text{if } x \in \mathbb{Q}, x \neq 0, x = p/q \text{ with } p, q \text{ coprime} \\ 0 & \text{if } x = 0 \\ 0 & \text{if } x \notin \mathbb{Q} \end{cases}$$

Proposition (Domain Additivity)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function and let $c \in (a, b)$. Then f is integrable on $[a, b]$ if and only if f is integrable on $[a, c]$ and on $[c, b]$. In this case,

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

Proposition (Domain Additivity)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function and let $c \in (a, b)$. Then f is integrable on $[a, b]$ if and only if f is integrable on $[a, c]$ and on $[c, b]$. In this case,

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx.$$

Proposition (Order Relations)

Let $f, g : [a, b] \rightarrow \mathbb{R}$ be integrable. (i) If $f \leq g$, then

$$\int_a^b f(x)dx \leq \int_a^b g(x)dx. \quad \text{(ii) The function } |f| \text{ is integrable and}$$
$$\left| \int_a^b f(x)dx \right| \leq \int_a^b |f|(x)dx.$$

Proposition (Algebraic and order relations)

Let $f, g : [a, b] \rightarrow \mathbb{R}$ be integrable functions. Then

- $f + g$ is integrable and $\int_a^b (f + g)(x)dx = \int_a^b f(x)dx + \int_a^b g(x)dx$,
- rf is integrable for any $r \in \mathbb{R}$ and $\int_a^b (rf)(x)dx = r \int_a^b f(x)dx$,
- fg is integrable,
- if there is $\delta > 0$ such that $|f(x)| \geq \delta$ and all $x \in [a, b]$, then $1/f$ is integrable,
- if $f(x) \geq 0$ for all $x \in [a, b]$, then for any $k \in \mathbb{N}$, the function $f^{1/k}$ is integrable.

Lemma

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function.

Lemma

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then,

- 1 If f is monotonic, it is integrable.

Integrable functions

Lemma

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then,

- 1 If f is monotonic, it is integrable.
- 2 If f is continuous, it is integrable.

Now,

Integrable functions

Lemma

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then,

- 1 If f is monotonic, it is integrable.
- 2 If f is continuous, it is integrable.

Now,

- Integrability \implies continuity?

Integrable functions

Lemma

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then,

- 1 If f is monotonic, it is integrable.
- 2 If f is continuous, it is integrable.

Now,

- Integrability \implies continuity? No!
- Integrability \implies boundedness?

Integrable functions

Lemma

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then,

- 1 If f is monotonic, it is integrable.
- 2 If f is continuous, it is integrable.

Now,

- Integrability \implies continuity? No!
- Integrability \implies boundedness? **Yes.**
- Integrability \implies monotonicity?

Integrable functions

Lemma

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Then,

- 1 If f is monotonic, it is integrable.
- 2 If f is continuous, it is integrable.

Now,

- Integrability \implies continuity? No!
- Integrability \implies boundedness? **Yes**.
- Integrability \implies monotonicity? No!

The Fundamental Theorem of Calculus

The FTC connects together the notions of integrability and differentiability. That is,

$$\text{Integrability} \xleftrightarrow{\text{FTC}} \text{Differentiability}$$

The Fundamental Theorem of Calculus

The FTC connects together the notions of integrability and differentiability. That is,

$$\text{Integrability} \xleftrightarrow{\text{FTC}} \text{Differentiability}$$

Also, a helpful definition.

Definition (Antiderivative)

Given a function $f : D \rightarrow \mathbb{R}$, we say that f has a **antiderivative** on D if there exists a differentiable function $F : D \rightarrow \mathbb{R}$ such that,

$$F'(x) = f(x) \quad \forall x \in D$$

FTC I

The FTC I connects integration to differentiation.

The FTC I connects integration to differentiation.

Theorem (FTC I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable, and define, $F : [a, b] \rightarrow \mathbb{R}$ as,

$$F(x) := \int_a^x f(x) dx$$

The FTC I connects integration to differentiation.

Theorem (FTC I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable, and define, $F : [a, b] \rightarrow \mathbb{R}$ as,

$$F(x) := \int_a^x f(x) dx$$

Then, the following holds,

- F is continuous on $[a, b]$.

The FTC I connects integration to differentiation.

Theorem (FTC I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable, and define, $F : [a, b] \rightarrow \mathbb{R}$ as,

$$F(x) := \int_a^x f(x) dx$$

Then, the following holds,

- F is continuous on $[a, b]$.
- If f is continuous at $c \in [a, b]$, then F is differentiable at c . Further, $F'(c) = f(c)$.

The FTC I connects integration to differentiation.

Theorem (FTC I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable, and define, $F : [a, b] \rightarrow \mathbb{R}$ as,

$$F(x) := \int_a^x f(x) dx$$

Then, the following holds,

- F is continuous on $[a, b]$.
- If f is continuous at $c \in [a, b]$, then F is differentiable at c . Further, $F'(c) = f(c)$.

That is, if your f is continuous, the proposed F is an antiderivative. Thus a continuous function on an interval always possesses an antiderivative.

Theorem (FTC II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable,

Theorem (FTC II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable, and let F be *any* antiderivative of f . That is, $F' = f$.

Theorem (FTC II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable, and let F be *any* antiderivative of f . That is, $F' = f$. Then,

$$\int_a^b f(x) dx = F(b) - F(a)$$

Theorem (FTC II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable, and let F be *any* antiderivative of f . That is, $F' = f$. Then,

$$\int_a^b f(x) dx = F(b) - F(a)$$

Another way to look at this, is that,

$$F(x) = F(a) + \int_a^x f(x) dx$$

Theorem (FTC II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable, and let F be *any* antiderivative of f . That is, $F' = f$. Then,

$$\int_a^b f(x) dx = F(b) - F(a)$$

Another way to look at this, is that,

$$F(x) = F(a) + \int_a^x f(x) dx$$

for *any* antiderivative F of f . This also relates to the fact that two antiderivatives differ by a constant.

Riemann Sums

Up till now, we dealt with some forms of defining integrability. While this enabled us to have a rigorous formalism, one may ask – how do you perform calculations? Is taking sup and inf every time not tedious?

Riemann Sums

Up till now, we dealt with some forms of defining integrability. While this enabled us to have a rigorous formalism, one may ask – how do you perform calculations? Is taking sup and inf every time not tedious?

Definition (Tagged Partition)

Let $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$ be a partition of $[a, b]$. Let $t_i \in [x_{i-1}, x_i]$ for $i = 1, 2, \dots, n$ be arbitrary, and denote $t := \{t_1 < t_2 < \cdots < t_n\}$. We call the tuple (\mathcal{P}, t) a **tagged partition**.

Riemann Sums

Up till now, we dealt with some forms of defining integrability. While this enabled us to have a rigorous formalism, one may ask – how do you perform calculations? Is taking sup and inf every time not tedious?

Definition (Tagged Partition)

Let $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$ be a partition of $[a, b]$. Let $t_i \in [x_{i-1}, x_i]$ for $i = 1, 2, \dots, n$ be arbitrary, and denote $t := \{t_1 < t_2 < \cdots < t_n\}$. We call the tuple (\mathcal{P}, t) a **tagged partition**.

Definition (Riemann Sum)

Given $f : [a, b] \rightarrow \mathbb{R}$, and a tagged partition (\mathcal{P}, t) of $[a, b]$, define

$$R(f, \mathcal{P}, t) := \sum_{i=1}^n f(t_i)(x_i - x_{i-1})$$

to be the associated **Riemann Sum**.

Riemann Sums

Up till now, we dealt with some forms of defining integrability. While this enabled us to have a rigorous formalism, one may ask – how do you perform calculations? Is taking sup and inf every time not tedious?

Definition (Tagged Partition)

Let $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$ be a partition of $[a, b]$. Let $t_i \in [x_{i-1}, x_i]$ for $i = 1, 2, \dots, n$ be arbitrary, and denote $t := \{t_1 < t_2 < \cdots < t_n\}$. We call the tuple (\mathcal{P}, t) a **tagged partition**.

Definition (Riemann Sum)

Given $f : [a, b] \rightarrow \mathbb{R}$, and a tagged partition (\mathcal{P}, t) of $[a, b]$, define

$$R(f, \mathcal{P}, t) := \sum_{i=1}^n f(t_i)(x_i - x_{i-1})$$

to be the associated **Riemann Sum**.

Note that, $L(f, \mathcal{P}) \leq R(f, \mathcal{P}, t) \leq U(f, \mathcal{P})$.

A Definition

Further for a partition $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$, define $\|\mathcal{P}\| := \max\{x_i - x_{i-1}\}$ to be the *norm* of that partition.

A Definition

Further for a partition $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$, define $\|\mathcal{P}\| := \max\{x_i - x_{i-1}\}$ to be the *norm* of that partition.

Definition (RS Def I)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable

A Definition

Further for a partition $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$, define $\|\mathcal{P}\| := \max\{x_i - x_{i-1}\}$ to be the *norm* of that partition.

Definition (RS Def I)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$

A Definition

Further for a partition $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$, define $\|\mathcal{P}\| := \max\{x_i - x_{i-1}\}$ to be the *norm* of that partition.

Definition (RS Def I)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$

A Definition

Further for a partition $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$, define $\|\mathcal{P}\| := \max\{x_i - x_{i-1}\}$ to be the *norm* of that partition.

Definition (RS Def I)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|R(f, \mathcal{P}, t) - R| < \epsilon,$$

whenever $\|\mathcal{P}\| < \delta$

A Definition

Further for a partition $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$, define $\|\mathcal{P}\| := \max\{x_i - x_{i-1}\}$ to be the *norm* of that partition.

Definition (RS Def I)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|R(f, \mathcal{P}, t) - R| < \epsilon,$$

whenever $\|\mathcal{P}\| < \delta$ and for all choice of tags t .

A Definition

Further for a partition $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$, define $\|\mathcal{P}\| := \max\{x_i - x_{i-1}\}$ to be the *norm* of that partition.

Definition (RS Def I)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|R(f, \mathcal{P}, t) - R| < \epsilon,$$

whenever $\|\mathcal{P}\| < \delta$ and for all choice of tags t . In this case R is called the Riemann integral of the function f on the interval $[a, b]$.

A Definition

Further for a partition $\mathcal{P} := \{x_0 = a < x_1 < x_2 < \cdots < x_{n-1} < x_n = b\}$, define $\|\mathcal{P}\| := \max\{x_i - x_{i-1}\}$ to be the *norm* of that partition.

Definition (RS Def I)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists $\delta > 0$ such that

$$|R(f, \mathcal{P}, t) - R| < \epsilon,$$

whenever $\|\mathcal{P}\| < \delta$ and for all choice of tags t . In this case R is called the Riemann integral of the function f on the interval $[a, b]$.

A weaker definition

Definition (RS Def II)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists a $\delta > 0$ **and a** partition \mathcal{P} such that **for every tagged refinement** (\mathcal{P}', t') of \mathcal{P} with $\|\mathcal{P}'\| \leq \delta$ we have,

$$|R(f, \mathcal{P}', t') - R| < \epsilon.$$

A weaker definition

Definition (RS Def II)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists a $\delta > 0$ **and a** partition \mathcal{P} such that **for every tagged refinement** (\mathcal{P}', t') of \mathcal{P} with $\|\mathcal{P}'\| \leq \delta$ we have,

$$|R(f, \mathcal{P}', t') - R| < \epsilon.$$

The nice thing about the above definition is that one only has to check that $|R(f, \mathcal{P}', t') - R|$ is small for refinements of a fixed partition, and not for all partitions.

A weaker definition

Definition (RS Def II)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists a $\delta > 0$ **and a** partition \mathcal{P} such that **for every tagged refinement** (\mathcal{P}', t') of \mathcal{P} with $\|\mathcal{P}'\| \leq \delta$ we have,

$$|R(f, \mathcal{P}', t') - R| < \epsilon.$$

The nice thing about the above definition is that one only has to check that $|R(f, \mathcal{P}', t') - R|$ is small for refinements of a fixed partition, and not for all partitions. Further, note that if $\mathcal{P} \subset \mathcal{P}^*$, then

A weaker definition

Definition (RS Def II)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists a $\delta > 0$ **and a** partition \mathcal{P} such that **for every tagged refinement** (\mathcal{P}', t') of \mathcal{P} with $\|\mathcal{P}'\| \leq \delta$ we have,

$$|R(f, \mathcal{P}', t') - R| < \epsilon.$$

The nice thing about the above definition is that one only has to check that $|R(f, \mathcal{P}', t') - R|$ is small for refinements of a fixed partition, and not for all partitions. Further, note that if $\mathcal{P} \subset \mathcal{P}^*$, then

- $\|\mathcal{P}^*\| \leq \|\mathcal{P}\|$

A weaker definition

Definition (RS Def II)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable if there exists some $R \in \mathbb{R}$ such that for every $\epsilon > 0$ there exists a $\delta > 0$ **and** a partition \mathcal{P} such that **for every tagged refinement** (\mathcal{P}', t') of \mathcal{P} with $\|\mathcal{P}'\| \leq \delta$ we have,

$$|R(f, \mathcal{P}', t') - R| < \epsilon.$$

The nice thing about the above definition is that one only has to check that $|R(f, \mathcal{P}', t') - R|$ is small for refinements of a fixed partition, and not for all partitions. Further, note that if $\mathcal{P} \subset \mathcal{P}^*$, then

- $\|\mathcal{P}^*\| \leq \|\mathcal{P}\|$
- $L(f, \mathcal{P}) \leq L(f, \mathcal{P}^*) \leq R(f, \mathcal{P}^*, t^*) \leq U(f, \mathcal{P}^*) \leq U(f, \mathcal{P})$

Theorem (Tying things up)

The three definitions, viz. The RI, RS Def I, and the RS Def II are **equivalent**.

Theorem (Tying things up)

The three definitions, viz. The RI, RS Def I, and the RS Def II are **equivalent**.

The RI definition, along with the RC help us in proving things about integrals. The RS Def (II) helps us in computing integrals (rigorously).

Application: Area




Figure: Area: Type 1

Application: Area




Figure: Area: Type 1

We compute the area as,

$$A = \int_a^b (f_2(x) - f_1(x)) dx$$

Application: Area




Figure: Area: Type 2

Application: Area




Figure: Area: Type 2

We compute the area as,

$$A = \int_c^d (g_2(y) - g_1(y)) dy$$

Application: Area




Figure: Area: Type 3, $\rho_1(\theta) = 2 \cos \theta$, $\rho_2(\theta) = \cos^2 \theta + \sin \theta$

Application: Area




Figure: Area: Type 3, $\rho_1(\theta) = 2 \cos \theta$, $\rho_2(\theta) = \cos^2 \theta + \sin \theta$

We compute the area as,

$$A = \frac{1}{2} \int_{\alpha}^{\beta} (\rho_2(\theta)^2 - \rho_1(\theta)^2) d\theta$$

Application: Arc Length

Definition (Curve)

A *curve* is a tuple $C = (x(t), y(t))$ wherein $x, y : [a, b] \rightarrow \mathbb{R}$ are continuous functions.

Application: Arc Length

Definition (Curve)

A *curve* is a tuple $C = (x(t), y(t))$ wherein $x, y : [a, b] \rightarrow \mathbb{R}$ are continuous functions.

That is, $C : \mathbb{R} \rightarrow \mathbb{R}^2$. We will assume that C is smooth. The image of a curve is called a *path*.

Application: Arc Length

Definition (Curve)

A *curve* is a tuple $C = (x(t), y(t))$ wherein $x, y : [a, b] \rightarrow \mathbb{R}$ are continuous functions.

That is, $C : \mathbb{R} \rightarrow \mathbb{R}^2$. We will assume that C is smooth. The image of a curve is called a *path*.

Steps:

Application: Arc Length

Definition (Curve)

A *curve* is a tuple $C = (x(t), y(t))$ wherein $x, y : [a, b] \rightarrow \mathbb{R}$ are continuous functions.

That is, $C : \mathbb{R} \rightarrow \mathbb{R}^2$. We will assume that C is smooth. The image of a curve is called a *path*.

Steps:

- Sum of lengths, $\sum_{i=1}^n \sqrt{(x(t_i) - x(t_{i-1}))^2 + (y(t_i) - y(t_{i-1}))^2}$

Application: Arc Length

Definition (Curve)

A *curve* is a tuple $C = (x(t), y(t))$ wherein $x, y : [a, b] \rightarrow \mathbb{R}$ are continuous functions.

That is, $C : \mathbb{R} \rightarrow \mathbb{R}^2$. We will assume that C is smooth. The image of a curve is called a *path*.

Steps:

- Sum of lengths, $\sum_{i=1}^n \sqrt{(x(t_i) - x(t_{i-1}))^2 + (y(t_i) - y(t_{i-1}))^2}$
- Apply MVT, $\sum_{i=1}^n \sqrt{(x'(s_i))^2 + (y'(u_i))^2} (t_i - t_{i-1});$

Application: Arc Length

Definition (Curve)

A *curve* is a tuple $C = (x(t), y(t))$ wherein $x, y : [a, b] \rightarrow \mathbb{R}$ are continuous functions.

That is, $C : \mathbb{R} \rightarrow \mathbb{R}^2$. We will assume that C is smooth. The image of a curve is called a *path*.

Steps:

- Sum of lengths, $\sum_{i=1}^n \sqrt{(x(t_i) - x(t_{i-1}))^2 + (y(t_i) - y(t_{i-1}))^2}$
- Apply MVT, $\sum_{i=1}^n \sqrt{(x'(s_i))^2 + (y'(u_i))^2} (t_i - t_{i-1})$;
recognize the **Riemann Sum**!

Application: Arc Length

Definition (Curve)

A *curve* is a tuple $C = (x(t), y(t))$ wherein $x, y : [a, b] \rightarrow \mathbb{R}$ are continuous functions.

That is, $C : \mathbb{R} \rightarrow \mathbb{R}^2$. We will assume that C is smooth. The image of a curve is called a *path*.

Steps:

- Sum of lengths, $\sum_{i=1}^n \sqrt{(x(t_i) - x(t_{i-1}))^2 + (y(t_i) - y(t_{i-1}))^2}$
- Apply MVT, $\sum_{i=1}^n \sqrt{(x'(s_i))^2 + (y'(u_i))^2} (t_i - t_{i-1})$;
recognize the **Riemann Sum**!
- Conclude,

$$\text{Arc Length}(C) := \int_{\alpha}^{\beta} \sqrt{x'(t)^2 + y'(t)^2} dt$$

Application: Surface Area

- Identify the area of a frustum, $A_F = \pi \lambda_2 (d_1 + d_2)$




Figure: A frustum

Application: Surface Area

- Identify the area of a frustum, $A_F = \pi \lambda_2 (d_1 + d_2)$




Figure: A frustum

- Use this to form the Riemann Sum, $\pi \sum (\rho(t_{i-1}) + \rho_{t_i}) \lambda_i$




Figure: A general area

Application: Surface Area

- Identify the area of a frustum, $A_F = \pi \lambda_2(d_1 + d_2)$




Figure: A frustum

- Use this to form the Riemann Sum, $\pi \sum (\rho(t_{i-1}) + \rho(t_i)) \lambda_i$




Figure: A general area

Thus,

$$\text{Area}(S) := 2\pi \int_{\alpha}^{\beta} \rho(t) \sqrt{x'(t)^2 + y'(t)^2} dt,$$

Application: Volume

Same procedure – partition, form a Riemann Sum, conclude hoping that the [area function](#) A is integrable.

Application: Volume

Same procedure – partition, form a Riemann Sum, conclude hoping that the [area function](#) A is integrable.

$$\text{Vol}(D) := \int_a^b A(x) dx$$

Application: Volume

Same procedure – partition, form a Riemann Sum, conclude hoping that the [area function](#) A is integrable.

$$\text{Vol}(D) := \int_a^b A(x) dx$$

We will deal with solids of revolution.

Application: Volume

Same procedure – partition, form a Riemann Sum, conclude hoping that the **area function** A is integrable.

$$\text{Vol}(D) := \int_a^b A(x) dx$$

We will deal with solids of revolution. Two methods, funny names,

- 1 Washer,
- 2 Shell.

Washer Method

Slices take **perpendicular to rotation axis**.

Washer Method

Slices take **perpendicular to rotation axis**.




Figure: Washers and Disks

Washer Method

Slices take **perpendicular to rotation axis**.




Figure: Washers and Disks

$$Vol(D) = \int_a^b \pi(f_2(x)^2 - f_1(x)^2) dx$$

Shell Method

Shells take **parallel to rotation axis**.

Shell Method

Shells take **parallel to rotation axis**.




Figure: Shells

Shell Method

Shells take **parallel to rotation axis**.




Figure: Shells

$$Vol(D) = 2\pi \int_a^b x(f_2(x) - f_1(x))dx$$