

MA 106 D1-T3 Recap-1

Siddhant Midha

30-03-2022

Recall

- Linear Systems

$$Ax = b$$

Recall

- Linear Systems

$$Ax = b$$

- Completely described by the *augmented matrix*

$$[A|b]$$

Recall

- Linear Systems

$$Ax = b$$

- Completely described by the *augmented matrix*

$$[A|b]$$

- EROs

Recall

- Linear Systems

$$Ax = b$$

- Completely described by the *augmented matrix*

$$[A|b]$$

- EROs

- Interchange the i^{th} and j^{th} row

$$P_{ij}$$

Recall

- Linear Systems

$$Ax = b$$

- Completely described by the *augmented matrix*

$$[A|b]$$

- EROs

- Interchange the i^{th} and j^{th} row

$$P_{ij}$$

- Multiply the j^{th} row by c and add to the i^{th} row.

$$E_{ij}$$

Recall

- Linear Systems

$$Ax = b$$

- Completely described by the *augmented matrix*

$$[A|b]$$

- EROs

- Interchange the i^{th} and j^{th} row

$$P_{ij}$$

- Multiply the j^{th} row by c and add to the i^{th} row.

$$E_{ij}$$

- Multiply the i^{th} row by $c \neq 0$.

REF

- Motivation?

REF

- Motivation?
- Each row starts with strictly more number of zeroes than the previous.

REF

- Motivation?
- Each row starts with strictly more number of zeroes than the previous.
- *PIVOT!* First non zero entry in the i^{th} row is the i^{th} pivot.

REF

- Motivation?
- Each row starts with strictly more number of zeroes than the previous.
- *PIVOT!* First non zero entry in the i^{th} row is the i^{th} pivot.
- No. of pivots $\leq \min(\text{no. of rows, no. of columns})$

REF

- Motivation?
- Each row starts with strictly more number of zeroes than the previous.
- *PIVOT!* First non zero entry in the i^{th} row is the i^{th} pivot.
- No. of pivots $\leq \min(\text{no. of rows, no. of columns})$
- Not unique.

REF

- Motivation?
- Each row starts with strictly more number of zeroes than the previous.
- *PIVOT!* First non zero entry in the i^{th} row is the i^{th} pivot.
- No. of pivots $\leq \min(\text{no. of rows, no. of columns})$
- Not unique.
- Can be *reduced*.

RREF

Theorem

Let A be a $n \times n$ square matrix. There exist ERM's E_i such that

$$\prod_i E_i A$$

is either the $n \times n$ identity matrix or has the last row zero.

RREF

Theorem

Let A be a $n \times n$ square matrix. There exist ERM's E_i such that

$$\prod_i E_i A$$

is either the $n \times n$ identity matrix or has the last row zero.

Definition

Let $A \in \mathbb{R}^{n \times n}$. It is said to be invertible if there exists $B \in \mathbb{R}^{n \times n}$ such that

$$AB = BA = I$$

where I is the $n \times n$ identity matrix. Such a B is unique and is denoted as A^{-1}

Gauss Jordan

Theorem

A square matrix A is invertible **iff** it is a product of ERM's.

Vector Subspaces

Definition

A subset V of \mathbb{R}^n is called a vector subspace of \mathbb{R}^n if

- 1 It is non empty.

Vector Subspaces

Definition

A subset V of \mathbb{R}^n is called a vector subspace of \mathbb{R}^n if

- 1 It is non empty.
- 2 It is closed under scalar multiplication.

Vector Subspaces

Definition

A subset V of \mathbb{R}^n is called a vector subspace of \mathbb{R}^n if

- 1 It is non empty.
- 2 It is closed under scalar multiplication.
- 3 It is closed under addition.

Vector Subspaces

Definition

A subset V of \mathbb{R}^n is called a vector subspace of \mathbb{R}^n if

- 1 It is non empty.
- 2 It is closed under scalar multiplication.
- 3 It is closed under addition.

Questions:

- Is \mathbb{Q}^2 a subspace of \mathbb{R}^2 ?
- Is \mathbb{R}^2 a subspace of \mathbb{R}^3 ?

To do: Categorize the subspaces of \mathbb{R}^2 and \mathbb{R}^3

Linear Combinations etc.

- $c_1 v_1 + c_2 v_2 + \dots$

Linear Combinations etc.

- $c_1 v_1 + c_2 v_2 + \dots$



$$V := \{v_1, v_2 \dots v_n\}$$

$$\text{Span}(V) := \left\{ \sum_i c_i v_i \mid c_i \in \mathbb{R} \forall i \right\}$$

Linear Combinations etc.

- $c_1 v_1 + c_2 v_2 + \dots$



$$V := \{v_1, v_2 \dots v_n\}$$

$$\text{Span}(V) := \left\{ \sum_i c_i v_i \mid c_i \in \mathbb{R} \forall i \right\}$$

- Show that $\text{Span}(V)$ is a vector subspace.

Linear Combinations etc.

- $c_1 v_1 + c_2 v_2 + \dots$



$$V := \{v_1, v_2 \dots v_n\}$$

$$\text{Span}(V) := \left\{ \sum_i c_i v_i \mid c_i \in \mathbb{R} \forall i \right\}$$

- Show that $\text{Span}(V)$ is a vector subspace.
- Given V as above, and w (all in \mathbb{R}^n), how will you decide if $w \in \text{Span}(V)$?

Linear Combinations etc.

- $c_1 v_1 + c_2 v_2 + \dots$



$$V := \{v_1, v_2 \dots v_n\}$$

$$\text{Span}(V) := \left\{ \sum_i c_i v_i \mid c_i \in \mathbb{R} \forall i \right\}$$

- Show that $\text{Span}(V)$ is a vector subspace.
- Given V as above, and w (all in \mathbb{R}^n), how will you decide if $w \in \text{Span}(V)$?
- Linear dependence: There exist a_i *not all* zero such that

$$\sum_i a_i v_i = 0$$

Linear Combinations etc.

- $c_1 v_1 + c_2 v_2 + \dots$



$$V := \{v_1, v_2 \dots v_n\}$$

$$\text{Span}(V) := \left\{ \sum_i c_i v_i \mid c_i \in \mathbb{R} \forall i \right\}$$

- Show that $\text{Span}(V)$ is a vector subspace.
- Given V as above, and w (all in \mathbb{R}^n), how will you decide if $w \in \text{Span}(V)$?
- Linear dependence: There exist a_i *not all* zero such that

$$\sum_i a_i v_i = 0$$

- Redundancy?

Linear Combinations etc.

- $c_1 v_1 + c_2 v_2 + \dots$



$$V := \{v_1, v_2 \dots v_n\}$$

$$\text{Span}(V) := \left\{ \sum_i c_i v_i \mid c_i \in \mathbb{R} \forall i \right\}$$

- Show that $\text{Span}(V)$ is a vector subspace.
- Given V as above, and w (all in \mathbb{R}^n), how will you decide if $w \in \text{Span}(V)$?
- Linear dependence: There exist a_i *not all* zero such that

$$\sum_i a_i v_i = 0$$

- Redundancy?
- Linear Independence

Fundamental Lemma

Lemma

Let V be a vector subspace of \mathbb{R}^n generated by k vectors. Any set of l vectors such that $l \geq k + 1$ is linearly dependent.

Basis and Dimension

- The question of optimality of generation.

Basis and Dimension

- The question of optimality of generation.
- Conventions:
 - \emptyset is linearly independent.
 - $\text{Span}(\emptyset) = \{0\}$

Basis and Dimension

- The question of optimality of generation.
- Conventions:
 - \emptyset is linearly independent.
 - $\text{Span}(\emptyset) = \{0\}$

Definition

Let V be a vector subspace of \mathbb{R}^n . A subset S of V is called a basis of V if

- S is linearly independent.
- $\text{Span}(S) = V$

Theorem

Let $V \subseteq \mathbb{R}^n$ be a vector subspace of \mathbb{R}^n .

- 1 V has a basis.
- 2 Each basis of V has the same number of elements.

Theorem

Let $V \subseteq \mathbb{R}^n$ be a vector subspace of \mathbb{R}^n .

- 1 V has a basis.
- 2 Each basis of V has the same number of elements.

With that in mind, we have the following definition.

Theorem

Let $V \subseteq \mathbb{R}^n$ be a vector subspace of \mathbb{R}^n .

- 1 V has a basis.
- 2 Each basis of V has the same number of elements.

With that in mind, we have the following definition.

Definition

Let V be a vector subspace of \mathbb{R}^n . Then, the dimension of V , denoted $\text{Dim}(V)$, is defined as the cardinality of any basis of V .

Theorem

Let $V \subseteq \mathbb{R}^n$ be a vector subspace of \mathbb{R}^n .

- 1 V has a basis.
- 2 Each basis of V has the same number of elements.

With that in mind, we have the following definition.

Definition

Let V be a vector subspace of \mathbb{R}^n . Then, the dimension of V , denoted $\text{Dim}(V)$, is defined as the cardinality of any basis of V .

(Revisiting the) Question: Show that $v_1, v_2, v_3 \in \mathbb{R}^3$ being linearly independent is equivalent to $\det\begin{bmatrix} v_1 & v_2 & v_3 \end{bmatrix} \neq 0$

Rank and all that

Definitions

Let $A \in \mathbb{R}^{m \times n}$. In terms of its columns and rows, A can be written as

$$\begin{bmatrix} c_1 & c_2 & \dots & c_n \end{bmatrix} \text{ or } \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix}$$

Rank and all that

Definitions

Let $A \in \mathbb{R}^{m \times n}$. In terms of its columns and rows, A can be written as

$$\begin{bmatrix} c_1 & c_2 & \dots & c_n \end{bmatrix} \text{ or } \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix}$$

- Column Space $:= \text{Span}(\{c_i\})$

Rank and all that

Definitions

Let $A \in \mathbb{R}^{m \times n}$. In terms of its columns and rows, A can be written as

$$\begin{bmatrix} c_1 & c_2 & \dots & c_n \end{bmatrix} \text{ or } \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix}$$

- Column Space $:= \text{Span}(\{c_i\})$
- Row Space $:= \text{Span}(\{r_i\})$

Rank and all that

Definitions

Let $A \in \mathbb{R}^{m \times n}$. In terms of its columns and rows, A can be written as

$$\begin{bmatrix} c_1 & c_2 & \dots & c_n \end{bmatrix} \text{ or } \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix}$$

- Column Space $:= \text{Span}(\{c_i\})$
- Row Space $:= \text{Span}(\{r_i\})$
- Column Rank $:= \text{Dim}(\text{Column Space})$

Rank and all that

Definitions

Let $A \in \mathbb{R}^{m \times n}$. In terms of its columns and rows, A can be written as

$$\begin{bmatrix} c_1 & c_2 & \dots & c_n \end{bmatrix} \text{ or } \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix}$$

- Column Space $:= \text{Span}(\{c_i\})$
- Row Space $:= \text{Span}(\{r_i\})$
- Column Rank $:= \text{Dim}(\text{Column Space})$
- Row Rank $:= \text{Dim}(\text{Row Space})$

Rank and all that

- Row rank and row space are preserved under EROs.

Rank and all that

- Row rank and row space are preserved under EROs.
- Column rank is preserved under EROs.

Rank and all that

- Row rank and row space are preserved under EROs.
- Column rank is preserved under EROs.
- Is the column *space* preserved?

Rank and all that

- Row rank and row space are preserved under EROs.
- Column rank is preserved under EROs.
- Is the column *space* preserved?
- Perform EROs to transform to RREF.

Rank and all that

- Row rank and row space are preserved under EROs.
- Column rank is preserved under EROs.
- Is the column *space* preserved?
- Perform EROs to transform to RREF.
- No. of pivots = Column Rank

Rank and all that

- Row rank and row space are preserved under EROs.
- Column rank is preserved under EROs.
- Is the column *space* preserved?
- Perform EROs to transform to RREF.
- No. of pivots = Column Rank
- No. of pivots = Row Rank

Rank and all that

- Row rank and row space are preserved under EROs.
- Column rank is preserved under EROs.
- Is the column *space* preserved?
- Perform EROs to transform to RREF.
- No. of pivots = Column Rank
- No. of pivots = Row Rank
- The *rank* of a matrix is the number of pivots in its RREF!

Rank and all that

- Row rank and row space are preserved under EROs.
- Column rank is preserved under EROs.
- Is the column *space* preserved?
- Perform EROs to transform to RREF.
- No. of pivots = Column Rank
- No. of pivots = Row Rank
- The *rank* of a matrix is the number of pivots in its RREF!

Kronecker-Capelli Theorem

Theorem

A system of linear equations

$$Ax = b$$

has a solution iff

$$\text{Rank}(A) = \text{Rank}([A|b])$$

Rank & Nullity

Definition

Given $A \in \mathbb{R}^{m \times n}$.

$$\text{Null Space}(A) := \{v \in \mathbb{R}^n \mid Av = 0\}$$

Rank & Nullity

Definition

Given $A \in \mathbb{R}^{m \times n}$.

$$\text{Null Space}(A) := \{v \in \mathbb{R}^n \mid Av = 0\}$$

$$N(A) = \text{Dim}(\text{Null Space}(A))$$

Rank & Nullity

Definition

Given $A \in \mathbb{R}^{m \times n}$.

$$\text{Null Space}(A) := \{v \in \mathbb{R}^n \mid Av = 0\}$$

$$N(A) = \text{Dim}(\text{Null Space}(A))$$

Rank Nullity Theorem

Given $A \in \mathbb{R}^{m \times n}$.

$$R(A) + N(A) = n$$

Determinantal Rank

Lemma

Let $A \in \mathbb{R}^{n \times n}$. A $k \times k$ submatrix of A has non zero determinant iff $\text{Rank}(A) \geq k$.

Determinantal Rank

Lemma

Let $A \in \mathbb{R}^{n \times n}$. A $k \times k$ submatrix of A has non zero determinant iff $\text{Rank}(A) \geq k$.

Definition

$A \in \mathbb{R}^{m \times n}$ is said to have determinantal rank k (denoted $\text{det rank}(A)$) if

Determinantal Rank

Lemma

Let $A \in \mathbb{R}^{n \times n}$. A $k \times k$ submatrix of A has non zero determinant iff $\text{Rank}(A) \geq k$.

Definition

$A \in \mathbb{R}^{m \times n}$ is said to have determinantal rank k (denoted $\text{det rank}(A)$) if

- 1 There exists some $k \times k$ submatrix of A with non zero determinant.

Determinantal Rank

Lemma

Let $A \in \mathbb{R}^{n \times n}$. A $k \times k$ submatrix of A has non zero determinant iff $\text{Rank}(A) \geq k$.

Definition

$A \in \mathbb{R}^{m \times n}$ is said to have determinantal rank k (denoted $\text{det rank}(A)$) if

- 1 There exists some $k \times k$ submatrix of A with non zero determinant.
- 2 All $(k + 1) \times (k + 1)$ submatrices of A have zero determinant.

Determinantal Rank

Lemma

Let $A \in \mathbb{R}^{n \times n}$. A $k \times k$ submatrix of A has non zero determinant iff $\text{Rank}(A) \geq k$.

Definition

$A \in \mathbb{R}^{m \times n}$ is said to have determinantal rank k (denoted $\text{det rank}(A)$) if

- 1 There exists some $k \times k$ submatrix of A with non zero determinant.
- 2 All $(k + 1) \times (k + 1)$ submatrices of A have zero determinant.

Fact:

$$\text{det rank}(A) = \text{rank}(A)$$