

# MA 109 D2 T1

## Week Five Recap

Siddhant Midha

<https://siddhant-midha.github.io/>

December 13, 2022

$$\mathbb{R} \rightarrow \mathbb{R}^2$$

$$\mathbb{R} \rightarrow \mathbb{R}^2$$

- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$

- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$
- Note,

$$|x_n - x_0|, |y_n - y_0| \leq ||(x_n, y_n) - (x_0, y_0)|| \leq |x_n - x_0| + |y_n - y_0|$$

- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$
- Note,

$$|x_n - x_0|, |y_n - y_0| \leq \|(x_n, y_n) - (x_0, y_0)\| \leq |x_n - x_0| + |y_n - y_0|$$

- That is,

$$(x_n, y_n) \rightarrow (x_0, y_0) \Leftrightarrow x_n \rightarrow x_0 \text{ and } y_n \rightarrow y_0$$

- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$
- Note,

$$|x_n - x_0|, |y_n - y_0| \leq ||(x_n, y_n) - (x_0, y_0)|| \leq |x_n - x_0| + |y_n - y_0|$$

- That is,

$$(x_n, y_n) \rightarrow (x_0, y_0) \Leftrightarrow x_n \rightarrow x_0 \text{ and } y_n \rightarrow y_0$$

- **Functions:** We will deal with  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ .

- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$
- Note,

$$|x_n - x_0|, |y_n - y_0| \leq ||(x_n, y_n) - (x_0, y_0)|| \leq |x_n - x_0| + |y_n - y_0|$$

- That is,

$$(x_n, y_n) \rightarrow (x_0, y_0) \Leftrightarrow x_n \rightarrow x_0 \text{ and } y_n \rightarrow y_0$$

- **Functions:** We will deal with  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ .
  - ①  $\{(x, y, f(x, y)) \mid (x, y) \in D\}$  is the **graph** of  $f$

- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$
- Note,

$$|x_n - x_0|, |y_n - y_0| \leq ||(x_n, y_n) - (x_0, y_0)|| \leq |x_n - x_0| + |y_n - y_0|$$

- That is,

$$(x_n, y_n) \rightarrow (x_0, y_0) \Leftrightarrow x_n \rightarrow x_0 \text{ and } y_n \rightarrow y_0$$

- **Functions:** We will deal with  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ .
  - 1  $\{(x, y, f(x, y)) \mid (x, y) \in D\}$  is the **graph** of  $f$
  - 2 For some fixed  $c \in \mathbb{R}$ ,  $\{(x, y, c) \mid (x, y) \in D \text{ and } f(x, y) = c\}$  is a **contour line**.



- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$
- Note,

$$|x_n - x_0|, |y_n - y_0| \leq ||(x_n, y_n) - (x_0, y_0)|| \leq |x_n - x_0| + |y_n - y_0|$$

- That is,

$$(x_n, y_n) \rightarrow (x_0, y_0) \Leftrightarrow x_n \rightarrow x_0 \text{ and } y_n \rightarrow y_0$$

- **Functions:** We will deal with  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ .
  - 1  $\{(x, y, f(x, y)) \mid (x, y) \in D\}$  is the **graph** of  $f$
  - 2 For some fixed  $c \in \mathbb{R}$ ,  $\{(x, y, c) \mid (x, y) \in D \text{ and } f(x, y) = c\}$  is a **contour line**.
  - 3 For some fixed  $c \in \mathbb{R}$ ,  $\{(x, y) \mid (x, y) \in D \text{ and } f(x, y) = c\}$  is a **level curve**.

- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$
- Note,

$$|x_n - x_0|, |y_n - y_0| \leq ||(x_n, y_n) - (x_0, y_0)|| \leq |x_n - x_0| + |y_n - y_0|$$

- That is,

$$(x_n, y_n) \rightarrow (x_0, y_0) \Leftrightarrow x_n \rightarrow x_0 \text{ and } y_n \rightarrow y_0$$

- **Functions:** We will deal with  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ .
  - 1  $\{(x, y, f(x, y)) \mid (x, y) \in D\}$  is the **graph** of  $f$
  - 2 For some fixed  $c \in \mathbb{R}$ ,  $\{(x, y, c) \mid (x, y) \in D \text{ and } f(x, y) = c\}$  is a **contour line**.
  - 3 For some fixed  $c \in \mathbb{R}$ ,  $\{(x, y) \mid (x, y) \in D \text{ and } f(x, y) = c\}$  is a **level curve**.
  - 4 **Limits and Continuity:** Again, two definitions for both. Again equivalent. Similar relations between limits and continuity.

- **Sequences:** A sequence in  $\mathbb{R}^2$  is of the form  $((x_n, y_n))$
- Note,

$$|x_n - x_0|, |y_n - y_0| \leq ||(x_n, y_n) - (x_0, y_0)|| \leq |x_n - x_0| + |y_n - y_0|$$

- That is,

$$(x_n, y_n) \rightarrow (x_0, y_0) \Leftrightarrow x_n \rightarrow x_0 \text{ and } y_n \rightarrow y_0$$

- **Functions:** We will deal with  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ .
  - 1  $\{(x, y, f(x, y)) \mid (x, y) \in D\}$  is the **graph** of  $f$
  - 2 For some fixed  $c \in \mathbb{R}$ ,  $\{(x, y, c) \mid (x, y) \in D \text{ and } f(x, y) = c\}$  is a **contour line**.
  - 3 For some fixed  $c \in \mathbb{R}$ ,  $\{(x, y) \mid (x, y) \in D \text{ and } f(x, y) = c\}$  is a **level curve**.
  - 4 **Limits and Continuity:** Again, two definitions for both. Again equivalent. Similar relations between limits and continuity.
  - 5 But note that, we will only deal with *interior points* here.

# Partial Derivatives

# Partial Derivatives

‘Partial’ Derivatives – Rate of change along the  $x$  and  $y$  axes.

# Partial Derivatives

'Partial' Derivatives – Rate of change along the  $x$  and  $y$  axes.

## Definition (Partial Derivatives)

Let  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , and let  $(x_0, y_0) \in D$  be an interior point of  $D$ .

# Partial Derivatives

'Partial' Derivatives – Rate of change along the  $x$  and  $y$  axes.

## Definition (Partial Derivatives)

Let  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , and let  $(x_0, y_0) \in D$  be an interior point of  $D$ . Then, we say that,

# Partial Derivatives

'Partial' Derivatives – Rate of change along the  $x$  and  $y$  axes.

## Definition (Partial Derivatives)

Let  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , and let  $(x_0, y_0) \in D$  be an interior point of  $D$ . Then, we say that,

- 1  $f$  has a partial derivative w.r.t.  $x$  at  $(x_0, y_0)$  if

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h, y_0) - f(x_0, y_0)}{h}$$

exists.



# Partial Derivatives

'Partial' Derivatives – Rate of change along the  $x$  and  $y$  axes.

## Definition (Partial Derivatives)

Let  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , and let  $(x_0, y_0) \in D$  be an interior point of  $D$ . Then, we say that,

- 1  $f$  has a partial derivative w.r.t.  $x$  at  $(x_0, y_0)$  if

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h, y_0) - f(x_0, y_0)}{h}$$

exists. We denote it as  $f_x(x_0, y_0)$ .

# Partial Derivatives

'Partial' Derivatives – Rate of change along the  $x$  and  $y$  axes.

## Definition (Partial Derivatives)

Let  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , and let  $(x_0, y_0) \in D$  be an interior point of  $D$ . Then, we say that,

- ①  $f$  has a partial derivative w.r.t.  $x$  at  $(x_0, y_0)$  if

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h, y_0) - f(x_0, y_0)}{h}$$

exists. We denote it as  $f_x(x_0, y_0)$ .

- ②  $f$  has a partial derivative w.r.t.  $y$  at  $(x_0, y_0)$  if

$$\lim_{h \rightarrow 0} \frac{f(x_0, y_0 + h) - f(x_0, y_0)}{h}$$

exists. We denote it as  $f_y(x_0, y_0)$ .

# Gradient

## Definition (Gradient)

Let  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , and let  $(x_0, y_0) \in D$  be an interior point of  $D$ . Suppose both partial derivatives of  $f$  exist at  $(x_0, y_0)$ , then we define the gradient of  $f$  at  $(x_0, y_0)$

$$\nabla f(x_0, y_0) := (f_x(x_0, y_0), f_y(x_0, y_0))^T \in \mathbb{R}^2$$

## Definition (Gradient)

Let  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , and let  $(x_0, y_0) \in D$  be an interior point of  $D$ . Suppose both partial derivatives of  $f$  exist at  $(x_0, y_0)$ , then we define the gradient of  $f$  at  $(x_0, y_0)$

$$\nabla f(x_0, y_0) := (f_x(x_0, y_0), f_y(x_0, y_0))^T \in \mathbb{R}^2$$

## Definition (Directional Derivatives)

Let  $f : D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ , and let  $(x_0, y_0) \in D$  be an interior point of  $D$ . For some  $u = (u_1, u_2)^T \in \mathbb{R}^2$ , we say that  $f$  has a directional derivative **along  $u$**  at  $(x_0, y_0)$  if

$$\lim_{t \rightarrow 0} \frac{f(x_0 + tu_1, y_0 + tu_2) - f(x_0, y_0)}{t}$$

exists, and denote it by  $D_u f(x_0, y_0)$ .

# 'Differentiability' ?

In the 1D case, we saw that, Differentiability implies Continuity.

# 'Differentiability' ?

In the 1D case, we saw that, Differentiability implies Continuity. Well, should we take the definitions of directional/partial derivatives as the definition of 2D Differentiability?

# 'Differentiability' ?

In the 1D case, we saw that, Differentiability implies Continuity. Well, should we take the definitions of directional/partial derivatives as the definition of 2D Differentiability?

Consider,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f(x, y) := 0 \forall (x, y) : x \neq 0 \text{ and } y \neq 0$ ,  
 $f(x, 0) := 1000 \forall x$  and  $f(0, y) := 1000 \forall y$ .



# 'Differentiability' ?

In the 1D case, we saw that, Differentiability implies Continuity. Well, should we take the definitions of directional/partial derivatives as the definition of 2D Differentiability?

Consider,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f(x, y) := 0 \forall (x, y) : x \neq 0 \text{ and } y \neq 0$ ,  $f(x, 0) := 1000 \forall x$  and  $f(0, y) := 1000 \forall y$ .

Does the 'derivative' exist at  $(0, 0)$ ? Continuity? Well?

# 'Differentiability' ?

In the 1D case, we saw that, Differentiability implies Continuity. Well, should we take the definitions of directional/partial derivatives as the definition of 2D Differentiability?

Consider,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f(x, y) := 0 \forall (x, y) : x \neq 0 \text{ and } y \neq 0$ ,  $f(x, 0) := 1000 \forall x$  and  $f(0, y) := 1000 \forall y$ .

Does the 'derivative' exist at  $(0, 0)$ ? Continuity? Well?

Thus, the directional derivative is not a *strong enough* definition! Can have functions for which DD's exist at a point, but is not continuous at that point.

# 'Differentiability' ?

In the 1D case, we saw that, Differentiability implies Continuity. Well, should we take the definitions of directional/partial derivatives as the definition of 2D Differentiability?

Consider,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f(x, y) := 0 \forall (x, y) : x \neq 0 \text{ and } y \neq 0$ ,  $f(x, 0) := 1000 \forall x$  and  $f(0, y) := 1000 \forall y$ .

Does the 'derivative' exist at  $(0, 0)$ ? Continuity? Well?

Thus, the directional derivative is not a *strong enough* definition! Can have functions for which DD's exist at a point, but is not continuous at that point.

## Tangent Lines

# 'Differentiability' ?

In the 1D case, we saw that, Differentiability implies Continuity. Well, should we take the definitions of directional/partial derivatives as the definition of 2D Differentiability?

Consider,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f(x, y) := 0 \forall (x, y) : x \neq 0 \text{ and } y \neq 0$ ,  $f(x, 0) := 1000 \forall x$  and  $f(0, y) := 1000 \forall y$ .

Does the 'derivative' exist at  $(0, 0)$ ? Continuity? Well?

Thus, the directional derivative is not a *strong enough* definition! Can have functions for which DD's exist at a point, but is not continuous at that point.

## Tangent Lines

- 1D:  $y = f(x_0) + f'(x_0)(x - x_0)$

# 'Differentiability' ?

In the 1D case, we saw that, Differentiability implies Continuity. Well, should we take the definitions of directional/partial derivatives as the definition of 2D Differentiability?

Consider,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f(x, y) := 0 \forall (x, y) : x \neq 0 \text{ and } y \neq 0$ ,  $f(x, 0) := 1000 \forall x$  and  $f(0, y) := 1000 \forall y$ .

Does the 'derivative' exist at  $(0, 0)$ ? Continuity? Well?

Thus, the directional derivative is not a *strong enough* definition! Can have functions for which DD's exist at a point, but is not continuous at that point.

## Tangent Lines

- 1D:  $y = f(x_0) + f'(x_0)(x - x_0)$
- 2D  $z = f(x_0, y_0) + \partial_x(x_0, y_0)(x - x_0) + \partial_y(x_0, y_0)(y - y_0)$

# 'Differentiability'?

In the 1D case, we saw that, Differentiability implies Continuity. Well, should we take the definitions of directional/partial derivatives as the definition of 2D Differentiability?

Consider,  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  such that  $f(x, y) := 0 \forall (x, y) : x \neq 0 \text{ and } y \neq 0$ ,  $f(x, 0) := 1000 \forall x$  and  $f(0, y) := 1000 \forall y$ .

Does the 'derivative' exist at  $(0, 0)$ ? Continuity? Well?

Thus, the directional derivative is not a *strong enough* definition! Can have functions for which DD's exist at a point, but is not continuous at that point.

## Tangent Lines

- 1D:  $y = f(x_0) + f'(x_0)(x - x_0)$
- 2D  $z = f(x_0, y_0) + \partial_x(x_0, y_0)(x - x_0) + \partial_y(x_0, y_0)(y - y_0)$

1D Definition:

$$\lim_{h \rightarrow 0} \frac{|f(x_0 + h) - f(x_0) - f'(x_0)h|}{|h|} = 0$$

What about 2D?

## Definition (Differentiability)

A function  $f : D \rightarrow \mathbb{R}$  is said to be differentiable at an interior point  $(x_0, y_0)$  if,  $\partial_x f$  and  $\partial_y f$  exist at that point and,

$$\lim_{(h,k) \rightarrow 0} \frac{|f(x_0 + h, y_0 + k) - f(x_0, y_0) - \partial_x(x_0, y_0)h - \partial_y(x_0, y_0)k|}{\|(h, k)\|} = 0$$

## Definition (Differentiability)

A function  $f : D \rightarrow \mathbb{R}$  is said to be differentiable at an interior point  $(x_0, y_0)$  if,  $\partial_x f$  and  $\partial_y f$  exist at that point and,

$$\lim_{(h,k) \rightarrow 0} \frac{|f(x_0 + h, y_0 + k) - f(x_0, y_0) - \partial_x(x_0, y_0)h - \partial_y(x_0, y_0)k|}{\|(h, k)\|} = 0$$

Differentiability at a point *implies* the following,



## Definition (Differentiability)

A function  $f : D \rightarrow \mathbb{R}$  is said to be differentiable at an interior point  $(x_0, y_0)$  if,  $\partial_x f$  and  $\partial_y f$  exist at that point and,

$$\lim_{(h,k) \rightarrow 0} \frac{|f(x_0 + h, y_0 + k) - f(x_0, y_0) - \partial_x(x_0, y_0)h - \partial_y(x_0, y_0)k|}{\|(h, k)\|} = 0$$

Differentiability at a point *implies* the following,

- Existence of partial derivatives at that point.

## Definition (Differentiability)

A function  $f : D \rightarrow \mathbb{R}$  is said to be differentiable at an interior point  $(x_0, y_0)$  if,  $\partial_x f$  and  $\partial_y f$  exist at that point and,

$$\lim_{(h,k) \rightarrow 0} \frac{|f(x_0 + h, y_0 + k) - f(x_0, y_0) - \partial_x(x_0, y_0)h - \partial_y(x_0, y_0)k|}{\|(h, k)\|} = 0$$

Differentiability at a point *implies* the following,

- Existence of partial derivatives at that point.
- Existence of ALL directional derivatives at that point.

## Definition (Differentiability)

A function  $f : D \rightarrow \mathbb{R}$  is said to be differentiable at an interior point  $(x_0, y_0)$  if,  $\partial_x f$  and  $\partial_y f$  exist at that point and,

$$\lim_{(h,k) \rightarrow 0} \frac{|f(x_0 + h, y_0 + k) - f(x_0, y_0) - \partial_x(x_0, y_0)h - \partial_y(x_0, y_0)k|}{\|(h, k)\|} = 0$$

Differentiability at a point *implies* the following,

- Existence of partial derivatives at that point.
- Existence of ALL directional derivatives at that point.
- The fact that  $D_u f(x_0, y_0) = \nabla f(x_0, y_0) \cdot u$  for every unit vector  $u$ .

## Definition (Differentiability)

A function  $f : D \rightarrow \mathbb{R}$  is said to be differentiable at an interior point  $(x_0, y_0)$  if,  $\partial_x f$  and  $\partial_y f$  exist at that point and,

$$\lim_{(h,k) \rightarrow 0} \frac{|f(x_0 + h, y_0 + k) - f(x_0, y_0) - \partial_x(x_0, y_0)h - \partial_y(x_0, y_0)k|}{\|(h, k)\|} = 0$$

Differentiability at a point *implies* the following,

- Existence of partial derivatives at that point.
- Existence of ALL directional derivatives at that point.
- The fact that  $D_u f(x_0, y_0) = \nabla f(x_0, y_0) \cdot u$  for every unit vector  $u$ .
- Continuity of  $f$  at that point.

# Alternate Condition

## Proposition

Let  $f : D \rightarrow \mathbb{R}$  and let  $(x_0, y_0)$  be an interior point of  $D$ . If the partial derivatives of  $f$  exist in some neighbourhood of  $(x_0, y_0)$  and are continuous at  $(x_0, y_0)$  then  $f$  is differentiable at  $(x_0, y_0)$ .

# Example

$$f(x, y) := xy \frac{x^2 - y^2}{x^2 + y^2} \forall (x, y) \neq (0, 0); \quad f(0, 0) := 0$$

# Example

$$f(x, y) := xy \frac{x^2 - y^2}{x^2 + y^2} \forall (x, y) \neq (0, 0); \quad f(0, 0) := 0$$




Figure: The graph of  $f$